



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 01:49 AM GMT

PDB ID : 2EFE
Title : Ara7-GDPNH2/AtVps9a
Authors : Ihara, K.; Uejima, T.; Wakatsuki, S.
Deposited on : 2007-02-22
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4546 (2.10-2.06)
Clashscore	102246	5101 (2.10-2.06)
Ramachandran outliers	100387	5048 (2.10-2.06)
Sidechain outliers	100360	5049 (2.10-2.06)
RSRZ outliers	91569	4556 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Similarity to vacuolar protein sorting-associated protein VPS9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	0	0	0
			1953	1242	324	375	12			

Mol	Chain	Residues	Total	C	N	O	S	ZeroOcc	AltConf	Trace
1	C	246	Total	C	N	O	S	0	0	0
			1953	1242	324	375	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP Q9LT31
A	0	SER	-	EXPRESSION TAG	UNP Q9LT31
C	-1	GLY	-	EXPRESSION TAG	UNP Q9LT31
C	0	SER	-	EXPRESSION TAG	UNP Q9LT31


- Molecule 2 is a protein called Small GTP-binding protein-like.

Mol	Chain	Residues	Total	C	N	O	S	ZeroOcc	AltConf	Trace
2	B	165	1287	820	218	245	4	0	0	0
2	D	165	1287	820	218	245	4	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	EXPRESSION TAG	UNP Q9LT31
B	0	SER	-	EXPRESSION TAG	UNP Q9LT31
D	-1	GLY	-	EXPRESSION TAG	UNP Q9LT31
D	0	SER	-	EXPRESSION TAG	UNP Q9LT31

- Molecule 3 is AMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNH) (formula: C₁₀H₁₆N₆O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	B	1	28	10	6	10	2	0	0
3	D	1	28	10	6	10	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	117	Total O 117 117		0	0
4	B	40	Total O 40 40		0	0
4	C	123	Total O 123 123		0	0
4	D	46	Total O 46 46		0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Similarity to vacuolar protein sorting-associated protein VPS9




- Molecule 1: Similarity to vacuolar protein sorting-associated protein VPS9



- Molecule 2: Small GTP-binding protein-like



- Molecule 2: Small GTP-binding protein-like





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	58.23Å 58.30Å 69.24Å 81.75° 86.83° 73.79°	Depositor
Resolution (Å)	68.52 – 2.08 46.23 – 2.08	Depositor EDS
% Data completeness (in resolution range)	95.8 (68.52-2.08) 92.1 (46.23-2.08)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) >$ ¹	2.35 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R , R_{free}	0.227 , 0.282 0.226 , 0.279	Depositor DCC
R_{free} test set	2529 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.34$	Xtriage
Outliers	0 of 49628 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6862	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GNH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/1991	0.67	0/2687
1	C	0.64	0/1991	0.69	0/2687
2	B	0.65	0/1310	0.72	0/1771
2	D	0.64	0/1310	0.73	0/1771
All	All	0.63	0/6602	0.70	0/8916

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	8	SER	Peptide

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1953	0	1937	31	0
1	C	1953	0	1937	30	0
2	B	1287	0	1273	24	0
2	D	1287	0	1273	24	0
3	B	28	0	14	0	0
3	D	28	0	14	0	0
4	A	117	0	0	18	0
4	B	40	0	0	7	0
4	C	123	0	0	15	0
4	D	46	0	0	3	0
All	All	6862	0	6448	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:232:SER:HB3	4:A:380:HOH:O	1.21	1.32
1:C:61:PHE:HA	4:C:348:HOH:O	1.46	1.14
1:A:133:GLU:HB2	4:A:312:HOH:O	1.61	1.00
2:D:18:ASP:HB2	4:D:2217:HOH:O	1.62	0.98
1:A:178:ASN:HB2	4:A:325:HOH:O	1.64	0.96
1:C:67:TRP:HD1	4:C:297:HOH:O	1.61	0.84
2:D:34:GLN:HB3	2:D:39:GLN:HB2	1.60	0.82
1:C:248:ASP:HB3	4:C:334:HOH:O	1.81	0.80
1:C:128:PRO:HA	1:C:131:GLN:HG3	1.64	0.78
1:A:70:CYS:SG	4:A:330:HOH:O	2.42	0.77
2:B:18:ASP:HB2	4:B:1205:HOH:O	1.88	0.72
1:C:131:GLN:HG2	4:C:331:HOH:O	1.89	0.70
2:D:115:ASN:HB2	4:D:2239:HOH:O	1.91	0.70
2:B:166:GLU:O	2:B:170:ARG:HG3	1.91	0.69
1:A:37:SER:HB3	4:A:323:HOH:O	1.92	0.69
2:D:14:VAL:HG21	2:D:80:TYR:HA	1.75	0.68
1:C:179:GLU:OE1	1:C:179:GLU:HA	1.95	0.67
1:A:129:THR:HG21	1:A:179:GLU:OE2	1.95	0.67
1:C:201:PRO:HD2	4:C:306:HOH:O	1.94	0.66
2:B:118:MET:HE1	4:B:1217:HOH:O	1.96	0.65
1:A:201:PRO:HD2	4:A:304:HOH:O	1.98	0.63
1:A:143:LEU:HD13	1:A:200:PRO:HG3	1.82	0.61
1:C:17:MET:N	4:C:367:HOH:O	2.32	0.61
1:A:178:ASN:ND2	4:A:331:HOH:O	2.33	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:153:ARG:NH1	1:C:154:ASP:OD1	2.36	0.59
1:A:131:GLN:HG2	4:A:375:HOH:O	2.02	0.58
2:B:118:MET:CE	4:B:1217:HOH:O	2.52	0.58
1:A:70:CYS:CB	4:A:330:HOH:O	2.52	0.57
2:D:52:LEU:HD21	2:D:165:TYR:CD1	2.40	0.57
1:A:108:LEU:HD21	1:A:234:GLU:HG3	1.88	0.56
1:C:67:TRP:CD1	4:C:297:HOH:O	2.47	0.55
2:B:14:VAL:HG21	2:B:80:TYR:HA	1.87	0.55
1:A:84:LYS:HG3	1:A:151:ALA:HB2	1.89	0.54
2:B:100:ARG:NH2	4:B:1225:HOH:O	2.40	0.54
1:A:178:ASN:ND2	4:A:280:HOH:O	2.40	0.54
1:A:60:ALA:O	1:A:64:HIS:HB2	2.07	0.54
1:A:189:PRO:HG2	4:A:295:HOH:O	2.07	0.53
2:B:113:ASN:OD1	2:B:113:ASN:N	2.39	0.53
1:C:60:ALA:O	1:C:64:HIS:HB2	2.09	0.53
2:D:166:GLU:O	2:D:170:ARG:HG3	2.09	0.53
1:C:108:LEU:HD21	1:C:234:GLU:HG3	1.91	0.53
2:D:102:LYS:HG2	2:D:141:TYR:CZ	2.44	0.52
1:C:70:CYS:HB3	4:C:332:HOH:O	2.09	0.52
1:C:144:GLN:O	4:C:387:HOH:O	2.19	0.52
1:C:128:PRO:HA	1:C:131:GLN:CG	2.39	0.52
1:C:70:CYS:CB	4:C:332:HOH:O	2.57	0.51
1:C:80:ASP:OD1	1:C:214:ARG:NH2	2.42	0.51
2:B:83:ALA:O	2:B:113:ASN:ND2	2.44	0.51
1:C:107:LYS:NZ	4:C:329:HOH:O	2.43	0.51
1:A:73:GLU:HG3	4:A:317:HOH:O	2.11	0.51
2:B:106:GLN:HG2	4:B:1220:HOH:O	2.10	0.51
2:D:131:ARG:NH1	2:D:151:GLU:OE1	2.41	0.51
2:B:126:ASP:OD1	2:B:153:SER:OG	2.24	0.50
2:D:102:LYS:HG2	2:D:141:TYR:CE1	2.46	0.50
2:D:59:VAL:HG11	2:D:168:ALA:HB1	1.94	0.50
2:D:143:GLN:HA	2:D:143:GLN:NE2	2.26	0.50
2:B:29:ARG:HB3	2:B:160:VAL:HG11	1.94	0.50
1:A:109:PHE:CE1	4:A:347:HOH:O	2.54	0.49
2:D:52:LEU:HD21	2:D:165:TYR:HD1	1.78	0.49
1:A:243:LYS:HG3	1:A:243:LYS:O	2.13	0.49
4:A:310:HOH:O	2:D:40:GLU:CG	2.60	0.49
2:B:102:LYS:O	2:B:106:GLN:HG3	2.12	0.48
1:A:109:PHE:HE1	4:A:347:HOH:O	1.93	0.48
2:D:43:ILE:HD13	2:D:43:ILE:H	1.76	0.48
2:B:105:VAL:O	2:B:109:GLN:HG3	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:9:ILE:HB	2:B:59:VAL:HG22	1.96	0.48
2:B:124:LYS:HB3	2:B:127:LEU:HD12	1.95	0.48
1:C:137:LEU:O	1:C:140:GLN:HB2	2.13	0.48
1:C:262:ARG:C	4:C:318:HOH:O	2.52	0.47
2:B:40:GLU:HG3	4:C:293:HOH:O	2.14	0.47
2:B:34:GLN:HB3	2:B:39:GLN:HB2	1.95	0.47
1:A:54:PHE:O	1:A:58:GLU:HB3	2.14	0.47
2:D:148:PHE:HD2	4:D:2216:HOH:O	1.98	0.47
1:A:61:PHE:O	1:A:67:TRP:HB2	2.14	0.47
1:C:170:LEU:HD21	1:C:190:VAL:HG21	1.97	0.46
1:C:58:GLU:HA	4:C:381:HOH:O	2.16	0.46
2:D:143:GLN:HA	2:D:143:GLN:HE21	1.81	0.45
2:B:81:ARG:HA	4:B:1224:HOH:O	2.16	0.45
2:B:100:ARG:NE	4:B:1226:HOH:O	2.38	0.45
2:B:32:LYS:HE2	2:B:43:ILE:HD12	2.00	0.44
1:C:123:ASN:HB3	1:C:245:ILE:HB	1.98	0.44
1:A:109:PHE:CD1	4:A:347:HOH:O	2.71	0.44
2:B:28:LEU:HD22	2:B:45:ALA:HB2	1.99	0.44
1:A:128:PRO:HA	1:A:131:GLN:HG3	1.99	0.44
2:D:12:LYS:HE3	2:D:64:TRP:CD2	2.53	0.44
1:A:123:ASN:HB3	1:A:245:ILE:HB	2.01	0.43
1:C:62:ARG:HG3	1:C:75:LEU:CD1	2.48	0.43
1:A:44:GLU:OE1	1:A:44:GLU:N	2.48	0.43
1:C:203:LEU:O	1:C:207:LEU:HG	2.18	0.43
2:D:29:ARG:HB3	2:D:160:VAL:HG11	2.01	0.43
1:C:121:PRO:HB3	1:C:126:ILE:HB	2.00	0.43
2:B:32:LYS:HE2	2:B:43:ILE:CD1	2.48	0.43
2:B:73:HIS:CE1	2:B:100:ARG:HH21	2.38	0.42
1:A:70:CYS:HB3	4:A:330:HOH:O	2.16	0.42
2:D:31:VAL:HG12	2:D:32:LYS:HG2	2.00	0.42
1:C:70:CYS:SG	4:C:332:HOH:O	2.62	0.42
1:A:138:LEU:O	1:A:142:GLU:HG2	2.19	0.42
2:D:19:VAL:HG11	2:D:69:GLN:HG3	2.01	0.42
2:D:135:ALA:O	2:D:139:GLN:HB3	2.20	0.42
1:C:62:ARG:HG3	1:C:75:LEU:HD13	2.03	0.41
1:C:108:LEU:HD21	1:C:234:GLU:CG	2.50	0.41
2:D:73:HIS:HE1	2:D:100:ARG:NH2	2.19	0.41
1:A:200:PRO:HA	1:A:201:PRO:HD3	1.97	0.40
2:B:29:ARG:O	2:B:33:ASP:HA	2.21	0.40
2:D:51:THR:HG23	2:D:58:THR:HG23	2.04	0.40
1:A:227:PHE:CE2	1:A:231:LEU:HD11	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:84:LYS:HG3	1:C:151:ALA:HB2	2.03	0.40
1:A:17:MET:N	4:A:379:HOH:O	2.53	0.40
2:D:88:ILE:HD13	2:D:105:VAL:HG22	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	244/267 (91%)	236 (97%)	8 (3%)	0	100 100
1	C	244/267 (91%)	241 (99%)	3 (1%)	0	100 100
2	B	163/181 (90%)	160 (98%)	3 (2%)	0	100 100
2	D	163/181 (90%)	157 (96%)	6 (4%)	0	100 100
All	All	814/896 (91%)	794 (98%)	20 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	217/235 (92%)	205 (94%)	12 (6%)	27 22
1	C	217/235 (92%)	206 (95%)	11 (5%)	29 25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	132/143 (92%)	125 (95%)	7 (5%)	28 24
2	D	132/143 (92%)	119 (90%)	13 (10%)	10 6
All	All	698/756 (92%)	655 (94%)	43 (6%)	23 18

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	PHE
1	A	56	LYS
1	A	133	GLU
1	A	138	LEU
1	A	143	LEU
1	A	156	LEU
1	A	171	LEU
1	A	178	ASN
1	A	179	GLU
1	A	243	LYS
1	A	248	ASP
1	A	254	LYS
2	B	8	SER
2	B	28	LEU
2	B	37	GLU
2	B	40	GLU
2	B	41	SER
2	B	43	ILE
2	B	113	ASN
1	C	25	PHE
1	C	36	PHE
1	C	37	SER
1	C	108	LEU
1	C	138	LEU
1	C	143	LEU
1	C	156	LEU
1	C	171	LEU
1	C	179	GLU
1	C	249	GLU
1	C	254	LYS
2	D	8	SER
2	D	19	VAL
2	D	28	LEU
2	D	34	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	36	VAL
2	D	43	ILE
2	D	49	SER
2	D	54	VAL
2	D	56	ASP
2	D	129	ASP
2	D	147	LEU
2	D	151	GLU
2	D	169	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	116	GLN
1	A	178	ASN
1	A	255	ASN
2	B	139	GLN
2	B	143	GLN
1	C	116	GLN
1	C	255	ASN
2	D	39	GLN
2	D	106	GLN
2	D	143	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GNH	B	1200	-	23,30,30	1.64	4 (17%)	27,47,47	1.90	9 (33%)
3	GNH	D	2200	-	23,30,30	1.53	3 (13%)	27,47,47	1.92	10 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GNH	B	1200	-	-	0/9/32/32	0/3/3/3
3	GNH	D	2200	-	-	0/9/32/32	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1200	GNH	PB-O2B	-2.12	1.50	1.56
3	D	2200	GNH	C5-C4	2.73	1.46	1.40
3	B	1200	GNH	C5-C4	3.13	1.47	1.40
3	D	2200	GNH	C6-C5	3.79	1.48	1.41
3	D	2200	GNH	PB-O1B	4.25	1.51	1.46
3	B	1200	GNH	C6-C5	4.36	1.50	1.41
3	B	1200	GNH	PB-O1B	4.54	1.51	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1200	GNH	C6-C5-C4	-4.03	116.08	120.90
3	D	2200	GNH	C6-C5-C4	-3.52	116.69	120.90
3	B	1200	GNH	C5-C6-N1	-3.51	118.78	123.59
3	D	2200	GNH	C5-C6-N1	-3.26	119.14	123.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2200	GNH	C4-C5-N7	-3.21	106.53	109.48
3	D	2200	GNH	C1'-N9-C4	-3.05	122.34	126.94
3	B	1200	GNH	C4-C5-N7	-2.84	106.87	109.48
3	B	1200	GNH	C1'-N9-C4	-2.68	122.90	126.94
3	D	2200	GNH	N3-C2-N1	-2.55	123.56	127.44
3	B	1200	GNH	N3-C2-N1	-2.53	123.59	127.44
3	D	2200	GNH	C2'-C1'-N9	-2.38	110.66	114.29
3	B	1200	GNH	C2'-C1'-N9	-2.13	111.03	114.29
3	B	1200	GNH	O2B-PB-O1B	2.12	115.44	110.07
3	D	2200	GNH	N2-C2-N1	2.55	121.43	117.20
3	D	2200	GNH	O4'-C1'-N9	2.64	113.63	108.10
3	D	2200	GNH	O2B-PB-O1B	2.66	116.83	110.07
3	B	1200	GNH	O4'-C1'-N9	2.85	114.06	108.10
3	D	2200	GNH	C6-N1-C2	3.44	120.71	115.94
3	B	1200	GNH	C6-N1-C2	4.47	122.14	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	246/267 (92%)	0.26	10 (4%) 41 49	19, 33, 58, 71	0
1	C	246/267 (92%)	0.28	10 (4%) 41 49	18, 32, 60, 73	0
2	B	165/181 (91%)	0.53	17 (10%) 9 11	25, 40, 65, 71	0
2	D	165/181 (91%)	0.55	19 (11%) 6 8	23, 41, 65, 71	0
All	All	822/896 (91%)	0.38	56 (6%) 20 27	18, 36, 61, 73	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	165	TYR	7.0
2	B	57	ALA	5.8
2	B	165	TYR	5.6
2	B	56	ASP	5.1
2	D	57	ALA	5.0
1	A	18	ARG	4.7
2	D	54	VAL	4.6
2	D	56	ASP	4.6
1	A	38	ASN	4.2
2	D	52	LEU	4.2
2	B	54	VAL	4.1
1	C	67	TRP	4.1
2	B	53	ALA	4.0
2	B	9	ILE	4.0
2	B	129	ASP	3.6
1	C	18	ARG	3.5
2	B	55	ASN	3.5
2	D	36	VAL	3.4
1	A	63	ALA	3.4
2	D	37	GLU	3.3
2	D	38	PHE	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	33	ILE	3.2
2	B	59	VAL	3.1
1	A	85	TYR	3.0
1	C	36	PHE	3.0
1	A	65	PRO	3.0
2	D	9	ILE	2.9
1	A	67	TRP	2.9
2	B	172	PRO	2.9
2	B	37	GLU	2.9
2	B	112	GLY	2.8
2	D	53	ALA	2.8
1	C	85	TYR	2.7
2	D	115	ASN	2.6
1	C	20	PRO	2.6
2	D	55	ASN	2.6
2	D	8	SER	2.4
2	D	58	THR	2.4
2	D	129	ASP	2.3
1	A	34	VAL	2.3
2	D	51	THR	2.3
1	A	19	LYS	2.3
1	C	23	GLY	2.2
2	D	39	GLN	2.2
1	C	38	ASN	2.2
1	C	40	ALA	2.2
2	B	113	ASN	2.2
2	B	169	ARG	2.2
1	A	36	PHE	2.2
1	C	63	ALA	2.2
2	B	52	LEU	2.2
1	C	33	ILE	2.2
2	B	58	THR	2.2
2	D	42	THR	2.1
2	D	113	ASN	2.1
2	B	111	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GNH	B	1200	28/28	0.96	0.11	-0.74	28,36,37,38	0
3	GNH	D	2200	28/28	0.95	0.10	-0.75	33,34,36,40	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.