



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:21 PM GMT

PDB ID : 4EGV
Title : Crystal structure of a monomeric SCP2-thiolase like protein type 1 (STLP1) from *Mycobacterium smegmatis*
Authors : Janardan, N.; Harijan, R.K.; Wierenga, R.K.; Murthy, M.R.N.
Deposited on : 2012-04-01
Resolution : 2.71 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2439 (2.74-2.70)
Clashscore	102246	2771 (2.74-2.70)
Ramachandran outliers	100387	2726 (2.74-2.70)
Sidechain outliers	100360	2727 (2.74-2.70)
RSRZ outliers	91569	2443 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	<div> <div>69%</div> <div>16%</div> <div>•</div> <div>14%</div> </div>
1	B	520	<div> <div>68%</div> <div>16%</div> <div>•</div> <div>13%</div> </div>
1	C	520	<div> <div>67%</div> <div>18%</div> <div>•</div> <div>13%</div> </div>
1	D	520	<div> <div>65%</div> <div>17%</div> <div>•</div> <div>14%</div> </div>
1	E	520	<div> <div>67%</div> <div>16%</div> <div>•</div> <div>14%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	520	<div><div><div>%</div><div><div></div><div></div><div></div><div></div></div><div>66%17%•14%</div></div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 20206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl-CoA acetyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	448	Total	C	N	O	S	Se	0	0	0
			3327	2086	579	650	3	9			
1	B	452	Total	C	N	O	S	Se	0	3	0
			3376	2115	588	661	3	9			
1	C	453	Total	C	N	O	S	Se	0	0	0
			3373	2116	588	656	3	10			
1	D	445	Total	C	N	O	S	Se	0	0	0
			3299	2066	576	645	3	9			
1	E	447	Total	C	N	O	S	Se	0	2	0
			3336	2090	585	648	3	10			
1	F	449	Total	C	N	O	S	Se	0	0	0
			3335	2090	581	652	3	9			

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MSE	-	EXPRESSION TAG	UNP A0R3M0
A	-11	ARG	-	EXPRESSION TAG	UNP A0R3M0
A	-10	GLY	-	EXPRESSION TAG	UNP A0R3M0
A	-9	SER	-	EXPRESSION TAG	UNP A0R3M0
A	-8	HIS	-	EXPRESSION TAG	UNP A0R3M0
A	-7	HIS	-	EXPRESSION TAG	UNP A0R3M0
A	-6	HIS	-	EXPRESSION TAG	UNP A0R3M0
A	-5	HIS	-	EXPRESSION TAG	UNP A0R3M0
A	-4	HIS	-	EXPRESSION TAG	UNP A0R3M0
A	-3	HIS	-	EXPRESSION TAG	UNP A0R3M0
A	-2	MSE	-	EXPRESSION TAG	UNP A0R3M0
A	-1	ALA	-	EXPRESSION TAG	UNP A0R3M0
A	0	SER	-	EXPRESSION TAG	UNP A0R3M0
B	-12	MSE	-	EXPRESSION TAG	UNP A0R3M0
B	-11	ARG	-	EXPRESSION TAG	UNP A0R3M0
B	-10	GLY	-	EXPRESSION TAG	UNP A0R3M0
B	-9	SER	-	EXPRESSION TAG	UNP A0R3M0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-8	HIS	-	EXPRESSION TAG	UNP A0R3M0
B	-7	HIS	-	EXPRESSION TAG	UNP A0R3M0
B	-6	HIS	-	EXPRESSION TAG	UNP A0R3M0
B	-5	HIS	-	EXPRESSION TAG	UNP A0R3M0
B	-4	HIS	-	EXPRESSION TAG	UNP A0R3M0
B	-3	HIS	-	EXPRESSION TAG	UNP A0R3M0
B	-2	MSE	-	EXPRESSION TAG	UNP A0R3M0
B	-1	ALA	-	EXPRESSION TAG	UNP A0R3M0
B	0	SER	-	EXPRESSION TAG	UNP A0R3M0
C	-12	MSE	-	EXPRESSION TAG	UNP A0R3M0
C	-11	ARG	-	EXPRESSION TAG	UNP A0R3M0
C	-10	GLY	-	EXPRESSION TAG	UNP A0R3M0
C	-9	SER	-	EXPRESSION TAG	UNP A0R3M0
C	-8	HIS	-	EXPRESSION TAG	UNP A0R3M0
C	-7	HIS	-	EXPRESSION TAG	UNP A0R3M0
C	-6	HIS	-	EXPRESSION TAG	UNP A0R3M0
C	-5	HIS	-	EXPRESSION TAG	UNP A0R3M0
C	-4	HIS	-	EXPRESSION TAG	UNP A0R3M0
C	-3	HIS	-	EXPRESSION TAG	UNP A0R3M0
C	-2	MSE	-	EXPRESSION TAG	UNP A0R3M0
C	-1	ALA	-	EXPRESSION TAG	UNP A0R3M0
C	0	SER	-	EXPRESSION TAG	UNP A0R3M0
D	-12	MSE	-	EXPRESSION TAG	UNP A0R3M0
D	-11	ARG	-	EXPRESSION TAG	UNP A0R3M0
D	-10	GLY	-	EXPRESSION TAG	UNP A0R3M0
D	-9	SER	-	EXPRESSION TAG	UNP A0R3M0
D	-8	HIS	-	EXPRESSION TAG	UNP A0R3M0
D	-7	HIS	-	EXPRESSION TAG	UNP A0R3M0
D	-6	HIS	-	EXPRESSION TAG	UNP A0R3M0
D	-5	HIS	-	EXPRESSION TAG	UNP A0R3M0
D	-4	HIS	-	EXPRESSION TAG	UNP A0R3M0
D	-3	HIS	-	EXPRESSION TAG	UNP A0R3M0
D	-2	MSE	-	EXPRESSION TAG	UNP A0R3M0
D	-1	ALA	-	EXPRESSION TAG	UNP A0R3M0
D	0	SER	-	EXPRESSION TAG	UNP A0R3M0
E	-12	MSE	-	EXPRESSION TAG	UNP A0R3M0
E	-11	ARG	-	EXPRESSION TAG	UNP A0R3M0
E	-10	GLY	-	EXPRESSION TAG	UNP A0R3M0
E	-9	SER	-	EXPRESSION TAG	UNP A0R3M0
E	-8	HIS	-	EXPRESSION TAG	UNP A0R3M0
E	-7	HIS	-	EXPRESSION TAG	UNP A0R3M0
E	-6	HIS	-	EXPRESSION TAG	UNP A0R3M0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-5	HIS	-	EXPRESSION TAG	UNP A0R3M0
E	-4	HIS	-	EXPRESSION TAG	UNP A0R3M0
E	-3	HIS	-	EXPRESSION TAG	UNP A0R3M0
E	-2	MSE	-	EXPRESSION TAG	UNP A0R3M0
E	-1	ALA	-	EXPRESSION TAG	UNP A0R3M0
E	0	SER	-	EXPRESSION TAG	UNP A0R3M0
F	-12	MSE	-	EXPRESSION TAG	UNP A0R3M0
F	-11	ARG	-	EXPRESSION TAG	UNP A0R3M0
F	-10	GLY	-	EXPRESSION TAG	UNP A0R3M0
F	-9	SER	-	EXPRESSION TAG	UNP A0R3M0
F	-8	HIS	-	EXPRESSION TAG	UNP A0R3M0
F	-7	HIS	-	EXPRESSION TAG	UNP A0R3M0
F	-6	HIS	-	EXPRESSION TAG	UNP A0R3M0
F	-5	HIS	-	EXPRESSION TAG	UNP A0R3M0
F	-4	HIS	-	EXPRESSION TAG	UNP A0R3M0
F	-3	HIS	-	EXPRESSION TAG	UNP A0R3M0
F	-2	MSE	-	EXPRESSION TAG	UNP A0R3M0
F	-1	ALA	-	EXPRESSION TAG	UNP A0R3M0
F	0	SER	-	EXPRESSION TAG	UNP A0R3M0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	36	Total O 36 36	0	0
2	B	29	Total O 29 29	0	0
2	C	37	Total O 37 37	0	0
2	D	18	Total O 18 18	0	0
2	E	25	Total O 25 25	0	0
2	F	15	Total O 15 15	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acetyl-CoA acetyltransferase




- Molecule 1: Acetyl-CoA acetyltransferase




- Molecule 1: Acetyl-CoA acetyltransferase





• Molecule 1: Acetyl-CoA acetyltransferase



• Molecule 1: Acetyl-CoA acetyltransferase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	96.31Å 102.00Å 102.42Å 116.46° 101.28° 97.44°	Depositor
Resolution (Å)	47.95 – 2.71 47.95 – 2.71	Depositor EDS
% Data completeness (in resolution range)	96.9 (47.95-2.71) 80.9 (47.95-2.71)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.27 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.234 , 0.261 0.244 , 0.271	Depositor DCC
R_{free} test set	4512 reflections (5.43%)	DCC
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.8	EDS
Estimated twinning fraction	0.018 for -h,-l,-k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 87684 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	20206	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.99	2/3377 (0.1%)	0.88	5/4572 (0.1%)
1	B	0.93	1/3435 (0.0%)	0.88	6/4649 (0.1%)
1	C	0.92	2/3424 (0.1%)	0.86	3/4634 (0.1%)
1	D	0.92	1/3348 (0.0%)	0.84	4/4532 (0.1%)
1	E	0.91	0/3388	0.87	5/4582 (0.1%)
1	F	0.92	1/3384 (0.0%)	0.88	7/4580 (0.2%)
All	All	0.93	7/20356 (0.0%)	0.87	30/27549 (0.1%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	477	GLU	CG-CD	5.93	1.60	1.51
1	A	44	CYS	CB-SG	-5.59	1.72	1.81
1	A	31	GLU	CG-CD	5.42	1.60	1.51
1	D	332	CYS	CB-SG	5.38	1.91	1.82
1	F	294	LEU	C-O	-5.26	1.13	1.23

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	ASP	CB-CG-OD2	6.00	123.70	118.30
1	F	32	LEU	CB-CG-CD1	5.97	121.15	111.00
1	F	174	GLY	N-CA-C	-5.95	98.23	113.10
1	C	487	ASP	CB-CG-OD1	5.94	123.65	118.30
1	A	466	ASP	CB-CG-OD1	5.86	123.58	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3327	0	3278	77	0
1	B	3376	0	3320	82	0
1	C	3373	0	3329	84	0
1	D	3299	0	3252	89	0
1	E	3336	0	3296	93	0
1	F	3335	0	3283	85	0
2	A	36	0	0	3	0
2	B	29	0	0	2	0
2	C	37	0	0	3	0
2	D	18	0	0	2	0
2	E	25	0	0	3	0
2	F	15	0	0	1	0
All	All	20206	0	19758	495	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 495 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:176:ILE:HD13	1:B:176:ILE:C	1.51	1.25
1:F:176:ILE:HD12	1:F:176:ILE:O	1.34	1.22
1:A:489:ILE:HD12	1:A:489:ILE:N	1.57	1.13
1:A:489:ILE:H	1:A:489:ILE:CD1	1.50	1.12
1:A:175:LEU:HD23	1:A:175:LEU:N	1.58	1.10

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/520 (84%)	417 (95%)	20 (5%)	1 (0%)	52	80
1	B	445/520 (86%)	425 (96%)	19 (4%)	1 (0%)	52	80
1	C	443/520 (85%)	423 (96%)	20 (4%)	0	100	100
1	D	435/520 (84%)	414 (95%)	21 (5%)	0	100	100
1	E	437/520 (84%)	420 (96%)	15 (3%)	2 (0%)	34	62
1	F	437/520 (84%)	420 (96%)	17 (4%)	0	100	100
All	All	2635/3120 (84%)	2519 (96%)	112 (4%)	4 (0%)	52	80

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	119	GLY
1	E	412	ALA
1	E	466	ASP
1	A	489	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/394 (87%)	318 (93%)	25 (7%)	17	39
1	B	347/394 (88%)	309 (89%)	38 (11%)	8	17
1	C	348/394 (88%)	319 (92%)	29 (8%)	14	31
1	D	340/394 (86%)	304 (89%)	36 (11%)	8	19
1	E	344/394 (87%)	310 (90%)	34 (10%)	10	22
1	F	344/394 (87%)	311 (90%)	33 (10%)	10	23
All	All	2066/2364 (87%)	1871 (91%)	195 (9%)	11	24

5 of 195 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	494	VAL
1	D	314	LEU
1	F	273	LEU
1	C	499	GLU
1	D	184	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	388	GLN
1	D	257	ASN
1	F	369	ASN
1	D	106	GLN
1	D	368	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/520 (84%)	0.10	4 (0%) 85 86	23, 46, 64, 69	0
1	B	442/520 (85%)	0.02	4 (0%) 85 86	25, 46, 64, 69	0
1	C	443/520 (85%)	0.08	7 (1%) 74 75	25, 46, 64, 69	0
1	D	435/520 (83%)	-0.10	7 (1%) 74 75	25, 46, 64, 69	0
1	E	437/520 (84%)	-0.03	4 (0%) 85 86	25, 46, 64, 69	0
1	F	440/520 (84%)	0.02	6 (1%) 78 78	25, 46, 64, 76	0
All	All	2636/3120 (84%)	0.02	32 (1%) 81 81	23, 46, 64, 76	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	119	GLY	6.0
1	C	119	GLY	4.3
1	C	25	ARG	3.9
1	F	133	SER	3.9
1	B	492	LYS	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.