



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:52 PM GMT

PDB ID : 5EHU
Title : sfGFP mutant with unnatural amino acid 4-azidoethoxy-L-phenylalanine incorporated at the 149 site
Authors : Tookmanian, E.M.; Phillips-Piro, C.M.; Fenlon, E.E.; Brewer, S.B.
Deposited on : 2015-10-28
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1278 (1.48-1.44)
Clashscore	102246	1336 (1.48-1.44)
Ramachandran outliers	100387	1320 (1.48-1.44)
Sidechain outliers	100360	1320 (1.48-1.44)
RSRZ outliers	91569	1279 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	<div> <div>3%</div> <div>92%</div> <div>• • •</div> </div>
1	B	237	<div> <div>%</div> <div>92%</div> <div>5% •</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7903 atoms, of which 3609 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	229	Total	C	H	N	O	S	0	10	0
			3678	1181	1808	324	360	5			
1	B	229	Total	C	H	N	O	S	0	7	0
			3672	1181	1801	325	360	5			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP A0A059PIQ0
A	1	VAL	-	expression tag	UNP A0A059PIQ0
A	2	SER	ARG	engineered mutation	UNP A0A059PIQ0
A	30	ARG	SER	engineered mutation	UNP A0A059PIQ0
A	66	CRO	THR	chromophore	UNP A0A059PIQ0
A	66	CRO	TYR	chromophore	UNP A0A059PIQ0
A	66	CRO	GLY	chromophore	UNP A0A059PIQ0
A	72	SER	ALA	engineered mutation	UNP A0A059PIQ0
A	80	ARG	GLN	engineered mutation	UNP A0A059PIQ0
A	149	4LZ	ASN	engineered mutation	UNP A0A059PIQ0
A	206	VAL	ALA	engineered mutation	UNP A0A059PIQ0
B	0	MET	-	expression tag	UNP A0A059PIQ0
B	1	VAL	-	expression tag	UNP A0A059PIQ0
B	2	SER	ARG	engineered mutation	UNP A0A059PIQ0
B	30	ARG	SER	engineered mutation	UNP A0A059PIQ0
B	66	CRO	THR	chromophore	UNP A0A059PIQ0
B	66	CRO	TYR	chromophore	UNP A0A059PIQ0
B	66	CRO	GLY	chromophore	UNP A0A059PIQ0
B	72	SER	ALA	engineered mutation	UNP A0A059PIQ0
B	80	ARG	GLN	engineered mutation	UNP A0A059PIQ0
B	149	4LZ	ASN	engineered mutation	UNP A0A059PIQ0
B	206	VAL	ALA	engineered mutation	UNP A0A059PIQ0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	279	Total 279	O 279	0	0
2	B	274	Total 274	O 274	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	129.92Å 37.49Å 91.85Å 90.00° 106.31° 90.00°	Depositor
Resolution (Å)	45.26 – 1.45 45.26 – 1.45	Depositor EDS
% Data completeness (in resolution range)	96.2 (45.26-1.45) 91.3 (45.26-1.45)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.45Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.175 , 0.208 0.178 , 0.205	Depositor DCC
R_{free} test set	1995 reflections (2.95%)	DCC
Wilson B-factor (Å ²)	14.7	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 43.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	4 of 73332 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7903	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 95.46 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3289e-09. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4LZ, CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/1911	0.67	1/2586 (0.0%)
1	B	0.44	0/1882	0.64	0/2543
All	All	0.45	0/3793	0.65	1/5129 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	153	THR	N-CA-C	5.82	126.71	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1870	1808	1769	10	0
1	B	1871	1801	1785	11	0
2	A	279	0	0	6	0
2	B	274	0	0	6	0
All	All	4294	3609	3554	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:17:GLU:OE2	1:B:122:ARG:NH1	2.09	0.86
1:B:73[A]:ARG:NH1	2:B:301:HOH:O	2.19	0.75
1:A:153:THR:O	2:A:301:HOH:O	2.07	0.72
1:B:166:LYS:NZ	2:B:302:HOH:O	2.26	0.64
1:A:115:GLU:OE2	1:A:122:ARG:NH2	2.29	0.64
1:A:80:ARG:NH2	2:A:307:HOH:O	2.32	0.62
1:B:131[A]:LYS:NZ	2:B:306:HOH:O	2.35	0.60
1:A:132:GLU:OE1	1:B:73[A]:ARG:NH2	2.36	0.58
1:A:76:ASP:OD1	2:A:302:HOH:O	2.17	0.56
1:B:17:GLU:OE1	1:B:30:ARG:NH2	2.39	0.56
1:B:80:ARG:NH1	2:B:309:HOH:O	2.41	0.53
1:B:73[A]:ARG:NE	2:B:308:HOH:O	2.41	0.53
1:A:180:ASP:OD2	2:A:304:HOH:O	2.20	0.47
1:B:180[B]:ASP:OD2	1:B:182:TYR:OH	2.21	0.47
1:A:142[A]:GLU:OE1	2:A:303:HOH:O	2.19	0.46
1:B:2:SER:N	2:B:312:HOH:O	2.50	0.44
1:A:153:THR:O	1:A:154:ALA:HB3	2.18	0.44
1:A:73[A]:ARG:NH1	2:A:315:HOH:O	2.53	0.42
1:A:133:ASP:OD1	1:A:133:ASP:N	2.46	0.41
1:B:133:ASP:N	1:B:133:ASP:OD2	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/237 (98%)	230 (99%)	2 (1%)	1 (0%)	39	13
1	B	230/237 (97%)	225 (98%)	5 (2%)	0	100	100
All	All	463/474 (98%)	455 (98%)	7 (2%)	1 (0%)	52	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/206 (99%)	204 (100%)	0	100	100
1	B	201/206 (98%)	201 (100%)	0	100	100
All	All	405/412 (98%)	405 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	4LZ	A	149	1	14,17,18	2.03	3 (21%)	15,20,22	1.09	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	A	66	1	23,23,24	1.82	3 (13%)	30,32,34	2.29	9 (30%)
1	4LZ	B	149	1	14,17,18	2.01	3 (21%)	15,20,22	1.19	1 (6%)
1	CRO	B	66	1	23,23,24	1.86	3 (13%)	30,32,34	2.27	8 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	4LZ	A	149	1	-	0/10/12/14	0/1/1/1
1	CRO	A	66	1	-	0/12/31/32	0/2/2/2
1	4LZ	B	149	1	-	0/10/12/14	0/1/1/1
1	CRO	B	66	1	-	0/12/31/32	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	CRO	CG2-CB2	2.19	1.51	1.46
1	A	66	CRO	CG2-CB2	2.26	1.51	1.46
1	B	149	4LZ	OH-CZ	2.28	1.43	1.37
1	A	149	4LZ	OH-CZ	2.43	1.43	1.37
1	A	149	4LZ	CB-CG	2.70	1.57	1.51
1	A	66	CRO	C2-N3	2.80	1.46	1.39
1	B	149	4LZ	CB-CG	2.93	1.58	1.51
1	B	66	CRO	C2-N3	3.19	1.47	1.39
1	A	149	4LZ	N7-N6	5.65	1.38	1.23
1	B	149	4LZ	N7-N6	5.66	1.38	1.23
1	B	66	CRO	CA2-C2	6.77	1.55	1.48
1	A	66	CRO	CA2-C2	6.88	1.55	1.48

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CRO	CA1-C1-N3	-5.46	118.37	124.98
1	B	66	CRO	CA1-C1-N3	-4.98	118.94	124.98
1	A	66	CRO	CG2-CB2-CA2	-3.86	125.38	130.27
1	B	66	CRO	C2-CA2-N2	-3.75	106.18	109.03
1	A	66	CRO	C2-CA2-N2	-3.34	106.49	109.03
1	B	66	CRO	CG2-CB2-CA2	-2.96	126.53	130.27
1	A	66	CRO	CA2-C2-N3	-2.58	102.09	103.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	149	4LZ	O-C-CA	-2.05	120.23	125.72
1	A	149	4LZ	C31-OH-CZ	-2.04	112.33	117.90
1	A	66	CRO	CD2-CG2-CD1	2.34	121.13	117.62
1	B	66	CRO	CB2-CA2-N2	2.38	132.93	128.71
1	A	66	CRO	CB2-CA2-N2	2.40	132.97	128.71
1	B	66	CRO	CD2-CG2-CD1	2.41	121.24	117.62
1	A	66	CRO	CA3-N3-C1	2.74	130.41	127.38
1	B	66	CRO	CA3-N3-C1	2.90	130.58	127.38
1	B	66	CRO	CA1-C1-N2	3.34	128.49	123.71
1	A	66	CRO	CA1-C1-N2	3.45	128.65	123.71
1	A	66	CRO	CA2-N2-C1	6.16	110.72	105.54
1	B	66	CRO	CA2-N2-C1	7.16	111.55	105.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/237 (95%)	-0.26	6 (2%) 59 60	10, 17, 33, 65	0
1	B	227/237 (95%)	-0.23	3 (1%) 79 80	10, 18, 35, 55	0
All	All	454/474 (95%)	-0.25	9 (1%) 68 69	10, 18, 36, 65	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	SER	5.5
1	B	2	SER	3.9
1	B	156	LYS	3.6
1	A	157	GLN	3.2
1	A	154	ALA	2.8
1	B	232	GLY	2.4
1	A	232	GLY	2.2
1	A	156	LYS	2.1
1	A	153	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	4LZ	B	149	17/18	0.96	0.08	-	14,21,36,39	0
1	4LZ	A	149	17/18	0.94	0.10	-	13,21,36,43	0
1	CRO	A	66	22/23	0.98	0.07	-	8,11,20,20	0
1	CRO	B	66	22/23	0.98	0.07	-	8,11,18,18	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.