Full wwPDB X-ray Structure Validation Report (i)

Jan 31, 2016 – 07:12 PM GMT

PDB ID	:	1EJU
Title	:	CRYSTAL STRUCTURE OF THE H320N VARIANT OF KLEBSIELLA AEROGENES UREASE
Authors	:	Pearson, M.A.; Park, I.S.; Schaller, R.A.; Michel, L.O.; Karplus, P.A.; Hausinger, R.P.
Deposited on	:	2000-03-04
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	567	% 85% 11% • •
2	B	101	2% 89% 11%
3	A	100	97% •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE ALPHA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	553	4128	2590	724	792	22	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1217	KCX	LYS	MODIFIED RESIDUE	UNP P18314
C	1320	ASN	HIS	ENGINEERED	UNP P18314

- Molecule 2 is a protein called UREASE BETA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	101	785	496	150	136	3	0	0	0

- Molecule 3 is a protein called UREASE GAMMA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	100	776	491	134	146	5	0	0	0

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ni		
4	C	2	2	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	54	54	54	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	B	48	48	48	0	0
			Total	O		
5	C	306	306	306	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UREASE ALPHA SUBUNIT

Chain C:



- Molecule 2: UREASE BETA SUBUNIT

Chain B:



- Molecule 3: UREASE GAMMA SUBUNIT

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α, β, γ	170.80Å 170.80Å 170.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 31.18 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.00) 86.8 (31.18-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.63 (at 2.00Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.187 , (Not available) 0.180 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å^2)	18.1	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 73.1	EDS
Estimated twinning fraction	0.020 for -l,-k,-h	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 48394 reflections	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6099	wwPDB-VP
Average B, all atoms (Å^2)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	C	0.33	0/4196	0.68	3/5713 (0.1%)
2	B	0.32	0/805	0.59	0/1087
3	A	0.31	0/787	0.54	0/1061
All	All	0.32	0/5788	0.65	3/7861 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	1299	ASN	N-CA-C	5.91	126.95	111.00
1	C	1316	LEU	CB-CG-CD1	-5.39	101.84	111.00
1	C	1372	LEU	CA-CB-CG	5.07	126.96	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	4128	0	4085	37	0
2	B	785	0	772	6	0
3	A	776	0	804	1	0
4	C	2	0	0	0	0
5	A	54	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	48	0	0	0	0
5	C	306	0	0	1	0
All	All	6099	0	5661	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1508:ARG:HH11	1:C:1508:ARG:HG2	1.25	0.98
1:C:1545:VAL:O	1:C:1548:GLU:HG2	1.90	0.72
2:B:2100:PRO:O	2:B:2101:LEU:HB2	1.90	0.70
1:C:1508:ARG:NH1	1:C:1508:ARG:HG2	2.04	0.68
1:C:1313:LEU:O	1:C:1317:MET:HG2	1.98	0.63
1:C:1201:GLN:HE21	1:C:1203:ASP:H	1.50	0.60
1:C:1460:ASP:HB3	1:C:1463:ALA:HB2	1.86	0.56
2:B:2089:ALA:HA	2:B:2100:PRO:HA	1.86	0.56
1:C:1022:ARG:O	2:B:2004:GLY:HA2	2.07	0.54
1:C:1363:ALA:O	1:C:1364:MET:HB2	2.08	0.54
2:B:2072:GLU:H	2:B:2075:GLN:NE2	2.09	0.51
1:C:1218:ILE:HD11	1:C:1232:ALA:CB	2.41	0.51
1:C:1442:VAL:HG12	1:C:1443:LYS:HG3	1.93	0.50
1:C:1311:GLU:HG3	1:C:1312:HIS:N	2.27	0.50
1:C:1508:ARG:CG	1:C:1508:ARG:HH11	2.11	0.49
1:C:1298:THR:CG2	1:C:1360:ASP:HB2	2.43	0.49
1:C:1020:LYS:HB2	2:B:2007:HIS:HB3	1.95	0.49
1:C:1474:ARG:HD3	5:C:106:HOH:O	2.12	0.48
1:C:1297:SER:OG	1:C:1349:HIS:HE1	1.96	0.48
1:C:1138:ILE:HD12	1:C:1362:GLN:HB2	1.96	0.48
1:C:1395:ASN:HD22	1:C:1397:ASN:H	1.62	0.47
1:C:1355:SER:HB2	1:C:1356:LEU:HD12	1.97	0.47
1:C:1545:VAL:HG23	1:C:1550:ILE:HG13	1.96	0.46
1:C:1465:ILE:HG13	1:C:1468:PRO:HD3	1.99	0.45
1:C:1304:TYR:OH	1:C:1339:ARG:HG3	2.17	0.45
1:C:1380:ARG:NE	1:C:1555:ALA:HB2	2.32	0.45
1:C:1218:ILE:HD13	1:C:1243:VAL:HG13	1.99	0.44
3:A:3081:GLN:HG2	3:A:3094:THR:HG23	2.00	0.43
1:C:1489:LEU:HD23	1:C:1510:ALA:HB3	2.01	0.43
1:C:1359:SER:O	1:C:1365:GLY:HA3	2.19	0.43
1:C:1300:PRO:HG2	1:C:1364:MET:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1170:CYS:O	1:C:1172:PRO:HD3	2.19	0.42
1:C:1395:ASN:C	1:C:1395:ASN:HD22	2.22	0.42
1:C:1080:GLY:HA2	1:C:1401:LYS:HE2	2.00	0.42
1:C:1075:ILE:O	1:C:1081:ILE:HA	2.19	0.42
1:C:1296:SER:HB3	1:C:1356:LEU:HB2	2.01	0.41
2:B:2039:HIS:HA	2:B:2060:ARG:HD2	2.01	0.41
1:C:1012:MET:O	1:C:1044:LYS:HE2	2.20	0.41
1:C:1154:MET:SD	1:C:1189:VAL:HG21	2.61	0.41
1:C:1306:LEU:HG	1:C:1556:ASP:HA	2.03	0.41
1:C:1246:HIS:CE1	1:C:1278:GLY:HA3	2.56	0.40
1:C:1249:THR:HG21	1:C:1336:ARG:HD2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	548/567 (97%)	514 (94%)	31 (6%)	3 (0%)	34 26
2	B	99/101 (98%)	93 (94%)	6 (6%)	0	100 100
3	A	98/100 (98%)	98 (100%)	0	0	100 100
All	All	745/768 (97%)	705 (95%)	37 (5%)	3 (0%)	39 33

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	1360	ASP
1	C	1364	MET
1	C	1481	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	430/443 (97%)	419 (97%)	11 (3%)	54	54
2	B	78/78 (100%)	76 (97%)	2 (3%)	54	54
3	A	85/85 (100%)	84 (99%)	1 (1%)	78	81
All	All	593/606 (98%)	579 (98%)	14 (2%)	57	58

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1179	ARG
1	C	1282	PRO
1	C	1332	PHE
1	C	1336	ARG
1	C	1349	HIS
1	C	1394	ASP
1	C	1395	ASN
1	C	1449	LYS
1	C	1506	ASN
1	C	1508	ARG
1	C	1539	GLN
2	B	2009	LYS
2	B	2077	ARG
3	A	3100	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1142	GLN
1	C	1166	HIS
1	C	1201	GLN
1	C	1349	HIS
1	C	1362	GLN
1	C	1395	ASN
1	C	1469	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	1486	HIS
2	B	2016	ASN
2	B	2054	GLN
2	B	2075	GLN
3	A	3081	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	KCX	C	1217	1,4	7,11,12	0.49	0	7,12,14	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	C	1217	1,4	-	0/6/10/12	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	C	552/567 (97%)	-0.53	6 (1%) 82 83	7, 15, 33, 98	0
2	B	101/101 (100%)	-0.24	2 (1%) 68 69	13, 22, 43, 61	0
3	A	100/100 (100%)	-0.55	0 100 100	8, 16, 34, 46	0
All	All	753/768 (98%)	-0.49	8 (1%) 82 83	7, 16, 37, 98	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1332	PHE	5.2
1	C	1313	LEU	4.8
1	C	1317	MET	4.8
1	C	1333	ALA	2.5
1	C	1499	ASN	2.2
2	B	2054	GLN	2.1
1	C	1316	LEU	2.1
2	B	2101	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
1	KCX	C	1217	12/13	0.97	0.10	-	6,13,17,17	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
4	NI	C	4775	1/1	1.00	0.03	-3.01	24,24,24,24	0
4	NI	C	4774	1/1	1.00	0.02	-6.68	21,21,21,21	0

6.5 Other polymers (i)

There are no such residues in this entry.