



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:25 AM GMT

PDB ID : 3ELA
Title : Crystal structure of active site inhibited coagulation factor VIIA mutant in complex with soluble tissue factor
Authors : Bjelke, J.R.; Fodje, M.; Svensson, L.A.
Deposited on : 2008-09-21
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	152	
2	H	254	
3	T	209	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4389 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor VIIA light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	95	703	423	122	145	13	0	0	0

- Molecule 2 is a protein called Coagulation factor VIIA heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	254	1911	1216	331	352	12	0	0	0


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	158	ASP	VAL	ENGINEERED	UNP P08709
H	296	VAL	GLU	ENGINEERED	UNP P08709
H	298	GLN	MET	ENGINEERED	UNP P08709

- Molecule 3 is a protein called Tissue factor.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	T	191	1505	956	242	303	4	0	0	0

- Molecule 4 is SUGAR (ALPHA-D-GLUCOSE) (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is SUGAR (ALPHA-L-FUCOSE) (three-letter code: FUC) (formula: $C_6H_{12}O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	1	Total Ca 1 1	0	0

- Molecule 7 is D-PHENYLALANYL-N-[(2S,3S)-6- {[AMINO(IMINIO)METHYL]AMINO}-1-CHLORO-2-HYDROXYHEXAN-3-YL]-L-PHENYLALANINAMIDE (three-letter code: OZ6) (formula: C₂₅H₃₆ClN₆O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	H	1	Total C N O 34 25 6 3	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	L	39	Total O 39 39	0	0
8	H	123	Total O 123 123	0	0
8	T	52	Total O 52 52	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.36Å 68.56Å 78.82Å 90.00° 90.22° 90.00°	Depositor
Resolution (Å)	29.19 – 2.20 29.19 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.19-2.20) 81.1 (29.19-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.233 , 0.294 0.242 , 0.295	Depositor DCC
R_{free} test set	1734 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	19.8	Xtriage
Anisotropy	1.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.8	EDS
Estimated twinning fraction	0.017 for l,k,-h 0.078 for h,-k,-l 0.037 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 34672 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4389	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC, FUC, OZ6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.91	3/716 (0.4%)	0.86	0/967
2	H	0.85	1/1959 (0.1%)	0.95	2/2674 (0.1%)
3	T	0.70	0/1536	0.86	3/2097 (0.1%)
All	All	0.81	4/4211 (0.1%)	0.91	5/5738 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	2
3	T	0	10
All	All	0	12

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	114	CYS	CB-SG	-8.96	1.67	1.82
2	H	357	TYR	CD2-CE2	5.88	1.48	1.39
1	L	116	GLU	CG-CD	5.05	1.59	1.51
1	L	127	CYS	CB-SG	-5.02	1.73	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	188	SER	CB-CA-C	-8.54	93.88	110.10
3	T	189	VAL	N-CA-C	7.47	131.18	111.00
2	H	271	ARG	NE-CZ-NH2	-5.83	117.39	120.30
3	T	44	ASP	CB-CG-OD2	-5.70	113.17	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	170	LEU	CB-CG-CD1	-5.30	102.00	111.00

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	218	GLY	Peptide
2	H	352	TYR	Peptide
3	T	81	GLY	Peptide
3	T	82	ASN	Peptide
3	T	83	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	703	0	622	21	0
2	H	1911	0	1816	85	0
3	T	1505	0	1421	78	0
4	L	11	0	10	1	0
5	L	10	0	10	1	0
6	H	1	0	0	0	0
7	H	34	0	32	6	0
8	H	123	0	0	18	0
8	L	39	0	0	3	0
8	T	52	0	0	3	0
All	All	4389	0	3911	178	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 178 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:T:87:GLY:CA	3:T:89:ALA:H	1.34	1.39

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:T:87:GLY:HA2	3:T:89:ALA:N	1.03	1.32
3:T:87:GLY:HA2	3:T:88:SER:C	1.58	1.23
3:T:85:SER:O	3:T:86:THR:HG22	1.37	1.20
3:T:87:GLY:CA	3:T:89:ALA:N	1.93	1.19

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	93/152 (61%)	79 (85%)	13 (14%)	1 (1%)	17	14
2	H	252/254 (99%)	222 (88%)	24 (10%)	6 (2%)	7	4
3	T	185/209 (88%)	160 (86%)	18 (10%)	7 (4%)	4	1
All	All	530/615 (86%)	461 (87%)	55 (10%)	14 (3%)	7	3

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	57	ASN
2	H	353	ARG
3	T	86	THR
3	T	88	SER
3	T	89	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	79/132 (60%)	69 (87%)	10 (13%)	5	4
2	H	199/216 (92%)	188 (94%)	11 (6%)	27	30
3	T	169/192 (88%)	148 (88%)	21 (12%)	6	5
All	All	447/540 (83%)	405 (91%)	42 (9%)	11	10

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	320	SER
3	T	22	ILE
3	T	197	THR
2	H	394	GLU
3	T	6	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	227	GLN
2	H	240	ASN
3	T	114	GLN
2	H	216	HIS
3	T	69	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
7	0Z6	H	407	2	31,35,36	2.02	2 (6%)	39,45,46	0.78	1 (2%)
4	GLC	L	215	-	11,11,12	0.57	0	14,15,17	1.71	2 (14%)
5	FUC	L	216	-	10,10,11	0.57	0	14,14,16	2.42	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	0Z6	H	407	2	-	0/33/35/37	0/2/2/2
4	GLC	L	215	-	-	0/2/19/22	0/1/1/1
5	FUC	L	216	-	-	0/0/17/20	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	H	407	0Z6	O2-C2	-8.89	1.23	1.43
7	H	407	0Z6	C3-C2	-5.51	1.38	1.51

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	L	216	FUC	O5-C5-C6	2.03	109.48	106.13
5	L	216	FUC	O5-C5-C4	2.23	113.39	109.53
4	L	215	GLC	O2-C2-C1	2.64	114.51	109.21
7	H	407	0Z6	O2-C2-CA2	2.90	115.10	108.58
4	L	215	GLC	C1-O5-C5	3.79	117.06	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	H	407	OZ6	6	0
4	L	215	GLC	1	0
5	L	216	FUC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	95/152 (62%)	-0.27	0 100 100	12, 29, 60, 65	0
2	H	254/254 (100%)	-0.30	1 (0%) 93 93	7, 22, 52, 61	4 (1%)
3	T	191/209 (91%)	0.46	22 (11%) 6 6	14, 46, 89, 92	10 (5%)
All	All	540/615 (87%)	-0.03	23 (4%) 39 38	7, 28, 79, 92	14 (2%)

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	T	119	VAL	8.8
3	T	123	VAL	6.6
3	T	138	ASN	5.8
3	T	158	TRP	5.6
3	T	118	GLN	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	FUC	L	216	10/11	0.93	0.14	0.29	67,68,69,70	0
7	OZ6	H	407	34/35	0.93	0.12	-0.06	14,31,41,42	0
4	GLC	L	215	11/12	0.80	0.11	-0.62	52,53,53,54	0
6	CA	H	1	1/1	0.96	0.05	-3.72	50,50,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.