



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 20, 2016 – 11:53 PM EDT

PDB ID : 5EYP
Title : TUBULIN-DARPIN COMPLEX
Authors : Ahmad, S.; Knossow, M.; Gigant, B.
Deposited on : 2015-11-25
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	
2	B	445	
3	F	169	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	GOL	A	505	-	-	-	X
7	GOL	A	506	-	-	-	X
7	GOL	A	507	-	-	-	X
7	GOL	B	505	-	-	-	X
7	GOL	F	202	-	-	-	X

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 8501 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	428	Total	C	N	O	S	0	1	0
			3328	2109	566	631	22			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	SER	GLY	variant	UNP D0VWZ0
A	340	SER	THR	variant	UNP D0VWZ0

- Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	425	Total	C	N	O	S	0	12	0
			3405	2140	581	657	27			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	203	CYS	SER	variant	UNP D0VWY9
B	318	ILE	VAL	variant	UNP D0VWY9

- Molecule 3 is a protein called DESIGNED ANKYRIN REPEAT PROTEIN (DARPIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	130	Total	C	N	O	S	0	1	0
			960	603	171	183	3			

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0


- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0
7	F	1	Total C O 6 3 3	0	0
7	F	1	Total C O 6 3 3	0	0
7	F	1	Total C O 6 3 3	0	0

- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:

C₁₀H₁₅N₅O₁₁P₂).




- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	281	Total 281	O 281	0	0
10	B	281	Total 281	O 281	0	0
10	F	93	Total 93	O 93	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tubulin alpha chain



• Molecule 2: Tubulin beta chain



• Molecule 3: DESIGNED ANKYRIN REPEAT PROTEIN (DARPIN)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.57Å 71.76Å 93.05Å 90.00° 99.85° 90.00°	Depositor
Resolution (Å)	49.74 – 1.90 49.74 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.74-1.90) 100.0 (49.74-1.90)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.90Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.173 , 0.208 0.174 , 0.207	Depositor DCC
R_{free} test set	4531 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8501	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, LOC, MG, GOL, GTP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.49	0/3404	0.64	0/4623
2	B	0.53	0/3509	0.63	0/4752
3	F	0.49	0/976	0.64	0/1327
All	All	0.51	0/7889	0.64	0/10702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3328	0	3240	19	0
2	B	3405	0	3296	28	0
3	F	960	0	979	5	0
4	A	32	0	12	0	0
5	A	1	0	0	0	0
6	A	10	0	0	0	0
6	B	5	0	0	0	0
7	A	18	0	24	1	0
7	B	12	0	16	0	0
7	F	18	0	24	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	28	0	12	0	0
9	B	29	0	25	1	0
10	A	281	0	0	3	0
10	B	281	0	0	4	0
10	F	93	0	0	0	0
All	All	8501	0	7628	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:ASN:HD22	2:B:258:ASN:HD21	1.47	0.63
2:B:318:ILE:HD11	9:B:502:LOC:H4	1.84	0.60
2:B:57:THR:HG22	10:B:764:HOH:O	2.00	0.59
2:B:106:GLY:O	2:B:111:GLY:HA3	2.03	0.58
2:B:345:GLU:OE1	10:B:601:HOH:O	2.17	0.56
2:B:400:ARG:HD2	3:F:112:TRP:NE1	2.22	0.54
2:B:295:MET:HG2	2:B:377:PHE:HB2	1.89	0.54
1:A:221:ARG:NE	2:B:325:MET:HB3	2.24	0.52
1:A:406:HIS:CG	2:B:263:PRO:HD3	2.47	0.50
2:B:214:PHE:O	2:B:218:LYS:HA	2.13	0.49
1:A:28:HIS:CE1	1:A:49:PHE:HB3	2.48	0.49
3:F:92:HIS:HE1	7:F:203:GOL:H32	1.77	0.48
1:A:266:HIS:HE1	10:A:675:HOH:O	1.96	0.48
2:B:147[A]:SER:HB2	2:B:190:SER:HG	1.79	0.48
2:B:75:MET:HG3	2:B:94:PHE:CD1	2.49	0.47
2:B:192:HIS:CD2	2:B:424[A]:ASN:HD22	2.32	0.47
1:A:307:PRO:HA	1:A:383:ALA:HB2	1.96	0.46
2:B:83:PHE:O	2:B:86:ILE:HG22	2.14	0.46
3:F:79:MET:HE2	3:F:111:THR:HG21	1.96	0.46
2:B:269:MET:HG3	2:B:303:ALA:HB3	1.96	0.46
2:B:294:GLN:HE21	2:B:300:ASN:HD21	1.63	0.45
2:B:396:THR:O	2:B:400:ARG:HG3	2.15	0.45
1:A:8:HIS:HE1	10:A:836:HOH:O	1.99	0.45
2:B:172:MET:HG3	2:B:387:LEU:HD21	1.99	0.45
1:A:210:TYR:CZ	1:A:214:ARG:HD2	2.52	0.45
2:B:395:PHE:CE1	2:B:422[B]:GLU:HB2	2.52	0.45
2:B:181:VAL:HG21	3:F:90:ILE:HG23	1.99	0.45
1:A:175:PRO:HA	1:A:179:THR:HG22	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:320:ARG:HA	1:A:356:ASN:O	2.17	0.44
1:A:406:HIS:HB3	7:A:506:GOL:H32	1.98	0.44
2:B:66:ILE:HG12	2:B:121:VAL:HG12	2.00	0.44
2:B:275:LEU:HA	2:B:275:LEU:HD12	1.87	0.43
2:B:400:ARG:HD3	10:B:844:HOH:O	2.18	0.43
3:F:79:MET:HA	3:F:79:MET:HE2	2.00	0.43
1:A:167:LEU:HG	1:A:200:CYS:HB3	2.02	0.42
1:A:406:HIS:HD2	10:A:849:HOH:O	2.02	0.42
2:B:215[B]:ARG:HH22	2:B:275:LEU:HD23	1.83	0.42
1:A:238:ILE:HG12	1:A:378:LEU:HD21	2.02	0.42
2:B:147[B]:SER:OG	2:B:190:SER:OG	2.27	0.42
2:B:247:GLN:NE2	10:B:603:HOH:O	2.47	0.42
1:A:307:PRO:HA	1:A:383:ALA:CB	2.50	0.42
1:A:31:GLN:HE21	1:A:37:PRO:HD3	1.84	0.42
2:B:308:ARG:HG3	2:B:342:TYR:CZ	2.55	0.42
1:A:263:PRO:O	1:A:266:HIS:HD2	2.04	0.41
1:A:192:HIS:CG	1:A:421:ALA:HA	2.56	0.40
1:A:71:GLU:HB3	1:A:98:ASP:HB3	2.03	0.40
2:B:298:SER:H	2:B:308:ARG:HH22	1.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	425/451 (94%)	408 (96%)	17 (4%)	0	100	100
2	B	433/445 (97%)	422 (98%)	10 (2%)	1 (0%)	52	42
3	F	129/169 (76%)	128 (99%)	1 (1%)	0	100	100
All	All	987/1065 (93%)	958 (97%)	28 (3%)	1 (0%)	56	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	276	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/379 (94%)	347 (98%)	9 (2%)	55	47
2	B	377/383 (98%)	362 (96%)	15 (4%)	38	26
3	F	99/131 (76%)	97 (98%)	2 (2%)	63	57
All	All	832/893 (93%)	806 (97%)	26 (3%)	47	37

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	84	ARG
1	A	181	VAL
1	A	280	LYS
1	A	297	GLU
1	A	302[A]	MET
1	A	302[B]	MET
1	A	335	ILE
1	A	349	THR
2	B	48	ARG
2	B	57	THR
2	B	75	MET
2	B	139	HIS
2	B	207	GLU
2	B	221	THR
2	B	247	GLN
2	B	275	LEU
2	B	276	THR
2	B	278	ARG
2	B	295	MET
2	B	325	MET
2	B	338	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	371	LEU
2	B	405	LEU
3	F	20	GLU
3	F	127	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	101	ASN
1	A	107	HIS
1	A	266	HIS
1	A	285	GLN
1	A	301	GLN
2	B	167	ASN
2	B	192	HIS
2	B	247	GLN
2	B	300	ASN
3	F	59	HIS
3	F	92	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GTP	A	501	5	26,34,34	1.01	1 (3%)	29,54,54	2.35	4 (13%)
6	SO4	A	503	-	4,4,4	0.07	0	6,6,6	0.08	0
6	SO4	A	504	-	4,4,4	0.19	0	6,6,6	0.10	0
7	GOL	A	505	-	5,5,5	0.14	0	5,5,5	0.19	0
7	GOL	A	506	-	5,5,5	0.13	0	5,5,5	0.34	0
7	GOL	A	507	-	5,5,5	0.10	0	5,5,5	0.22	0
8	GDP	B	501	-	24,30,30	1.06	2 (8%)	26,47,47	2.51	4 (15%)
9	LOC	B	502	-	29,31,31	0.98	2 (6%)	27,44,44	0.53	0
6	SO4	B	503	-	4,4,4	0.16	0	6,6,6	0.09	0
7	GOL	B	504	-	5,5,5	0.12	0	5,5,5	0.29	0
7	GOL	B	505	-	5,5,5	0.13	0	5,5,5	0.51	0
7	GOL	F	201	-	5,5,5	0.12	0	5,5,5	0.22	0
7	GOL	F	202	-	5,5,5	0.11	0	5,5,5	0.23	0
7	GOL	F	203	-	5,5,5	0.11	0	5,5,5	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	A	501	5	-	0/18/38/38	0/3/3/3
6	SO4	A	503	-	-	0/0/0/0	0/0/0/0
6	SO4	A	504	-	-	0/0/0/0	0/0/0/0
7	GOL	A	505	-	-	0/4/4/4	0/0/0/0
7	GOL	A	506	-	-	0/4/4/4	0/0/0/0
7	GOL	A	507	-	-	0/4/4/4	0/0/0/0
8	GDP	B	501	-	-	0/12/32/32	0/3/3/3
9	LOC	B	502	-	-	0/10/25/25	0/3/3/3
6	SO4	B	503	-	-	0/0/0/0	0/0/0/0
7	GOL	B	504	-	-	0/4/4/4	0/0/0/0
7	GOL	B	505	-	-	0/4/4/4	0/0/0/0
7	GOL	F	201	-	-	0/4/4/4	0/0/0/0
7	GOL	F	202	-	-	0/4/4/4	0/0/0/0
7	GOL	F	203	-	-	0/4/4/4	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	501	GDP	C6-C5	2.25	1.45	1.41
9	B	502	LOC	C15-C16	2.48	1.44	1.39
9	B	502	LOC	C14-C11	2.95	1.57	1.52
4	A	501	GTP	C6-N1	3.11	1.38	1.33
8	B	501	GDP	C6-N1	3.34	1.39	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	501	GDP	C5-C6-N1	-8.59	112.29	123.52
4	A	501	GTP	C5-C6-N1	-8.47	112.44	123.52
4	A	501	GTP	N3-C2-N1	-3.08	123.37	127.56
8	B	501	GDP	N3-C2-N1	-3.06	123.39	127.56
8	B	501	GDP	C6-C5-C4	-2.84	117.61	120.86
4	A	501	GTP	C6-C5-C4	-2.40	118.11	120.86
8	B	501	GDP	C6-N1-C2	7.97	125.23	115.88
4	A	501	GTP	C6-N1-C2	7.98	125.23	115.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	506	GOL	1	0
9	B	502	LOC	1	0
7	F	203	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/451 (94%)	0.50	44 (10%) 9 9	30, 49, 86, 106	0
2	B	425/445 (95%)	0.21	20 (4%) 35 38	28, 41, 67, 120	0
3	F	130/169 (76%)	0.16	9 (6%) 20 22	32, 46, 83, 123	0
All	All	983/1065 (92%)	0.33	73 (7%) 17 19	28, 45, 81, 123	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	277	SER	9.2
2	B	285	ALA	8.7
1	A	282	TYR	7.8
3	F	140	ASN	7.4
2	B	276	THR	6.6
1	A	283	HIS	5.8
3	F	141	ALA	5.2
1	A	281	ALA	5.1
1	A	59	GLY	4.9
3	F	139	VAL	4.4
1	A	365	GLY	4.3
1	A	279	GLU	4.2
2	B	75	MET	4.0
1	A	338	LYS	4.0
2	B	286	LEU	3.8
1	A	58	ALA	3.8
3	F	142	GLN	3.7
1	A	35	GLN	3.6
1	A	49	PHE	3.5
1	A	53	PHE	3.5
1	A	57	GLY	3.4
1	A	61	HIS	3.4
1	A	32	PRO	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	370	GLY	3.4
1	A	1	MET	3.4
3	F	32	VAL	3.3
2	B	94	PHE	3.3
1	A	37	PRO	3.3
1	A	244	PHE	3.2
1	A	89	PRO	3.2
1	A	85	GLN	3.2
1	A	83	TYR	3.1
1	A	346	TRP	3.1
1	A	437	VAL	3.1
1	A	48	SER	3.0
2	B	441	ASP	3.0
1	A	56	THR	3.0
1	A	82	THR	3.0
3	F	36	ASN	3.0
1	A	31	GLN	3.0
1	A	26	LEU	2.9
1	A	335	ILE	2.9
2	B	278	ARG	2.9
1	A	33	ASP	2.8
1	A	348	PRO	2.8
2	B	416	MET	2.8
1	A	50	ASN	2.8
3	F	37	GLY	2.6
1	A	284	GLU	2.5
1	A	372	GLN	2.5
1	A	345	ASP	2.4
1	A	60	LYS	2.4
2	B	247	GLN	2.4
2	B	72	PRO	2.4
2	B	322	ARG	2.3
1	A	36	MET	2.3
1	A	391	LEU	2.3
2	B	248	LEU	2.2
1	A	339	ARG	2.2
1	A	340	SER	2.2
2	B	275	LEU	2.2
2	B	287	THR	2.1
3	F	35	ALA	2.1
2	B	58	GLY	2.1
3	F	33	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	55	GLU	2.1
1	A	367	ASP	2.1
1	A	358	GLN	2.1
2	B	242	LEU	2.1
1	A	131	GLY	2.1
2	B	250	ALA	2.0
1	A	47	ASP	2.0
2	B	324	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	GOL	A	507	6/6	0.68	0.24	18.35	76,80,81,82	0
7	GOL	B	505	6/6	0.70	0.39	9.69	69,78,80,82	0
7	GOL	A	506	6/6	0.52	0.27	7.47	65,69,70,71	0
7	GOL	F	202	6/6	0.90	0.16	3.33	50,59,62,64	0
7	GOL	A	505	6/6	0.80	0.21	2.07	49,53,56,60	0
6	SO4	A	504	5/5	0.66	0.23	1.72	116,117,117,118	0
7	GOL	B	504	6/6	0.69	0.21	1.17	65,68,69,69	0
7	GOL	F	201	6/6	0.97	0.11	0.42	45,46,48,51	0
4	GTP	A	501	32/32	0.98	0.12	-0.59	31,35,41,43	0
9	LOC	B	502	29/29	0.95	0.12	-0.61	28,33,38,39	0
8	GDP	B	501	28/28	0.98	0.09	-0.71	27,31,32,33	0
5	MG	A	502	1/1	0.97	0.12	-0.79	68,68,68,68	0
6	SO4	A	503	5/5	0.97	0.12	-	79,80,81,81	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	GOL	F	203	6/6	0.69	0.18	-	83,84,84,85	0
6	SO4	B	503	5/5	0.91	0.16	-	121,122,122,122	0

6.5 Other polymers [i](#)

There are no such residues in this entry.