



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 4, 2017 – 07:27 PM EST

PDB ID : 5EZY  
Title : Crystal structure of T2R-TTL-taccalonolide AJ complex  
Authors : Wang, Y.; Yu, Y.; Chen, Q.; Yang, J.  
Deposited on : 2015-11-27  
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1192 (2.04-2.04)
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	<div> <div>2%</div> <div>95%</div> <div>• •</div> </div>
1	C	450	<div> <div>%</div> <div>94%</div> <div>• •</div> </div>
2	B	445	<div> <div>4%</div> <div>93%</div> <div>• •</div> </div>
2	D	445	<div> <div>5%</div> <div>93%</div> <div>• •</div> </div>
3	E	143	<div> <div>6%</div> <div>84%</div> <div>• 15%</div> </div>
4	F	384	<div> <div>17%</div> <div>85%</div> <div>5% • 10%</div> </div>

## 2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 36311 atoms, of which 16959 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	437	Total	C	H	N	O	S	0	3	0
			6756	2169	3328	583	652	24			
1	C	440	Total	C	H	N	O	S	0	9	0
			6836	2192	3368	589	662	25			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	427	Total	C	H	N	O	S	0	0	0
			6588	2110	3227	576	649	26			
2	D	431	Total	C	H	N	O	S	0	0	0
			6645	2126	3256	580	656	27			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	E	121	Total	C	H	N	O	S	0	4	0
			2046	627	1028	186	200	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	-	cloning artifact	UNP P63043
E	4	ALA	-	cloning artifact	UNP P63043


- Molecule 4 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	F	347	Total	C	H	N	O	S	0	1	0
			5546	1832	2686	494	520	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	P	0	0
			42	10	10	5	14	3		
5	C	1	Total	C	H	N	O	P	0	0
			42	10	10	5	14	3		
5	D	1	Total	C	H	N	O	P	0	0
			42	10	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	A	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		
7	C	1	Total	Ca	0	0
			1	1		


- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).






Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
9	B	1	Total 25	C 6	H 13	N 1	O 4	S 1	0	0
9	B	1	Total 25	C 6	H 13	N 1	O 4	S 1	0	0

- Molecule 10 is taccalonolide AJ (three-letter code: TAJ) (formula:  $C_{34}H_{46}O_{14}$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total 48	C 34	O 14	0	0
10	D	1	Total 48	C 34	O 14	0	0


- Molecule 11 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (three-letter code: ACP) (formula: C<sub>11</sub>H<sub>18</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>).



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain





- Molecule 1: Tubulin alpha-1B chain




- Molecule 2: Tubulin beta-2B chain




- Molecule 2: Tubulin beta-2B chain





- Molecule 3: Stathmin-4







● Molecule 4: Uncharacterized protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.30Å 158.50Å 180.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.20 – 2.05 45.20 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.20-2.05) 99.5 (45.20-2.05)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.67 (at 2.05Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, $R_{free}$	0.187 , 0.208 0.187 , 0.193	Depositor DCC
$R_{free}$ test set	9234 reflections (4.90%)	DCC
Wilson B-factor (Å <sup>2</sup> )	25.1	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 56.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	36311	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG, CA, GTP, ACP, MES, TAJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/3514	0.42	0/4770
1	C	0.23	0/3577	0.42	0/4857
2	B	0.23	0/3436	0.41	0/4654
2	D	0.23	0/3464	0.40	0/4692
3	E	0.22	0/1044	0.33	0/1385
4	F	0.22	0/2931	0.40	0/3958
All	All	0.23	0/17966	0.40	0/24316

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3428	3328	3329	5	0
1	C	3468	3368	3349	7	0
2	B	3361	3227	3238	6	0
2	D	3389	3256	3266	8	0
3	E	1018	1028	1007	0	0
4	F	2860	2686	2816	21	0
5	A	32	10	12	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	32	10	12	0	0
5	D	32	10	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	1	0	0	0	0
7	C	1	0	0	0	0
8	B	28	10	12	0	0
9	B	24	26	26	0	0
10	B	48	0	0	3	0
10	D	48	0	0	3	0
11	F	31	0	14	5	0
12	A	331	0	0	2	0
12	B	275	0	0	2	0
12	C	454	0	0	2	0
12	D	218	0	0	1	0
12	E	91	0	0	0	0
12	F	178	0	0	1	0
All	All	19352	16959	17093	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:F:150:LYS:HE3	4:F:160:ILE:HD13	1.48	0.94
4:F:148:ILE:HD11	4:F:150:LYS:HE3	1.54	0.88
2:B:424:ASN:ND2	12:B:601:HOH:O	2.21	0.74
4:F:40:MET:SD	12:F:662:HOH:O	2.45	0.73
1:C:128:GLN:NE2	12:C:601:HOH:O	2.23	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/450 (97%)	430 (98%)	8 (2%)	0	100	100
1	C	447/450 (99%)	437 (98%)	10 (2%)	0	100	100
2	B	425/445 (96%)	418 (98%)	7 (2%)	0	100	100
2	D	429/445 (96%)	421 (98%)	8 (2%)	0	100	100
3	E	121/143 (85%)	120 (99%)	1 (1%)	0	100	100
4	F	340/384 (88%)	327 (96%)	13 (4%)	0	100	100
All	All	2200/2317 (95%)	2153 (98%)	47 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/378 (98%)	370 (100%)	1 (0%)	94	95
1	C	380/378 (100%)	373 (98%)	7 (2%)	66	62
2	B	369/383 (96%)	363 (98%)	6 (2%)	70	67
2	D	372/383 (97%)	369 (99%)	3 (1%)	86	86
3	E	113/127 (89%)	112 (99%)	1 (1%)	84	84
4	F	314/342 (92%)	307 (98%)	7 (2%)	60	53
All	All	1919/1991 (96%)	1894 (99%)	25 (1%)	76	74

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	293	ASN
1	C	347[B]	CYS
4	F	248	GLU
1	C	347[A]	CYS
2	D	19	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 6 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GTP	A	501	6	26,34,34	1.04	1 (3%)	26,54,54	0.87	1 (3%)
8	GDP	B	501	6	24,30,30	1.42	4 (16%)	23,47,47	0.79	1 (4%)
9	MES	B	503	-	12,12,12	1.95	1 (8%)	16,16,16	1.93	5 (31%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
9	MES	B	504	-	12,12,12	2.04	1 (8%)	16,16,16	1.35	2 (12%)
10	TAJ	B	505	2	52,54,54	2.71	10 (19%)	57,94,94	2.27	17 (29%)
5	GTP	C	501	6	26,34,34	1.02	1 (3%)	26,54,54	0.81	0
5	GTP	D	501	6	26,34,34	1.05	1 (3%)	26,54,54	0.91	1 (3%)
10	TAJ	D	503	2	52,54,54	2.49	8 (15%)	57,94,94	2.40	18 (31%)
11	ACP	F	401	-	29,33,33	2.12	11 (37%)	29,52,52	1.63	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	A	501	6	-	0/18/38/38	0/3/3/3
8	GDP	B	501	6	-	0/12/32/32	0/3/3/3
9	MES	B	503	-	-	0/6/14/14	0/1/1/1
9	MES	B	504	-	-	0/6/14/14	0/1/1/1
10	TAJ	B	505	2	-	0/12/145/145	0/0/7/7
5	GTP	C	501	6	-	0/18/38/38	0/3/3/3
5	GTP	D	501	6	-	0/18/38/38	0/3/3/3
10	TAJ	D	503	2	-	0/12/145/145	0/0/7/7
11	ACP	F	401	-	-	0/15/38/38	0/3/3/3

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	B	505	TAJ	CAT-CAS	-14.18	1.38	1.52
10	D	503	TAJ	CAT-CAS	-12.14	1.40	1.52
10	D	503	TAJ	CAM-CAS	-7.53	1.38	1.51
10	B	505	TAJ	CAM-CAS	-7.49	1.38	1.51
9	B	504	MES	C8-S	-6.19	1.66	1.78

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	B	505	TAJ	OBS-CBI-CBH	-7.75	120.87	128.24
10	B	505	TAJ	CAX-CBC-CBB	-7.64	98.19	108.00
10	D	503	TAJ	OBS-CBI-CBH	-7.60	121.01	128.24
10	D	503	TAJ	CAX-CBC-CBB	-7.30	98.61	108.00
10	D	503	TAJ	CAI-CAM-CAS	-6.60	105.03	113.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	B	505	TAJ	3	0
10	D	503	TAJ	3	0
11	F	401	ACP	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	437/450 (97%)	-0.11	11 (2%) 61 67	14, 26, 51, 64	0
1	C	440/450 (97%)	-0.27	4 (0%) 85 89	11, 20, 45, 70	0
2	B	427/445 (95%)	0.08	16 (3%) 45 51	13, 29, 59, 90	0
2	D	431/445 (96%)	0.27	24 (5%) 28 32	14, 32, 69, 128	0
3	E	121/143 (84%)	0.28	8 (6%) 22 24	17, 39, 67, 80	0
4	F	347/384 (90%)	0.70	66 (19%) 2 1	19, 40, 90, 115	0
All	All	2203/2317 (95%)	0.12	129 (5%) 26 29	11, 30, 68, 128	0

The worst 5 of 129 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	283	TYR	9.2
2	B	57	THR	8.1
2	D	284	ARG	7.8
4	F	142	ARG	7.6
2	D	285	ALA	7.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	MG	C	502	1/1	0.95	0.18	1.99	15,15,15,15	0
9	MES	B	504	12/12	0.92	0.17	1.93	35,44,52,54	0
6	MG	A	502	1/1	0.97	0.21	1.93	18,18,18,18	0
5	GTP	C	501	32/32	0.98	0.15	1.36	11,14,17,17	0
10	TAJ	B	505	48/48	0.80	0.19	1.05	53,62,71,79	0
5	GTP	A	501	32/32	0.97	0.19	0.75	14,18,21,24	0
9	MES	B	503	12/12	0.93	0.13	0.66	28,38,56,68	0
8	GDP	B	501	28/28	0.96	0.14	0.62	14,20,26,26	0
10	TAJ	D	503	48/48	0.88	0.14	-0.16	44,57,71,77	0
5	GTP	D	501	32/32	0.96	0.10	-0.46	20,27,37,40	0
11	ACP	F	401	31/31	0.93	0.12	-0.57	37,50,86,90	0
7	CA	A	503	1/1	0.95	0.05	-2.17	37,37,37,37	0
6	MG	D	502	1/1	0.98	0.05	-2.17	33,33,33,33	0
7	CA	C	503	1/1	0.99	0.04	-3.96	26,26,26,26	0
6	MG	B	502	1/1	0.99	0.15	-	11,11,11,11	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.