



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:35 AM GMT

PDB ID : 3FB6
Title : KcsA Potassium channel in the partially open state with 16 Å opening at T112
Authors : Cuello, L.G.; Jogini, V.; Cortes, D.M.; Perozo, E.
Deposited on : 2008-11-18
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	<div> <div>3%</div> <div>78%</div> <div>21%</div> <div>.</div> </div>
2	B	212	<div> <div>2%</div> <div>73%</div> <div>26%</div> </div>
3	C	104	<div> <div>%</div> <div>67%</div> <div>19%</div> <div>13%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody fab fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1648	1042	275	325	6			

- Molecule 2 is a protein called antibody fab fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	212	Total	C	N	O	S	0	0	0
			1649	1023	283	338	5			

- Molecule 3 is a protein called Voltage-gated potassium channel.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	90	Total	C	N	O	S	0	0	0
			632	415	103	112	2			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	25	GLN	HIS	ENGINEERED	UNP P0A334
C	90	CYS	LEU	ENGINEERED	UNP P0A334
C	117	GLN	ARG	ENGINEERED	UNP P0A334
C	120	GLN	GLU	ENGINEERED	UNP P0A334
C	121	GLN	ARG	ENGINEERED	UNP P0A334
C	122	GLN	ARG	ENGINEERED	UNP P0A334
C	124	GLN	HIS	ENGINEERED	UNP P0A334


- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	5	Total	K	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody fab fragment heavy chain



- Molecule 2: antibody fab fragment light chain



- Molecule 3: Voltage-gated potassium channel



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	155.71Å 155.71Å 74.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 3.00 36.70 – 3.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-3.00) 95.4 (36.70-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.96 (at 3.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.248 , 0.255 0.252 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	89.8	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 66.9	EDS
Estimated twinning fraction	0.028 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 17778 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3934	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.34	0/1692	0.66	0/2312
2	B	0.34	0/1686	0.64	0/2287
3	C	0.39	0/645	0.69	0/891
All	All	0.35	0/4023	0.66	0/5490

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1648	0	1620	41	3
2	B	1649	0	1580	57	3
3	C	632	0	635	17	0
4	C	5	0	0	0	0
All	All	3934	0	3835	110	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:134:CYS:HG	2:B:194:CYS:HG	0.94	0.87
2:B:167:ASP:OD2	2:B:169:LYS:HB2	1.79	0.83
1:A:65:GLN:H	1:A:65:GLN:NE2	1.79	0.80
1:A:120:LYS:H	1:A:120:LYS:HE3	1.49	0.78
1:A:22:CYS:HG	1:A:96:CYS:HG	0.88	0.77
2:B:21:PHE:HE1	2:B:75:ILE:HD13	1.56	0.70
2:B:183:LYS:O	2:B:187:GLU:HG2	1.91	0.70
2:B:74:SER:C	2:B:75:ILE:HD12	2.12	0.69
3:C:86:LEU:HD12	3:C:86:LEU:O	1.93	0.69
1:A:124:PRO:HB3	1:A:150:TYR:HB3	1.74	0.68
2:B:2:ILE:HD11	2:B:93:ARG:HB2	1.75	0.67
3:C:91:VAL:O	3:C:95:VAL:HG23	1.96	0.66
2:B:183:LYS:HB3	2:B:183:LYS:NZ	2.11	0.65
2:B:58:ILE:N	2:B:58:ILE:HD12	2.11	0.65
1:A:31:SER:HB2	3:C:62:TYR:CE1	2.31	0.65
2:B:61:ARG:HG3	2:B:76:ASN:O	1.96	0.64
1:A:148:LYS:HB3	1:A:181:THR:HG23	1.79	0.64
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.79	0.63
1:A:120:LYS:HE3	1:A:120:LYS:N	2.14	0.62
1:A:51:ILE:HD13	1:A:52:ILE:C	2.21	0.61
2:B:21:PHE:HE1	2:B:75:ILE:CD1	2.13	0.61
2:B:92:ASN:HD21	3:C:58:GLN:NE2	1.98	0.61
1:A:120:LYS:H	1:A:120:LYS:CE	2.13	0.60
3:C:89:ARG:O	3:C:93:VAL:HG23	2.01	0.60
2:B:2:ILE:CD1	2:B:93:ARG:HB2	2.32	0.59
2:B:75:ILE:N	2:B:75:ILE:HD12	2.18	0.59
2:B:13:VAL:CG2	2:B:17:GLU:HB3	2.31	0.59
3:C:44:SER:OG	3:C:66:LEU:HA	2.03	0.58
2:B:48:ILE:HD12	2:B:73:LEU:CD1	2.33	0.58
2:B:55:ILE:HB	2:B:58:ILE:HD13	1.86	0.57
1:A:34:ILE:HD13	1:A:79:ALA:HB2	1.86	0.57
1:A:6:GLN:HE21	1:A:109:GLY:HA3	1.69	0.57
2:B:124:GLN:HG2	2:B:129:GLY:O	2.05	0.57
1:A:23:LYS:HD3	1:A:78:THR:CG2	2.35	0.57
2:B:2:ILE:H	2:B:2:ILE:HD12	1.70	0.56
2:B:2:ILE:N	2:B:2:ILE:HD12	2.21	0.56
1:A:64:ILE:HD12	1:A:65:GLN:O	2.06	0.56
1:A:215:ILE:HD12	1:A:215:ILE:N	2.21	0.55
2:B:85:ASN:ND2	2:B:103:LYS:HD3	2.22	0.55
2:B:21:PHE:CE1	2:B:75:ILE:HD13	2.41	0.54
1:A:101:GLY:HA3	3:C:62:TYR:CE1	2.43	0.53
2:B:19:VAL:HG12	2:B:75:ILE:HB	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:LEU:HD12	1:A:182:LEU:C	2.29	0.53
2:B:2:ILE:HD11	2:B:93:ARG:HD2	1.90	0.53
1:A:51:ILE:HD13	1:A:52:ILE:N	2.24	0.52
3:C:29:ALA:C	3:C:31:ALA:H	2.11	0.52
1:A:83:LEU:HB3	1:A:86:LEU:HD21	1.92	0.52
1:A:140:MET:HE3	1:A:187:THR:HG22	1.90	0.52
2:B:18:ARG:NH1	2:B:76:ASN:OD1	2.43	0.51
2:B:93:ARG:HG2	3:C:58:GLN:HG2	1.91	0.51
2:B:58:ILE:N	2:B:58:ILE:CD1	2.72	0.50
1:A:23:LYS:HD3	1:A:78:THR:HG23	1.94	0.50
1:A:65:GLN:H	1:A:65:GLN:HE21	1.58	0.50
3:C:51:GLU:HG3	3:C:59:LEU:HB3	1.94	0.50
1:A:17:SER:CB	1:A:84:SER:HA	2.42	0.50
2:B:79:GLU:HG3	2:B:81:GLU:HG2	1.95	0.49
2:B:161:ASN:HB3	2:B:175:MET:HE3	1.95	0.49
2:B:115:VAL:HG22	2:B:136:LEU:CD1	2.43	0.49
2:B:134:CYS:HG	2:B:194:CYS:CB	2.26	0.48
2:B:142:LYS:HB3	2:B:173:TYR:CG	2.49	0.48
2:B:136:LEU:HD22	2:B:136:LEU:N	2.29	0.48
1:A:148:LYS:NZ	1:A:148:LYS:HB2	2.29	0.47
1:A:1:GLN:N	1:A:1:GLN:OE1	2.42	0.47
2:B:150:ILE:HD12	2:B:155:ARG:HH11	1.78	0.47
2:B:13:VAL:HG22	2:B:14:SER:N	2.29	0.47
3:C:86:LEU:C	3:C:86:LEU:HD12	2.36	0.46
1:A:12:VAL:O	1:A:116:VAL:HA	2.15	0.46
3:C:29:ALA:C	3:C:31:ALA:N	2.69	0.46
2:B:142:LYS:HD3	2:B:173:TYR:CE2	2.50	0.46
1:A:145:CYS:HG	1:A:200:CYS:HG	1.64	0.46
2:B:92:ASN:ND2	3:C:58:GLN:NE2	2.64	0.46
2:B:183:LYS:HZ3	2:B:183:LYS:HB3	1.79	0.46
1:A:34:ILE:CD1	1:A:79:ALA:HB2	2.46	0.45
1:A:198:VAL:HG12	1:A:215:ILE:HD13	1.97	0.45
2:B:83:ILE:HD11	2:B:106:ILE:HD12	1.99	0.45
2:B:55:ILE:HB	2:B:58:ILE:CD1	2.45	0.45
2:B:48:ILE:HD12	2:B:73:LEU:HD13	1.98	0.44
1:A:120:LYS:H	1:A:120:LYS:CD	2.29	0.44
1:A:98:ARG:HD3	1:A:100:ARG:CZ	2.48	0.44
2:B:40:THR:HG22	2:B:41:ASN:ND2	2.33	0.44
3:C:62:TYR:HB2	3:C:63:PRO:HD3	2.00	0.44
1:A:3:GLN:O	1:A:24:ALA:HA	2.18	0.44
2:B:84:ALA:O	2:B:103:LYS:HD2	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:GLN:O	1:A:4:LEU:HD12	2.18	0.43
1:A:52:ILE:O	1:A:52:ILE:HG23	2.18	0.43
2:B:193:THR:HG22	2:B:208:SER:CB	2.49	0.43
2:B:193:THR:HG22	2:B:208:SER:HB3	1.99	0.43
1:A:124:PRO:CB	1:A:150:TYR:HB3	2.46	0.43
1:A:140:MET:CE	1:A:187:THR:HG22	2.48	0.43
2:B:139:PHE:HE1	2:B:142:LYS:HA	1.83	0.43
2:B:75:ILE:CD1	2:B:75:ILE:N	2.81	0.42
1:A:35:HIS:CG	1:A:105:PHE:HE2	2.37	0.42
2:B:118:PHE:HA	2:B:119:PRO:HD3	1.87	0.42
2:B:61:ARG:NH1	2:B:82:ASP:OD1	2.53	0.42
2:B:83:ILE:HD13	2:B:166:GLN:HG2	2.01	0.42
2:B:21:PHE:CE1	2:B:75:ILE:CD1	2.99	0.42
1:A:1:GLN:CD	1:A:1:GLN:N	2.73	0.42
2:B:55:ILE:O	2:B:58:ILE:HD13	2.19	0.42
3:C:96:MET:O	3:C:100:ILE:HG13	2.19	0.42
1:A:128:PRO:HG3	1:A:213:LYS:HD2	2.01	0.42
2:B:55:ILE:CB	2:B:58:ILE:HD13	2.50	0.41
1:A:23:LYS:HD3	1:A:78:THR:OG1	2.20	0.41
3:C:31:ALA:O	3:C:34:VAL:HB	2.20	0.41
2:B:13:VAL:HG22	2:B:17:GLU:HB3	1.99	0.41
2:B:136:LEU:CD1	2:B:196:ALA:HB2	2.51	0.41
2:B:13:VAL:HG23	2:B:17:GLU:CD	2.40	0.41
3:C:66:LEU:O	3:C:69:SER:OG	2.38	0.40
2:B:85:ASN:ND2	2:B:103:LYS:CD	2.84	0.40
1:A:91:SER:O	1:A:92:ALA:HB2	2.21	0.40
1:A:40:ARG:NH2	1:A:89:GLU:HB3	2.36	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:GLN:OE1	2:B:18:ARG:NE[3_755]	1.37	0.83
1:A:65:GLN:OE1	2:B:18:ARG:CD[3_755]	1.70	0.50
1:A:65:GLN:OE1	2:B:18:ARG:CZ[3_755]	1.81	0.39

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/219 (99%)	212 (98%)	5 (2%)	0	100	100
2	B	210/212 (99%)	203 (97%)	7 (3%)	0	100	100
3	C	88/104 (85%)	84 (96%)	2 (2%)	2 (2%)	8	36
All	All	515/535 (96%)	499 (97%)	14 (3%)	2 (0%)	39	80

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	77	GLY
3	C	113	TRP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/185 (100%)	178 (96%)	7 (4%)	40	78
2	B	190/190 (100%)	184 (97%)	6 (3%)	46	82
3	C	58/75 (77%)	58 (100%)	0	100	100
All	All	433/450 (96%)	420 (97%)	13 (3%)	48	83

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	65	GLN
1	A	120	LYS
1	A	148	LYS
1	A	152	PRO
1	A	154	PRO
1	A	194	PRO
2	B	4	LEU
2	B	58	ILE
2	B	95	PRO
2	B	165	ASP
2	B	184	ASP
2	B	190	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	5	GLN
1	A	6	GLN
1	A	65	GLN
2	B	41	ASN
2	B	85	ASN
2	B	92	ASN
2	B	137	ASN
2	B	190	ASN
2	B	210	ASN
3	C	58	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	219/219 (100%)	0.10	6 (2%)	58	28	76, 111, 136, 149	0
2	B	212/212 (100%)	0.12	4 (1%)	70	41	67, 108, 143, 158	0
3	C	90/104 (86%)	-0.27	1 (1%)	82	58	56, 74, 120, 138	0
All	All	521/535 (97%)	0.04	11 (2%)	67	36	56, 107, 141, 158	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	157	ASN	4.6
2	B	209	PHE	3.7
2	B	145	ASN	3.5
1	A	27	TYR	3.2
1	A	1	GLN	2.7
1	A	136	GLN	2.6
1	A	219	ASP	2.6
1	A	55	TYR	2.4
3	C	112	THR	2.3
1	A	28	THR	2.2
2	B	192	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	K	C	1	1/1	0.82	0.14	-	100,100,100,100	1
4	K	C	4	1/1	0.97	0.51	-	64,64,64,64	1
4	K	C	2	1/1	0.96	0.63	-	80,80,80,80	1
4	K	C	5	1/1	0.87	0.64	-	129,129,129,129	1
4	K	C	3	1/1	0.86	0.68	-	86,86,86,86	1

6.5 Other polymers [i](#)

There are no such residues in this entry.