



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:38 AM GMT

PDB ID : 3FFS
Title : The Crystal Structure of Cryptosporidium parvum Inosine-5'-Monophosphate Dehydrogenase
Authors : Riera, T.V.; D'Aquino, J.A.; Lu, J.; Petsko, G.A.; Hedstrom, L.
Deposited on : 2008-12-04
Resolution : 3.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	400	<div> <div>54%</div> <div>18%</div> <div>•</div> <div>25%</div> </div>
1	B	400	<div> <div>53%</div> <div>19%</div> <div>•</div> <div>24%</div> </div>
1	C	400	<div> <div>4%</div> <div>49%</div> <div>21%</div> <div>•</div> <div>26%</div> </div>
1	D	400	<div> <div>4%</div> <div>51%</div> <div>21%</div> <div>• •</div> <div>25%</div> </div>


2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 8561 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inosine-5-monophosphate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	0	0	0
			2188	1388	371	418	11			
1	B	303	Total	C	N	O	S	0	0	0
			2190	1383	364	433	10			
1	C	298	Total	C	N	O	S	0	0	0
			2140	1348	363	418	11			
1	D	299	Total	C	N	O	S	0	0	0
			2043	1290	346	397	10			



• Molecule 1: Inosine-5-monophosphate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	119.07Å 153.32Å 98.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.34 – 3.19 48.32 – 3.19	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.34-3.19) 99.1 (48.32-3.19)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.269 , 0.328 0.272 , 0.326	Depositor DCC
R_{free} test set	1534 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	74.5	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 38.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 30483 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	8561	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/2206	0.71	1/2976 (0.0%)
1	B	0.52	0/2211	0.71	0/2994
1	C	0.54	0/2155	0.70	0/2909
1	D	0.56	0/2060	0.69	1/2803 (0.0%)
All	All	0.54	0/8632	0.70	2/11682 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	162	LEU	CA-CB-CG	6.58	130.44	115.30
1	D	129	LEU	CA-CB-CG	5.26	127.40	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2188	0	2259	49	0
1	B	2190	0	2189	66	0
1	C	2140	0	2157	57	0
1	D	2043	0	1945	50	0
All	All	8561	0	8550	210	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 210 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:25:LEU:HD13	1:C:26:PRO:HD2	1.37	1.05
1:B:15:ILE:HD11	1:B:232:ILE:HG21	1.49	0.95
1:C:185:ILE:HG13	1:C:186:ASP:H	1.32	0.94
1:C:334:ARG:HG2	1:C:334:ARG:HH11	1.39	0.87
1:B:21:TYR:HE2	1:B:23:GLU:HG3	1.40	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/400 (73%)	262 (90%)	25 (9%)	4 (1%)	14	57
1	B	295/400 (74%)	262 (89%)	25 (8%)	8 (3%)	6	39
1	C	286/400 (72%)	231 (81%)	43 (15%)	12 (4%)	3	26
1	D	287/400 (72%)	223 (78%)	46 (16%)	18 (6%)	2	13
All	All	1159/1600 (72%)	978 (84%)	139 (12%)	42 (4%)	4	30

5 of 42 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	244	LYS
1	B	143	VAL
1	B	184	ASN
1	B	185	ILE
1	B	378	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/334 (70%)	197 (84%)	36 (16%)	3	15
1	B	230/334 (69%)	204 (89%)	26 (11%)	7	31
1	C	224/334 (67%)	191 (85%)	33 (15%)	4	18
1	D	194/334 (58%)	170 (88%)	24 (12%)	6	27
All	All	881/1336 (66%)	762 (86%)	119 (14%)	5	22

5 of 119 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	209	ILE
1	C	25	LEU
1	D	204	ASN
1	B	249	ILE
1	B	366	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	231	GLN
1	C	5	ASN
1	C	231	GLN
1	B	171	ASN
1	C	204	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	301/400 (75%)	-0.02	0	100 100	24, 50, 65, 82	0
1	B	303/400 (75%)	-0.08	0	100 100	25, 53, 72, 81	0
1	C	298/400 (74%)	0.03	3 (1%)	84 75	26, 55, 68, 81	0
1	D	299/400 (74%)	0.32	16 (5%)	29 17	25, 56, 73, 83	0
All	All	1201/1600 (75%)	0.06	19 (1%)	74 62	24, 54, 70, 83	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	190	GLY	4.7
1	D	189	VAL	4.0
1	D	187	VAL	3.5
1	D	163	ASP	2.9
1	D	162	LEU	2.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.