



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 10, 2016 – 11:31 PM EDT

PDB ID : 5FGS
Title : Crystal structure of C-terminal domain of shaft pilin spaA from *Lactobacillus rhamnosus* GG - P21212 space group
Authors : Chaurasia, P.; Pratap, S.; von Ossowski, I.; Palva, A.; Krishnan, V.
Deposited on : 2015-12-21
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027790
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027790


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 87%, yellow 87%, yellow 10%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 87% 10% </div> </div>
1	B	136	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 4%, green 4%, green 83%, yellow 83%, yellow 7%, yellow 10%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 4% 83% 7% • 10% </div> </div>
1	C	136	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, green 3%, green 83%, yellow 83%, yellow 8%, yellow 9%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 3% 83% 8% 9% </div> </div>
1	D	136	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 12%, green 12%, green 81%, yellow 81%, yellow 9%, yellow 10%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 12% 81% 9% • 10% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3993 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell surface protein SpaA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	123	Total	C	N	O	0	0	0
			931	580	149	202			
1	B	123	Total	C	N	O	0	0	0
			931	580	149	202			
1	C	124	Total	C	N	O	0	0	0
			935	582	150	203			
1	D	123	Total	C	N	O	0	0	0
			931	580	149	202			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	303	LEU	-	expression tag	UNP C7T9P4
A	304	GLU	-	expression tag	UNP C7T9P4
A	305	HIS	-	expression tag	UNP C7T9P4
A	306	HIS	-	expression tag	UNP C7T9P4
A	307	HIS	-	expression tag	UNP C7T9P4
A	308	HIS	-	expression tag	UNP C7T9P4
A	309	HIS	-	expression tag	UNP C7T9P4
A	310	HIS	-	expression tag	UNP C7T9P4
B	303	LEU	-	expression tag	UNP C7T9P4
B	304	GLU	-	expression tag	UNP C7T9P4
B	305	HIS	-	expression tag	UNP C7T9P4
B	306	HIS	-	expression tag	UNP C7T9P4
B	307	HIS	-	expression tag	UNP C7T9P4
B	308	HIS	-	expression tag	UNP C7T9P4
B	309	HIS	-	expression tag	UNP C7T9P4
B	310	HIS	-	expression tag	UNP C7T9P4
C	303	LEU	-	expression tag	UNP C7T9P4
C	304	GLU	-	expression tag	UNP C7T9P4
C	305	HIS	-	expression tag	UNP C7T9P4
C	306	HIS	-	expression tag	UNP C7T9P4
C	307	HIS	-	expression tag	UNP C7T9P4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	308	HIS	-	expression tag	UNP C7T9P4
C	309	HIS	-	expression tag	UNP C7T9P4
C	310	HIS	-	expression tag	UNP C7T9P4
D	303	LEU	-	expression tag	UNP C7T9P4
D	304	GLU	-	expression tag	UNP C7T9P4
D	305	HIS	-	expression tag	UNP C7T9P4
D	306	HIS	-	expression tag	UNP C7T9P4
D	307	HIS	-	expression tag	UNP C7T9P4
D	308	HIS	-	expression tag	UNP C7T9P4
D	309	HIS	-	expression tag	UNP C7T9P4
D	310	HIS	-	expression tag	UNP C7T9P4

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	3	Total Zn 3 3	0	0
2	A	3	Total Zn 3 3	0	0
2	D	3	Total Zn 3 3	0	0



- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	99	Total O 99 99	0	0
3	B	68	Total O 68 68	0	0
3	C	56	Total O 56 56	0	0
3	D	33	Total O 33 33	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Cell surface protein SpaA





- Molecule 1: Cell surface protein SpaA



- Molecule 1: Cell surface protein SpaA



- Molecule 1: Cell surface protein SpaA



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	57.07Å 74.34Å 116.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.17 – 2.00 37.17 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (37.17-2.00) 99.2 (37.17-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.217 , 0.250 0.223 , 0.259	Depositor DCC
R_{free} test set	1697 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3993	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/946	0.62	0/1281
1	B	0.46	0/946	0.60	0/1281
1	C	0.43	0/950	0.63	0/1286
1	D	0.37	0/946	0.58	0/1281
All	All	0.44	0/3788	0.61	0/5129

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	931	0	875	2	0
1	B	931	0	874	5	0
1	C	935	0	875	6	0
1	D	931	0	874	9	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
2	D	3	0	0	0	0
3	A	99	0	0	0	1
3	B	68	0	0	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	56	0	0	0	0
3	D	33	0	0	0	0
All	All	3993	0	3498	20	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:222:GLU:OE2	1:D:233:ARG:NH2	2.24	0.70
1:B:260:ASP:OD2	1:B:263:THR:OG1	2.13	0.67
1:A:260:ASP:OD2	1:A:263:THR:OG1	2.13	0.65
1:C:260:ASP:OD2	1:C:263:THR:OG1	2.13	0.65
1:C:201:SER:HB3	1:C:207:PHE:CE2	2.36	0.60
1:D:263:THR:CG2	1:D:265:TYR:CZ	2.85	0.60
1:B:201:SER:HB3	1:B:207:PHE:CE2	2.39	0.58
1:A:201:SER:HB3	1:A:207:PHE:CE2	2.39	0.58
1:D:201:SER:HB3	1:D:207:PHE:CE2	2.39	0.58
1:D:189:THR:OG1	1:D:191:GLU:HG2	2.04	0.57
1:C:189:THR:OG1	1:C:191:GLU:HG2	2.06	0.56
1:C:271:ASN:HA	1:D:237:VAL:HG12	1.91	0.51
1:B:283:ASP:N	1:B:283:ASP:OD1	2.47	0.45
1:D:176:THR:HG23	1:D:176:THR:O	2.17	0.45
1:B:213:LYS:HE3	1:B:231:ASN:O	2.17	0.44
1:D:263:THR:HG22	1:D:265:TYR:CZ	2.52	0.44
1:B:176:THR:O	1:B:176:THR:HG23	2.19	0.43
1:D:194:GLU:OE1	1:D:249:LYS:HD3	2.20	0.40
1:C:194:GLU:OE1	1:C:249:LYS:HD3	2.21	0.40
1:D:188:GLU:HG2	1:D:189:THR:HG23	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:526:HOH:O	3:B:557:HOH:O[2_655]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	121/136 (89%)	119 (98%)	2 (2%)	0	100	100
1	B	121/136 (89%)	119 (98%)	2 (2%)	0	100	100
1	C	122/136 (90%)	121 (99%)	1 (1%)	0	100	100
1	D	121/136 (89%)	119 (98%)	2 (2%)	0	100	100
All	All	485/544 (89%)	478 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/113 (89%)	100 (99%)	1 (1%)	82	85
1	B	101/113 (89%)	98 (97%)	3 (3%)	48	47
1	C	101/113 (89%)	100 (99%)	1 (1%)	82	85
1	D	101/113 (89%)	99 (98%)	2 (2%)	63	65
All	All	404/452 (89%)	397 (98%)	7 (2%)	68	71

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	277	ASP
1	B	277	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	283	ASP
1	B	298	SER
1	C	277	ASP
1	D	263	THR
1	D	277	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	123/136 (90%)	0.21	2 (1%) 74 75	20, 29, 50, 61	0
1	B	123/136 (90%)	0.33	5 (4%) 41 42	20, 37, 64, 78	0
1	C	124/136 (91%)	0.39	4 (3%) 51 52	32, 46, 64, 70	0
1	D	123/136 (90%)	0.90	16 (13%) 5 5	37, 55, 76, 85	0
All	All	493/544 (90%)	0.46	27 (5%) 29 30	20, 43, 70, 85	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	280	ALA	4.8
1	D	262	THR	3.3
1	D	190	LYS	3.1
1	D	191	GLU	3.1
1	D	188	GLU	3.0
1	D	216	GLN	3.0
1	D	231	ASN	3.0
1	A	176	THR	2.8
1	C	261	ASN	2.7
1	D	260	ASP	2.7
1	B	263	THR	2.6
1	C	176	THR	2.6
1	D	263	THR	2.5
1	D	213	LYS	2.5
1	D	205	GLY	2.4
1	C	262	THR	2.4
1	D	264	THR	2.3
1	D	249	LYS	2.2
1	D	204	ASP	2.2
1	A	298	SER	2.2
1	B	259	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	265	TYR	2.1
1	C	264	THR	2.1
1	D	215	GLY	2.1
1	B	298	SER	2.1
1	B	178	TYR	2.1
1	B	177	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ZN	A	401	1/1	0.98	0.06	-1.88	37,37,37,37	0
2	ZN	A	403	1/1	0.95	0.04	-3.67	79,79,79,79	0
2	ZN	D	402	1/1	0.95	0.05	-4.79	58,58,58,58	0
2	ZN	D	403	1/1	0.96	0.11	-	73,73,73,73	0
2	ZN	A	402	1/1	0.92	0.09	-	82,82,82,82	0
2	ZN	B	402	1/1	0.97	0.08	-	45,45,45,45	0
2	ZN	D	401	1/1	0.89	0.07	-	80,80,80,80	0
2	ZN	B	401	1/1	0.91	0.07	-	92,92,92,92	0
2	ZN	B	403	1/1	0.76	0.25	-	86,86,86,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.