Full wwPDB NMR Structure Validation Report (i)

Apr 26, 2016 – 02:27 PM BST

PDB ID	:	1FJB
Title	:	NMR Study of an 11-Mer DNA Duplex Containing 7,8-Dihydro-8-Oxoadenine (AOXO) Opposite Thymine
Authors	:	Chen, H.; Johnson, F.; Grollman, A.P.; Patel, D.J.
Deposited on	:	1995-12-15

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	11	73% 27%
2	B	11	9% 82% 9%

2 Ensemble composition and analysis (i)

This entry contains 2 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 697 atoms, of which 251 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*CP*AP*TP*GP*TP*GP*TP*AP*CP*G)-3').

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	11	351	108	126	42	65	10	0

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*GP*TP*AP*CP*(A38)P*CP*AP*TP*GP*C)-3').

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
2	B	11	346	106	125	41	64	10	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA (5'-D(*GP*CP*AP*TP*GP*TP*GP*TP*AP*CP*G)-3')

Chain A: 73% 27%

G1 C2 A3 T4 G5 T6 G7 T8 A9 C10 G11

- Molecule 2: DNA (5'-D(*CP*GP*TP*AP*CP*(A38)P*CP*AP*TP*GP*C)-3')

Chain B: 9% 82% 9%

C12 G13 T14 A15 C16 A17 C18 A19 T20 G21 C22

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: DNA (5'-D(*GP*CP*AP*TP*GP*TP*GP*TP*AP*CP*G)-3')

Chain A: 73% 27%

G1 C2 A3 T4 G5 T6 G7 T8 A9 C10 G11

- Molecule 2: DNA (5'-D(*CP*GP*TP*AP*CP*(A38)P*CP*AP*TP*GP*C)-3')

Chain B: 9% 82% 9%

C12 G13 T14 A15 C16 A17 C18 A19 T20 G21 C22

4.2.2 Score per residue for model 2

- Molecule 1: DNA (5'-D(*GP*CP*AP*TP*GP*TP*GP*TP*AP*CP*G)-3')

Chain A:





- Molecule 2: DNA (5'-D(*CP*GP*TP*AP*CP*(A38)P*CP*AP*TP*GP*C)-3')

Chain B:





5 Refinement protocol and experimental data overview (i)

Of the ? calculated structures, 2 were deposited, based on the following criterion: *?*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: A38

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.29±0.06	3±1/252 (1.0±0.2%)	2.50±0.03	29±1/388 (7.3±0.1%)
2	B	1.32±0.02	3±1/221 (1.1±0.2%)	2.43±0.03	23±1/336 (6.7±0.1%)
All	All	1.31	10/946 (1.1%)	2.47	102/1448 (7.0%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	8	DT	C5-C7	6.61	1.54	1.50	1	2
1	A	6	DT	C5-C7	6.44	1.53	1.50	1	2
1	A	4	DT	C5-C7	6.28	1.53	1.50	1	1
2	B	14	DT	C5-C7	6.08	1.53	1.50	2	2
2	B	20	DT	C5-C7	5.36	1.53	1.50	2	2
2	B	13	DG	C2'-C1'	-5.27	1.47	1.52	1	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	1	DG	P-O3'-C3'	8.97	130.46	119.70	1	1
2	B	15	DA	O4'-C1'-N9	8.25	113.78	108.00	2	2
2	B	20	DT	C6-C5-C7	-8.21	117.98	122.90	2	2
2	B	12	DC	O4'-C1'-C2'	-8.18	99.35	105.90	2	2
1	A	2	DC	O4'-C1'-C2'	-8.16	99.37	105.90	2	2
2	B	13	DG	O4'-C1'-N9	8.13	113.69	108.00	1	2
1	A	4	DT	C6-C5-C7	-8.08	118.05	122.90	2	2
2	B	21	DG	N7-C8-N9	7.98	117.09	113.10	1	2
1	A	7	DG	N7-C8-N9	7.98	117.09	113.10	1	2
1	A	6	DT	O4'-C1'-C2'	-7.87	99.61	105.90	2	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	4	DT	O4'-C1'-C2'	-7.79	99.67	105.90	1	2
1	A	2	DC	O4'-C1'-N1	7.79	113.45	108.00	2	1
2	B	19	DA	O4'-C1'-N9	7.60	113.32	108.00	2	2
1	A	11	DG	N7-C8-N9	7.40	116.80	113.10	1	2
1	A	6	DT	C6-C5-C7	-7.27	118.54	122.90	2	2
1	A	5	DG	N7-C8-N9	7.14	116.67	113.10	2	2
2	B	14	DT	C6-C5-C7	-7.01	118.69	122.90	2	2
1	A	2	DC	C4'-C3'-C2'	-6.94	96.86	103.10	1	1
1	A	4	DT	O4'-C1'-N1	6.88	112.82	108.00	1	2
1	A	7	DG	O4'-C1'-C2'	-6.83	100.43	105.90	2	2
2	B	18	DC	N1-C2-O2	6.76	122.96	118.90	1	2
1	A	1	DG	N7-C8-N9	6.71	116.45	113.10	2	2
1	A	10	DC	O4'-C1'-C2'	-6.62	100.60	105.90	1	2
2	B	13	DG	N7-C8-N9	6.54	116.37	113.10	2	2
1	A	2	DC	N1-C2-O2	6.52	122.81	118.90	2	2
2	B	16	DC	N1-C2-O2	6.52	122.81	118.90	2	2
1	A	5	DG	O4'-C1'-C2'	-6.45	100.74	105.90	1	1
2	B	18	DC	O4'-C4'-C3'	6.37	109.82	106.00	2	2
1	A	1	DG	C8-N9-C4	-6.29	103.88	106.40	1	2
2	B	13	DG	O4'-C1'-C2'	-6.17	100.96	105.90	2	2
1	A	10	DC	N1-C2-O2	6.17	122.60	118.90	1	2
2	B	22	DC	N1-C2-O2	6.09	122.56	118.90	2	2
1	A	11	DG	C5-N7-C8	-6.01	101.30	104.30	1	2
2	B	21	DG	C5-N7-C8	-6.00	101.30	104.30	1	2
2	B	19	DA	N7-C8-N9	5.98	116.79	113.80	2	2
1	A	7	DG	C5-N7-C8	-5.93	101.34	104.30	1	2
2	B	13	DG	P-O3'-C3'	5.93	126.81	119.70	1	2
2	B	15	DA	P-O3'-C3'	5.92	126.80	119.70	2	2
2	B	20	DT	O4'-C1'-N1	5.86	112.10	108.00	1	1
2	B	12	DC	N1-C2-O2	5.84	122.40	118.90	2	2
2	B	21	DG	C8-N9-C4	-5.80	104.08	106.40	2	2
1	A	7	DG	C8-N9-C4	-5.75	104.10	106.40	1	2
1	A	11	DG	C8-N9-C4	-5.71	104.11	106.40	1	2
1	A	11	DG	O4'-C1'-C2'	-5.70	101.34	105.90	2	1
2	B	21	DG	O4'-C1'-C2'	-5.55	101.46	105.90	2	1
1	A	5	DG	C8-N9-C4	-5.52	104.19	106.40	2	2
1	A	8	DT	C4'-C3'-C2'	-5.51	98.14	103.10	2	2
1	A	9	DA	N7-C8-N9	5.49	116.55	113.80	2	2
2	B	15	DA	N7-C8-N9	5.33	116.46	113.80	2	2
1	A	5	DG	C5-N7-C8	-5.31	101.65	104.30	1	2
1	A	6	DT	O4'-C1'-N1	5.26	111.69	108.00	2	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	3	DA	C8-N9-C4	-5.26	103.70	105.80	2	2
1	A	7	DG	O4'-C1'-N9	5.23	111.66	108.00	2	2
2	B	13	DG	C8-N9-C4	-5.15	104.34	106.40	2	2
1	A	9	DA	P-O3'-C3'	5.13	125.85	119.70	1	1
2	B	12	DC	C1'-O4'-C4'	-5.12	104.98	110.10	1	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	225	126	126	2±0
2	B	221	125	125	1±0
All	All	892	502	502	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:9:DA:C2	2:B:15:DA:C2	0.62	2.88	1	2
1:A:1:DG:H1'	1:A:2:DC:O5'	0.45	2.12	1	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

There are no protein molecules in this entry.

6.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	A38	B	17	1,2	15,24,25	0.98±0.02	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	A38	B	17	1,2	19,35,38	1.52±0.01	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A38	B	17	1,2	-	0±0,3,21,22	0±0,3,3,3

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided