



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:07 AM GMT

PDB ID : 2FK0
Title : Crystal Structure of a H5N1 influenza virus hemagglutinin.
Authors : Stevens, J.; Wilson, I.A.
Deposited on : 2006-01-03
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2184 (3.00-2.92)
Clashscore	102246	2552 (3.00-2.92)
Ramachandran outliers	100387	2468 (3.00-2.92)
Sidechain outliers	100360	2471 (3.00-2.92)
RSRZ outliers	91569	2201 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>16%</div> <div>61%</div> <div>29%</div> <div>6%</div> <div>• •</div> </div>
1	C	334	<div> <div>15%</div> <div>60%</div> <div>31%</div> <div>• • •</div> </div>
1	E	334	<div> <div>14%</div> <div>56%</div> <div>34%</div> <div>6%</div> <div>• •</div> </div>
1	G	334	<div> <div>12%</div> <div>58%</div> <div>32%</div> <div>6%</div> <div>• •</div> </div>
1	I	334	<div> <div>16%</div> <div>57%</div> <div>32%</div> <div>7%</div> <div>• •</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	334	
1	M	334	
1	O	334	
1	Q	334	
2	B	181	
2	D	181	
2	F	181	
2	H	181	
2	J	181	
2	L	181	
2	N	181	
2	P	181	
2	R	181	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	C	3	X	-	-	-
3	NAG	I	3	X	-	-	-
3	NAG	K	3	X	-	-	X
3	NAG	M	3	X	-	-	X
3	NAG	O	3	X	-	-	X
4	NAG	A	3	X	-	-	-
4	NAG	E	3	X	-	-	-
4	BMA	E	5	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 36202 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			
1	C	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			
1	E	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			
1	G	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			
1	I	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			
1	K	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			
1	M	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			
1	O	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			
1	Q	322	Total	C	N	O	S	0	0	0
			2553	1613	441	484	15			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ALA	-	CLONING ARTIFACT	GB 50296053
A	8	ASP	-	CLONING ARTIFACT	GB 50296053
A	9	PRO	-	CLONING ARTIFACT	GB 50296053
A	10	GLY	-	CLONING ARTIFACT	GB 50296053
C	7	ALA	-	CLONING ARTIFACT	GB 50296053
C	8	ASP	-	CLONING ARTIFACT	GB 50296053
C	9	PRO	-	CLONING ARTIFACT	GB 50296053
C	10	GLY	-	CLONING ARTIFACT	GB 50296053
E	7	ALA	-	CLONING ARTIFACT	GB 50296053
E	8	ASP	-	CLONING ARTIFACT	GB 50296053
E	9	PRO	-	CLONING ARTIFACT	GB 50296053

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	10	GLY	-	CLONING ARTIFACT	GB 50296053
G	7	ALA	-	CLONING ARTIFACT	GB 50296053
G	8	ASP	-	CLONING ARTIFACT	GB 50296053
G	9	PRO	-	CLONING ARTIFACT	GB 50296053
G	10	GLY	-	CLONING ARTIFACT	GB 50296053
I	7	ALA	-	CLONING ARTIFACT	GB 50296053
I	8	ASP	-	CLONING ARTIFACT	GB 50296053
I	9	PRO	-	CLONING ARTIFACT	GB 50296053
I	10	GLY	-	CLONING ARTIFACT	GB 50296053
K	7	ALA	-	CLONING ARTIFACT	GB 50296053
K	8	ASP	-	CLONING ARTIFACT	GB 50296053
K	9	PRO	-	CLONING ARTIFACT	GB 50296053
K	10	GLY	-	CLONING ARTIFACT	GB 50296053
M	7	ALA	-	CLONING ARTIFACT	GB 50296053
M	8	ASP	-	CLONING ARTIFACT	GB 50296053
M	9	PRO	-	CLONING ARTIFACT	GB 50296053
M	10	GLY	-	CLONING ARTIFACT	GB 50296053
O	7	ALA	-	CLONING ARTIFACT	GB 50296053
O	8	ASP	-	CLONING ARTIFACT	GB 50296053
O	9	PRO	-	CLONING ARTIFACT	GB 50296053
O	10	GLY	-	CLONING ARTIFACT	GB 50296053
Q	7	ALA	-	CLONING ARTIFACT	GB 50296053
Q	8	ASP	-	CLONING ARTIFACT	GB 50296053
Q	9	PRO	-	CLONING ARTIFACT	GB 50296053
Q	10	GLY	-	CLONING ARTIFACT	GB 50296053

- Molecule 2 is a protein called hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			
2	D	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			
2	F	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			
2	H	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			
2	J	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			
2	L	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			
2	N	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			
2	R	175	Total	C	N	O	S	0	0	0
			1416	880	246	282	8			

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	175	SER	-	CLONING ARTIFACT	GB 58618438
B	176	GLY	-	CLONING ARTIFACT	GB 58618438
B	177	ARG	-	CLONING ARTIFACT	GB 58618438
B	178	LEU	-	CLONING ARTIFACT	GB 58618438
B	179	VAL	-	CLONING ARTIFACT	GB 58618438
B	180	PRO	-	CLONING ARTIFACT	GB 58618438
B	181	ARG	-	CLONING ARTIFACT	GB 58618438
D	175	SER	-	CLONING ARTIFACT	GB 58618438
D	176	GLY	-	CLONING ARTIFACT	GB 58618438
D	177	ARG	-	CLONING ARTIFACT	GB 58618438
D	178	LEU	-	CLONING ARTIFACT	GB 58618438
D	179	VAL	-	CLONING ARTIFACT	GB 58618438
D	180	PRO	-	CLONING ARTIFACT	GB 58618438
D	181	ARG	-	CLONING ARTIFACT	GB 58618438
F	175	SER	-	CLONING ARTIFACT	GB 58618438
F	176	GLY	-	CLONING ARTIFACT	GB 58618438
F	177	ARG	-	CLONING ARTIFACT	GB 58618438
F	178	LEU	-	CLONING ARTIFACT	GB 58618438
F	179	VAL	-	CLONING ARTIFACT	GB 58618438
F	180	PRO	-	CLONING ARTIFACT	GB 58618438
F	181	ARG	-	CLONING ARTIFACT	GB 58618438
H	175	SER	-	CLONING ARTIFACT	GB 58618438
H	176	GLY	-	CLONING ARTIFACT	GB 58618438
H	177	ARG	-	CLONING ARTIFACT	GB 58618438
H	178	LEU	-	CLONING ARTIFACT	GB 58618438
H	179	VAL	-	CLONING ARTIFACT	GB 58618438
H	180	PRO	-	CLONING ARTIFACT	GB 58618438
H	181	ARG	-	CLONING ARTIFACT	GB 58618438
J	175	SER	-	CLONING ARTIFACT	GB 58618438
J	176	GLY	-	CLONING ARTIFACT	GB 58618438
J	177	ARG	-	CLONING ARTIFACT	GB 58618438
J	178	LEU	-	CLONING ARTIFACT	GB 58618438
J	179	VAL	-	CLONING ARTIFACT	GB 58618438
J	180	PRO	-	CLONING ARTIFACT	GB 58618438

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	181	ARG	-	CLONING ARTIFACT	GB 58618438
L	175	SER	-	CLONING ARTIFACT	GB 58618438
L	176	GLY	-	CLONING ARTIFACT	GB 58618438
L	177	ARG	-	CLONING ARTIFACT	GB 58618438
L	178	LEU	-	CLONING ARTIFACT	GB 58618438
L	179	VAL	-	CLONING ARTIFACT	GB 58618438
L	180	PRO	-	CLONING ARTIFACT	GB 58618438
L	181	ARG	-	CLONING ARTIFACT	GB 58618438
N	175	SER	-	CLONING ARTIFACT	GB 58618438
N	176	GLY	-	CLONING ARTIFACT	GB 58618438
N	177	ARG	-	CLONING ARTIFACT	GB 58618438
N	178	LEU	-	CLONING ARTIFACT	GB 58618438
N	179	VAL	-	CLONING ARTIFACT	GB 58618438
N	180	PRO	-	CLONING ARTIFACT	GB 58618438
N	181	ARG	-	CLONING ARTIFACT	GB 58618438
P	175	SER	-	CLONING ARTIFACT	GB 58618438
P	176	GLY	-	CLONING ARTIFACT	GB 58618438
P	177	ARG	-	CLONING ARTIFACT	GB 58618438
P	178	LEU	-	CLONING ARTIFACT	GB 58618438
P	179	VAL	-	CLONING ARTIFACT	GB 58618438
P	180	PRO	-	CLONING ARTIFACT	GB 58618438
P	181	ARG	-	CLONING ARTIFACT	GB 58618438
R	175	SER	-	CLONING ARTIFACT	GB 58618438
R	176	GLY	-	CLONING ARTIFACT	GB 58618438
R	177	ARG	-	CLONING ARTIFACT	GB 58618438
R	178	LEU	-	CLONING ARTIFACT	GB 58618438
R	179	VAL	-	CLONING ARTIFACT	GB 58618438
R	180	PRO	-	CLONING ARTIFACT	GB 58618438
R	181	ARG	-	CLONING ARTIFACT	GB 58618438

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	C	2	Total	C	N	O	0	0
			28	16	2	10		
3	C	2	Total	C	N	O	0	0
			28	16	2	10		
3	E	2	Total	C	N	O	0	0
			28	16	2	10		
3	G	2	Total	C	N	O	0	0
			28	16	2	10		


Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	I	2	Total	C	N	O	0	0
			28	16	2	10		
3	I	2	Total	C	N	O	0	0
			28	16	2	10		
3	K	2	Total	C	N	O	0	0
			28	16	2	10		
3	K	2	Total	C	N	O	0	0
			28	16	2	10		
3	M	2	Total	C	N	O	0	0
			28	16	2	10		
3	M	2	Total	C	N	O	0	0
			28	16	2	10		
3	O	2	Total	C	N	O	0	0
			28	16	2	10		
3	O	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 4 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	3	Total	C	N	O	0	0
			39	22	2	15		
4	E	3	Total	C	N	O	0	0
			39	22	2	15		
4	G	3	Total	C	N	O	0	0
			39	22	2	15		







• Molecule 1: hemagglutinin




• Molecule 2: hemagglutinin






• Molecule 2: hemagglutinin

Chain D:




• Molecule 2: hemagglutinin

Chain F:




• Molecule 2: hemagglutinin


Chain H:




• Molecule 2: hemagglutinin




• Molecule 2: hemagglutinin




• Molecule 2: hemagglutinin





• Molecule 2: hemagglutinin





● Molecule 2: hemagglutinin



4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	197.94Å 197.94Å 134.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	171.50 – 2.95 49.49 – 2.95	Depositor EDS
% Data completeness (in resolution range)	98.9 (171.50-2.95) 98.9 (49.49-2.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.268 , 0.319 0.266 , 0.314	Depositor DCC
R_{free} test set	1233 reflections (1.02%)	DCC
Wilson B-factor (Å ²)	69.3	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 98.8	EDS
Estimated twinning fraction	0.012 for -h,-k,l 0.008 for h,-h-k,-l 0.000 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 122499 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	36202	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/2615	0.83	3/3551 (0.1%)
1	C	0.84	3/2615 (0.1%)	0.87	2/3551 (0.1%)
1	E	0.84	1/2615 (0.0%)	0.87	4/3551 (0.1%)
1	G	0.84	3/2615 (0.1%)	0.85	3/3551 (0.1%)
1	I	0.80	2/2615 (0.1%)	0.85	4/3551 (0.1%)
1	K	0.79	1/2615 (0.0%)	0.83	2/3551 (0.1%)
1	M	1.12	21/2615 (0.8%)	0.87	5/3551 (0.1%)
1	O	1.49	22/2615 (0.8%)	0.91	9/3551 (0.3%)
1	Q	0.65	2/2615 (0.1%)	0.70	0/3551
2	B	0.67	0/1443	0.70	3/1939 (0.2%)
2	D	0.68	0/1443	0.71	3/1939 (0.2%)
2	F	0.70	3/1443 (0.2%)	0.71	3/1939 (0.2%)
2	H	0.67	1/1443 (0.1%)	0.70	3/1939 (0.2%)
2	J	0.64	0/1443	0.68	2/1939 (0.1%)
2	L	0.67	1/1443 (0.1%)	0.67	2/1939 (0.1%)
2	N	1.29	7/1443 (0.5%)	0.86	5/1939 (0.3%)
2	P	1.38	13/1443 (0.9%)	0.88	5/1939 (0.3%)
2	R	0.74	5/1443 (0.3%)	0.69	0/1939
All	All	0.91	85/36522 (0.2%)	0.81	58/49410 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	2
1	E	0	2
1	G	0	2
1	I	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	K	0	3
1	M	0	1
1	O	0	3
1	Q	0	1
2	B	0	3
2	D	0	3
2	F	0	3
2	H	0	3
2	J	0	2
2	L	0	3
2	N	0	2
2	P	0	2
3	C	1	0
3	I	1	0
3	K	1	0
3	M	1	0
3	O	1	0
4	A	1	0
4	E	1	0
All	All	7	38

The worst 5 of 85 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	O	125(B)	SER	CB-OG	34.55	1.87	1.42
2	N	38	LYS	CE-NZ	31.98	2.29	1.49
2	P	143	LYS	CE-NZ	27.78	2.18	1.49
1	O	157	LYS	CD-CE	22.51	2.07	1.51
1	O	130	HIS	CE1-NE2	20.34	1.79	1.32

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	N	38	LYS	CD-CE-NZ	-14.03	79.42	111.70
1	M	166	ARG	NE-CZ-NH2	-12.85	113.88	120.30
1	O	125(A)	LYS	CD-CE-NZ	-11.74	84.70	111.70
1	M	154	LEU	CB-CG-CD2	11.21	130.05	111.00
2	P	143	LYS	CD-CE-NZ	-10.31	87.98	111.70

5 of 7 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	3	NAG	C1
3	C	3	NAG	C1
4	E	3	NAG	C1
3	I	3	NAG	C1
3	K	3	NAG	C1

5 of 38 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	248	ASN	Peptide
1	A	276	ASN	Peptide
2	B	60	ASN	Peptide
2	B	61	THR	Peptide
2	B	68	ARG	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2553	0	2496	117	2
1	C	2553	0	2496	118	3
1	E	2553	0	2496	169	0
1	G	2553	0	2496	131	3
1	I	2553	0	2496	140	0
1	K	2553	0	2496	117	2
1	M	2553	0	2496	166	0
1	O	2553	0	2496	191	0
1	Q	2553	0	2498	132	0
2	B	1416	0	1320	47	0
2	D	1416	0	1320	46	0
2	F	1416	0	1320	72	0
2	H	1416	0	1320	60	0
2	J	1416	0	1320	42	0
2	L	1416	0	1320	39	0
2	N	1416	0	1320	75	0
2	P	1416	0	1320	82	0
2	R	1416	0	1320	77	0
3	A	28	0	25	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	56	0	50	1	0
3	E	28	0	25	0	0
3	G	28	0	25	2	0
3	I	56	0	50	1	0
3	K	56	0	50	0	0
3	M	56	0	50	4	0
3	O	56	0	50	2	0
4	A	39	0	34	3	0
4	E	39	0	34	3	0
4	G	39	0	34	0	0
All	All	36202	0	34773	1624	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 1624 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:154:LEU:CD1	1:M:154:LEU:CG	1.79	1.60
1:M:130:HIS:CG	1:M:130:HIS:ND1	1.71	1.54
1:M:130:HIS:NE2	1:M:130:HIS:CE1	1.75	1.53
1:O:125(A):LYS:CE	1:O:125(A):LYS:CD	1.83	1.52
2:P:143:LYS:CE	2:P:143:LYS:CD	1.86	1.51

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:GLU:OE2	1:K:142:GLN:OE1[3_455]	1.69	0.51
1:C:142:GLN:OE1	1:G:78:GLU:OE2[2_555]	1.87	0.33
1:C:79:PHE:CE2	1:G:125(B):SER:C[2_555]	1.94	0.26
1:C:142:GLN:OE1	1:G:149:ARG:NH1[2_555]	2.17	0.03
1:A:125(B):SER:C	1:K:79:PHE:CE2[3_455]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/334 (96%)	289 (90%)	25 (8%)	6 (2%)	10	40
1	C	320/334 (96%)	290 (91%)	26 (8%)	4 (1%)	15	51
1	E	320/334 (96%)	289 (90%)	25 (8%)	6 (2%)	10	40
1	G	320/334 (96%)	285 (89%)	31 (10%)	4 (1%)	15	51
1	I	320/334 (96%)	285 (89%)	29 (9%)	6 (2%)	10	40
1	K	320/334 (96%)	287 (90%)	28 (9%)	5 (2%)	12	45
1	M	320/334 (96%)	255 (80%)	52 (16%)	13 (4%)	3	17
1	O	320/334 (96%)	248 (78%)	57 (18%)	15 (5%)	3	15
1	Q	320/334 (96%)	237 (74%)	63 (20%)	20 (6%)	2	8
2	B	173/181 (96%)	161 (93%)	10 (6%)	2 (1%)	16	53
2	D	173/181 (96%)	164 (95%)	7 (4%)	2 (1%)	16	53
2	F	173/181 (96%)	162 (94%)	9 (5%)	2 (1%)	16	53
2	H	173/181 (96%)	162 (94%)	8 (5%)	3 (2%)	11	43
2	J	173/181 (96%)	163 (94%)	8 (5%)	2 (1%)	16	53
2	L	173/181 (96%)	164 (95%)	7 (4%)	2 (1%)	16	53
2	N	173/181 (96%)	132 (76%)	35 (20%)	6 (4%)	4	22
2	P	173/181 (96%)	126 (73%)	37 (21%)	10 (6%)	2	10
2	R	173/181 (96%)	124 (72%)	29 (17%)	20 (12%)	0	2
All	All	4437/4635 (96%)	3823 (86%)	486 (11%)	128 (3%)	6	27

5 of 128 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	60	ASN
2	B	127	ARG
2	D	60	ASN
2	D	127	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	80	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/300 (96%)	257 (89%)	32 (11%)	8	27
1	C	289/300 (96%)	260 (90%)	29 (10%)	9	32
1	E	289/300 (96%)	257 (89%)	32 (11%)	8	27
1	G	289/300 (96%)	258 (89%)	31 (11%)	8	29
1	I	289/300 (96%)	256 (89%)	33 (11%)	7	26
1	K	289/300 (96%)	255 (88%)	34 (12%)	6	24
1	M	289/300 (96%)	239 (83%)	50 (17%)	2	10
1	O	289/300 (96%)	246 (85%)	43 (15%)	4	15
1	Q	289/300 (96%)	238 (82%)	51 (18%)	2	10
2	B	149/155 (96%)	135 (91%)	14 (9%)	11	36
2	D	149/155 (96%)	135 (91%)	14 (9%)	11	36
2	F	149/155 (96%)	135 (91%)	14 (9%)	11	36
2	H	149/155 (96%)	134 (90%)	15 (10%)	9	32
2	J	149/155 (96%)	135 (91%)	14 (9%)	11	36
2	L	149/155 (96%)	135 (91%)	14 (9%)	11	36
2	N	149/155 (96%)	121 (81%)	28 (19%)	2	8
2	P	149/155 (96%)	121 (81%)	28 (19%)	2	8
2	R	149/155 (96%)	125 (84%)	24 (16%)	3	12
All	All	3942/4095 (96%)	3442 (87%)	500 (13%)	5	21

5 of 500 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	169	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	M	53(A)	LEU
1	Q	238	LYS
1	K	37	THR
1	K	217	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 77 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	172	ASN
2	N	53	ASN
2	R	28	ASN
1	M	191	GLN
1	M	312	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

35 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1	1,3	14,14,15	0.61	0	15,19,21	1.74	4 (26%)
3	NAG	A	2	3	14,14,15	0.93	1 (7%)	15,19,21	0.74	0
4	NAG	A	3	1,4	14,14,15	0.68	0	15,19,21	1.33	1 (6%)
4	NAG	A	4	4	14,14,15	0.74	0	15,19,21	1.76	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BMA	A	5	4	11,11,12	0.69	0	14,15,17	1.51	2 (14%)
3	NAG	C	1	1,3	14,14,15	0.69	0	15,19,21	2.92	5 (33%)
3	NAG	C	2	3	14,14,15	0.60	0	15,19,21	1.51	1 (6%)
3	NAG	C	3	1,3	14,14,15	0.85	1 (7%)	15,19,21	2.38	3 (20%)
3	NAG	C	4	3	14,14,15	0.58	0	15,19,21	1.89	3 (20%)
3	NAG	E	1	1,3	14,14,15	0.58	0	15,19,21	1.88	4 (26%)
3	NAG	E	2	3	14,14,15	0.92	1 (7%)	15,19,21	1.79	2 (13%)
4	NAG	E	3	1,4	14,14,15	0.63	0	15,19,21	1.08	1 (6%)
4	NAG	E	4	4	14,14,15	0.57	0	15,19,21	1.35	3 (20%)
4	BMA	E	5	4	11,11,12	0.85	0	14,15,17	2.71	4 (28%)
3	NAG	G	1	1,3	14,14,15	0.66	1 (7%)	15,19,21	1.22	1 (6%)
3	NAG	G	2	3	14,14,15	1.01	1 (7%)	15,19,21	1.46	2 (13%)
4	NAG	G	3	1,4	14,14,15	0.77	0	15,19,21	1.92	3 (20%)
4	NAG	G	4	4	14,14,15	0.73	0	15,19,21	1.46	2 (13%)
4	BMA	G	5	4	11,11,12	0.54	0	14,15,17	2.60	1 (7%)
3	NAG	I	1	1,3	14,14,15	0.80	1 (7%)	15,19,21	2.84	5 (33%)
3	NAG	I	2	3	14,14,15	0.81	1 (7%)	15,19,21	1.53	3 (20%)
3	NAG	I	3	1,3	14,14,15	0.82	1 (7%)	15,19,21	1.55	4 (26%)
3	NAG	I	4	3	14,14,15	0.63	0	15,19,21	1.23	1 (6%)
3	NAG	K	1	1,3	14,14,15	0.75	0	15,19,21	2.21	5 (33%)
3	NAG	K	2	3	14,14,15	1.03	1 (7%)	15,19,21	2.69	5 (33%)
3	NAG	K	3	1,3	14,14,15	0.70	0	15,19,21	1.86	2 (13%)
3	NAG	K	4	3	14,14,15	0.52	0	15,19,21	1.32	2 (13%)
3	NAG	M	1	1,3	14,14,15	0.79	0	15,19,21	1.86	4 (26%)
3	NAG	M	2	3	14,14,15	0.94	1 (7%)	15,19,21	1.14	0
3	NAG	M	3	1,3	14,14,15	0.91	1 (7%)	15,19,21	1.30	1 (6%)
3	NAG	M	4	3	14,14,15	0.86	0	15,19,21	2.20	4 (26%)
3	NAG	O	1	1,3	14,14,15	0.91	1 (7%)	15,19,21	1.94	4 (26%)
3	NAG	O	2	3	14,14,15	1.24	1 (7%)	15,19,21	1.84	2 (13%)
3	NAG	O	3	1,3	14,14,15	0.71	0	15,19,21	2.48	4 (26%)
3	NAG	O	4	3	14,14,15	0.88	1 (7%)	15,19,21	1.21	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	2	3	-	0/6/23/26	0/1/1/1
4	NAG	A	3	1,4	1/1/5/7	0/6/23/26	0/1/1/1
4	NAG	A	4	4	-	0/6/23/26	0/1/1/1
4	BMA	A	5	4	-	0/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	NAG	C	3	1,3	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	C	4	3	-	0/6/23/26	0/1/1/1
3	NAG	E	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
4	NAG	E	3	1,4	1/1/5/7	0/6/23/26	0/1/1/1
4	NAG	E	4	4	-	0/6/23/26	0/1/1/1
4	BMA	E	5	4	-	0/2/19/22	0/1/1/1
3	NAG	G	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	1/6/23/26	0/1/1/1
4	NAG	G	3	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	4	4	-	0/6/23/26	0/1/1/1
4	BMA	G	5	4	-	0/2/19/22	0/1/1/1
3	NAG	I	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	0/6/23/26	0/1/1/1
3	NAG	I	3	1,3	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	I	4	3	-	0/6/23/26	0/1/1/1
3	NAG	K	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	K	2	3	-	0/6/23/26	0/1/1/1
3	NAG	K	3	1,3	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	K	4	3	-	0/6/23/26	0/1/1/1
3	NAG	M	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	M	2	3	-	1/6/23/26	0/1/1/1
3	NAG	M	3	1,3	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	M	4	3	-	2/6/23/26	0/1/1/1
3	NAG	O	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	O	2	3	-	0/6/23/26	0/1/1/1
3	NAG	O	3	1,3	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	O	4	3	-	1/6/23/26	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	3	NAG	O5-C1	-2.11	1.40	1.43
3	G	1	NAG	C1-C2	2.01	1.55	1.52
3	I	2	NAG	C1-C2	2.16	1.55	1.52
3	I	3	NAG	C1-C2	2.23	1.55	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	1	NAG	C1-C2	2.34	1.55	1.52

The worst 5 of 92 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3	NAG	C1-O5-C5	-6.18	104.40	112.25
3	O	3	NAG	C4-C3-C2	-5.58	102.56	111.23
3	C	4	NAG	C2-N2-C7	-5.36	116.16	123.04
3	C	3	NAG	C2-N2-C7	-4.91	116.73	123.04
4	A	4	NAG	C1-O5-C5	-4.82	106.13	112.25

5 of 7 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	E	3	NAG	C1
3	I	3	NAG	C1
3	C	3	NAG	C1
4	A	3	NAG	C1
3	O	3	NAG	C1

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	2	NAG	O7-C7-N2-C2
3	M	2	NAG	O7-C7-N2-C2
3	O	4	NAG	O7-C7-N2-C2
3	M	4	NAG	C8-C7-N2-C2
3	C	2	NAG	C8-C7-N2-C2

There are no ring outliers.

15 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1	NAG	1	0
3	A	2	NAG	1	0
4	A	3	NAG	1	0
4	A	4	NAG	3	0
3	C	1	NAG	1	0
4	E	5	BMA	3	0
3	G	1	NAG	2	0
3	G	2	NAG	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	1	NAG	1	0
3	M	1	NAG	1	0
3	M	2	NAG	1	0
3	M	3	NAG	2	0
3	M	4	NAG	3	0
3	O	1	NAG	2	0
3	O	2	NAG	1	0

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/334 (96%)	1.31	55 (17%) 2 1	70, 71, 72, 73	0
1	C	322/334 (96%)	1.21	49 (15%) 3 1	70, 71, 72, 73	0
1	E	322/334 (96%)	1.38	48 (14%) 3 1	70, 71, 72, 73	0
1	G	322/334 (96%)	1.34	41 (12%) 5 2	70, 71, 72, 73	0
1	I	322/334 (96%)	1.30	53 (16%) 2 1	70, 71, 72, 73	0
1	K	322/334 (96%)	1.40	61 (18%) 2 1	70, 71, 72, 74	0
1	M	322/334 (96%)	2.24	154 (47%) 0 0	54, 67, 75, 84	0
1	O	322/334 (96%)	2.10	140 (43%) 0 0	57, 68, 76, 83	0
1	Q	322/334 (96%)	3.62	231 (71%) 0 0	63, 72, 81, 86	0
2	B	175/181 (96%)	2.32	73 (41%) 0 0	70, 71, 72, 73	0
2	D	175/181 (96%)	2.67	90 (51%) 0 0	70, 71, 72, 73	0
2	F	175/181 (96%)	2.59	77 (44%) 0 0	70, 71, 72, 73	0
2	H	175/181 (96%)	2.83	90 (51%) 0 0	70, 71, 72, 73	0
2	J	175/181 (96%)	2.57	86 (49%) 0 0	70, 71, 72, 73	0
2	L	175/181 (96%)	2.83	89 (50%) 0 0	70, 71, 72, 73	0
2	N	175/181 (96%)	2.06	68 (38%) 0 0	55, 71, 82, 87	0
2	P	175/181 (96%)	2.16	65 (37%) 0 0	58, 68, 80, 82	0
2	R	175/181 (96%)	3.99	123 (70%) 0 0	60, 70, 85, 87	0
All	All	4473/4635 (96%)	2.08	1593 (35%) 0 0	54, 71, 77, 87	0

The worst 5 of 1593 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	324	PRO	21.9
2	R	134	GLY	20.1
1	I	16	GLY	18.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	Q	324	PRO	17.9
1	Q	323	SER	16.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	O	3	14/15	0.67	0.48	4.93	87,91,94,99	0
3	NAG	K	3	14/15	0.84	0.29	2.38	61,71,74,80	0
3	NAG	C	3	14/15	0.73	0.33	1.64	70,73,77,78	0
4	NAG	E	3	14/15	0.81	0.33	1.06	79,81,84,89	0
3	NAG	I	3	14/15	0.79	0.30	-0.27	64,72,76,79	0
4	BMA	E	5	11/12	0.46	0.47	-0.32	105,106,107,107	0
3	NAG	M	3	14/15	0.78	0.41	-0.52	82,85,87,89	0
4	NAG	G	3	14/15	0.85	0.22	-1.04	67,69,73,78	0
4	NAG	A	3	14/15	0.88	0.20	-2.12	57,64,69,75	0
4	NAG	G	4	14/15	0.64	0.37	-	82,85,88,91	0
3	NAG	I	1	14/15	0.64	0.42	-	90,97,100,103	0
3	NAG	K	1	14/15	0.62	0.44	-	90,97,100,106	0
4	BMA	G	5	11/12	0.53	0.59	-	94,96,97,97	0
4	BMA	A	5	11/12	0.31	0.56	-	96,100,101,102	0
3	NAG	I	4	14/15	0.70	0.44	-	82,85,88,88	0
3	NAG	G	2	14/15	0.62	0.46	-	103,106,107,108	0
3	NAG	C	4	14/15	0.75	0.54	-	80,82,86,86	0
3	NAG	K	4	14/15	0.74	0.49	-	82,84,85,87	0
3	NAG	O	4	14/15	0.46	0.52	-	104,106,111,112	0
3	NAG	A	2	14/15	0.63	0.56	-	97,100,103,104	0
3	NAG	C	1	14/15	0.77	0.43	-	85,91,92,94	0
3	NAG	K	2	14/15	0.46	0.44	-	109,112,114,115	0
3	NAG	O	1	14/15	0.60	0.40	-	99,104,107,113	0
3	NAG	I	2	14/15	0.62	0.69	-	107,108,109,110	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	M	4	14/15	0.51	0.33	-	92,93,94,94	0
3	NAG	C	2	14/15	0.70	0.51	-	94,95,97,97	0
3	NAG	G	1	14/15	0.80	0.35	-	87,91,94,100	0
3	NAG	M	1	14/15	0.73	0.50	-	92,98,102,105	0
4	NAG	A	4	14/15	0.69	0.41	-	80,84,88,92	0
3	NAG	M	2	14/15	0.58	0.63	-	109,110,110,110	0
4	NAG	E	4	14/15	0.61	0.40	-	94,97,100,103	0
3	NAG	O	2	14/15	0.07	0.68	-	115,117,118,118	0
3	NAG	A	1	14/15	0.70	0.48	-	87,93,97,98	0
3	NAG	E	1	14/15	0.71	0.36	-	87,91,96,102	0
3	NAG	E	2	14/15	0.28	0.49	-	108,111,113,113	0

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.