



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:06 AM GMT

PDB ID : 2FN3
Title : High resolution structure of s26a mutant of benzoylformate decarboxylase from pseudomonas putida complexed with thiamine thiazolone diphosphate
Authors : Bera, A.K.; Hasson, M.S.
Deposited on : 2006-01-10
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1235 (1.10-0.90)
Clashscore	102246	1333 (1.10-0.90)
Ramachandran outliers	100387	1247 (1.10-0.90)
Sidechain outliers	100360	1246 (1.10-0.90)
RSRZ outliers	91569	1239 (1.10-0.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	528	<div> <div>2%</div> <div>90%</div> <div>8% ..</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Benzoylformate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	524	Total	C	N	O	S	0	10	0
			3946	2490	682	752	22			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	ALA	SER	ENGINEERED	UNP P20906


- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ca	0	0
			3	3		

- Molecule 4 is 2-{3-[(4-AMINO-2-METHYLPYRIMIDIN-5-YL)METHYL]-4-METHYL-2-OXO-2,3-DIHYDRO-1,3-THIAZOL-5-YL}ETHYL TRIHYDROGEN DIPHOSPHATE (three-letter code: TZD) (formula: C₁₂H₁₈N₄O₈P₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	A	1	Total	C	N	O	P	S	0	0
			27	12	4	8	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	550	Total	O	0	0
			550	550		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Benzoylformate decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	80.93 Å 95.73 Å 136.86 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.00 27.84 – 1.00	Depositor EDS
% Data completeness (in resolution range)	83.1 (20.00-1.00) 83.4 (27.84-1.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.67 (at 1.00 Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.128 , (Not available) 0.158 , 0.166	Depositor DCC
R_{free} test set	12422 reflections (5.51%)	DCC
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 62.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 248207 reflections	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4527	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MG, TZD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	3/4094 (0.1%)	1.20	37/5593 (0.7%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	126[A]	SER	CB-OG	-6.79	1.33	1.42
1	A	126[B]	SER	CB-OG	-6.79	1.33	1.42
1	A	420	ARG	CZ-NH2	-6.01	1.25	1.33

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	120	ARG	NE-CZ-NH1	-17.08	111.76	120.30
1	A	226	ARG	NE-CZ-NH1	13.49	127.04	120.30
1	A	245	ARG	NE-CZ-NH2	-11.30	114.65	120.30
1	A	226	ARG	NE-CZ-NH2	-11.10	114.75	120.30
1	A	120	ARG	NE-CZ-NH2	10.49	125.55	120.30
1	A	462	ARG	NE-CZ-NH2	-10.26	115.17	120.30
1	A	469	GLU	OE1-CD-OE2	-8.95	112.56	123.30
1	A	13	ARG	NE-CZ-NH2	-8.75	115.92	120.30
1	A	352	ARG	CD-NE-CZ	7.47	134.06	123.60
1	A	281	HIS	CA-CB-CG	7.37	126.12	113.60
1	A	250	ARG	NE-CZ-NH2	-7.31	116.64	120.30
1	A	127	TYR	CB-CG-CD2	7.23	125.34	121.00
1	A	458	TYR	CB-CG-CD1	7.12	125.27	121.00
1	A	120	ARG	CD-NE-CZ	7.00	133.40	123.60
1	A	294	ARG	CD-NE-CZ	6.91	133.27	123.60
1	A	141	ARG	NE-CZ-NH2	-6.84	116.88	120.30
1	A	283	TYR	CB-CG-CD1	-6.84	116.90	121.00
1	A	245	ARG	NE-CZ-NH1	6.74	123.67	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	40	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	A	40	ARG	NE-CZ-NH1	-6.10	117.25	120.30
1	A	40	ARG	NH1-CZ-NH2	5.80	125.79	119.40
1	A	103[A]	MET	CG-SD-CE	-5.71	91.07	100.20
1	A	103[B]	MET	CG-SD-CE	-5.71	91.07	100.20
1	A	166	ASP	CB-CG-OD1	5.69	123.42	118.30
1	A	294	ARG	NE-CZ-NH1	-5.65	117.47	120.30
1	A	191	ASP	CB-CG-OD1	5.63	123.37	118.30
1	A	433	TYR	CB-CG-CD2	5.54	124.32	121.00
1	A	458	TYR	CB-CG-CD2	-5.54	117.68	121.00
1	A	420	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	484	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	A	249	PHE	CB-CG-CD1	-5.20	117.16	120.80
1	A	498	ASP	CB-CG-OD2	5.17	122.95	118.30
1	A	17	ILE	CB-CG1-CD1	5.14	128.28	113.90
1	A	232	TRP	CH2-CZ2-CE2	-5.08	112.32	117.40
1	A	189	ASP	CB-CG-OD1	5.07	122.86	118.30
1	A	62	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	A	397	PHE	CD1-CE1-CZ	-5.01	114.08	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3946	0	3873	16	0
2	A	1	0	0	0	0
3	A	3	0	0	0	0
4	A	27	0	15	0	0
5	A	550	0	0	7	0
All	All	4527	0	3888	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:GLU:HG2	5:A:957:HOH:O	1.90	0.70
1:A:496:LYS:HG3	5:A:826:HOH:O	2.01	0.61
1:A:232:TRP:CH2	1:A:263:LEU:HD11	2.36	0.60
1:A:232:TRP:CZ2	1:A:263:LEU:HD11	2.43	0.54
1:A:439:TRP:CZ2	1:A:443:GLN:HG3	2.44	0.53
1:A:40:ARG:HG3	5:A:917:HOH:O	2.09	0.52
1:A:439:TRP:CE2	1:A:443:GLN:HG3	2.45	0.51
1:A:61[B]:SER:OG	1:A:63:LYS:HG2	2.11	0.51
1:A:46:GLN:NE2	1:A:48:ALA:H	2.08	0.50
1:A:496:LYS:HE3	5:A:826:HOH:O	2.15	0.47
1:A:345:LYS:HE3	5:A:846:HOH:O	2.14	0.47
1:A:232:TRP:HZ3	5:A:965:HOH:O	1.99	0.45
1:A:352:ARG:HG2	1:A:522[B]:SER:OG	2.19	0.42
1:A:222[B]:MET:HE2	5:A:963:HOH:O	2.18	0.42
1:A:79[B]:MET:SD	1:A:119:PRO:HA	2.60	0.41
1:A:100:THR:CG2	1:A:103[B]:MET:HG3	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	532/528 (101%)	524 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	419/413 (102%)	414 (99%)	5 (1%)	78	43

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	GLN
1	A	217	ASN
1	A	281	HIS
1	A	310	MET
1	A	388	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	GLN
1	A	68	ASN
1	A	281	HIS
1	A	335	GLN
1	A	388	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TZD	A	530	3	22,28,28	1.16	1 (4%)	31,42,42	1.42	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TZD	A	530	3	-	0/16/17/17	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	530	TZD	OC2-C2	3.74	1.27	1.21

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	530	TZD	C4A-C4-N3	-3.02	118.83	122.82
4	A	530	TZD	N1'-C2'-N3'	-2.94	120.17	125.60
4	A	530	TZD	C5'-C6'-N1'	-2.10	120.22	123.86
4	A	530	TZD	C5-C4-N3	2.05	112.59	107.83
4	A	530	TZD	C2A-C2'-N1'	2.56	120.10	117.03
4	A	530	TZD	C6'-N1'-C2'	3.69	122.22	115.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	524/528 (99%)	0.01	13 (2%) 61 50	12, 15, 25, 42	32 (6%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	463	TRP	4.3
1	A	466	GLY	3.4
1	A	469	GLU	3.3
1	A	464	PHE	3.1
1	A	468	LEU	2.8
1	A	467	VAL	2.8
1	A	514	LYS	2.8
1	A	525	SER	2.7
1	A	2	ALA	2.3
1	A	283	TYR	2.2
1	A	350	ALA	2.1
1	A	471	GLU	2.1
1	A	207	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CA	A	533	1/1	0.95	0.08	0.64	18,18,18,18	1
3	CA	A	534	1/1	0.99	0.04	-1.09	15,15,15,15	1
4	TZD	A	530	27/27	0.99	0.04	-1.28	12,13,15,17	1
2	MG	A	531	1/1	0.99	0.04	-2.17	13,13,13,13	1
3	CA	A	532	1/1	1.00	0.05	-2.45	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.