



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:42 AM GMT

PDB ID : 3FOK  
Title : Crystal Structure of Cgl0159 From *Corynebacterium glutamicum* (*Brevibacterium flavum*). Northeast Structural Genomics Target CgR115  
Authors : Seetharaman, J.; Neely, H.; Wang, H.; Janjua, H.; Foote, E.L.; Xiao, R.; Everett, J.K.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2008-12-30  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	<div> <div>5%</div> <div>63% 28% 6% .</div> </div>
1	B	307	<div> <div>5%</div> <div>65% 25% 6% .</div> </div>
1	C	307	<div> <div>4%</div> <div>61% 29% 6% .</div> </div>
1	D	307	<div> <div>6%</div> <div>64% 26% 6% .</div> </div>
1	E	307	<div> <div>6%</div> <div>65% 26% 6% .</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	307	<div><div></div><div>4%</div><div>64%</div><div>27%</div><div>6%</div><div></div></div>
1	G	307	<div><div></div><div>5%</div><div>64%</div><div>26%</div><div>6%</div><div></div></div>
1	H	307	<div><div></div><div>8%</div><div>63%</div><div>28%</div><div>6%</div><div></div></div>
1	I	307	<div><div></div><div>8%</div><div>62%</div><div>28%</div><div>6%</div><div></div></div>
1	J	307	<div><div></div><div>7%</div><div>62%</div><div>28%</div><div>7%</div><div></div></div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 22795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein Cgl0159.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	B	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	C	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	D	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	E	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	F	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	G	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	H	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	I	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			
1	J	295	Total	C	N	O	Se	0	0	0
			2208	1379	387	429	13			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	72	Total	O	0	0
			72	72		
2	B	90	Total	O	0	0
			90	90		
2	C	77	Total	O	0	0
			77	77		
2	D	95	Total	O	0	0
			95	95		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	79	Total 79	O 79	0	0
2	F	62	Total 62	O 62	0	0
2	G	66	Total 66	O 66	0	0
2	H	74	Total 74	O 74	0	0
2	I	47	Total 47	O 47	0	0
2	J	53	Total 53	O 53	0	0







- Molecule 1: uncharacterized protein Cgl0159




- Molecule 1: uncharacterized protein Cgl0159




- Molecule 1: uncharacterized protein Cgl0159




- Molecule 1: uncharacterized protein Cgl0159



• Molecule 1: uncharacterized protein Cgl0159




• Molecule 1: uncharacterized protein Cgl0159



• Molecule 1: uncharacterized protein Cgl0159







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.00Å 176.21Å 104.24Å 90.00° 101.94° 90.00°	Depositor
Resolution (Å)	39.00 – 2.50 49.50 – 2.39	Depositor EDS
% Data completeness (in resolution range)	94.9 (39.00-2.50) 97.4 (49.50-2.39)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.13 (at 2.39Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.225 , 0.254 0.228 , 0.256	Depositor DCC
$R_{free}$ test set	5276 reflections (4.91%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.7	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 255933 reflections (0.000%)	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	22795	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.30	0/2230	0.68	1/3005 (0.0%)
1	B	0.30	0/2230	0.68	1/3005 (0.0%)
1	C	0.30	0/2230	0.68	1/3005 (0.0%)
1	D	0.31	0/2230	0.67	2/3005 (0.1%)
1	E	0.30	0/2230	0.67	1/3005 (0.0%)
1	F	0.30	0/2230	0.67	1/3005 (0.0%)
1	G	0.69	11/2230 (0.5%)	0.71	2/3005 (0.1%)
1	H	0.30	0/2230	0.67	1/3005 (0.0%)
1	I	0.57	7/2230 (0.3%)	0.70	3/3005 (0.1%)
1	J	0.76	10/2230 (0.4%)	0.69	2/3005 (0.1%)
All	All	0.45	28/22300 (0.1%)	0.68	15/30050 (0.0%)

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	J	278	TYR	CD1-CE1	-17.43	1.13	1.39
1	J	278	TYR	CD2-CE2	-16.39	1.14	1.39
1	G	278	TYR	CD1-CE1	-14.28	1.18	1.39
1	G	278	TYR	CD2-CE2	-12.96	1.20	1.39
1	J	278	TYR	CG-CD1	-11.49	1.24	1.39

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	277	LEU	CA-CB-CG	7.91	133.49	115.30
1	I	277	LEU	CA-CB-CG	7.56	132.69	115.30
1	G	277	LEU	CB-CG-CD1	-6.85	99.36	111.00
1	I	277	LEU	N-CA-C	-5.93	94.99	111.00
1	D	277	LEU	N-CA-C	-5.57	95.96	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2208	0	2202	109	0
1	B	2208	0	2202	105	0
1	C	2208	0	2202	118	0
1	D	2208	0	2202	112	0
1	E	2208	0	2202	103	0
1	F	2208	0	2202	108	0
1	G	2208	0	2202	114	0
1	H	2208	0	2202	107	0
1	I	2208	0	2202	109	0
1	J	2208	0	2202	115	0
2	A	72	0	0	4	0
2	B	90	0	0	1	0
2	C	77	0	0	4	0
2	D	95	0	0	3	0
2	E	79	0	0	2	0
2	F	62	0	0	4	0
2	G	66	0	0	6	0
2	H	74	0	0	2	0
2	I	47	0	0	2	0
2	J	53	0	0	1	0
All	All	22795	0	22020	1090	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 1090 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:76:MSE:CE	1:J:277:LEU:HD21	1.71	1.20
1:C:137:MSE:HE2	1:C:142:VAL:HG21	1.20	1.16
1:F:137:MSE:HE2	1:F:142:VAL:HG21	1.22	1.15
1:A:137:MSE:HE2	1:A:142:VAL:HG21	1.22	1.14
1:J:137:MSE:HE2	1:J:142:VAL:HG21	1.23	1.12

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/307 (95%)	266 (91%)	19 (6%)	6 (2%)	9	14
1	B	291/307 (95%)	269 (92%)	17 (6%)	5 (2%)	11	19
1	C	291/307 (95%)	269 (92%)	17 (6%)	5 (2%)	11	19
1	D	291/307 (95%)	266 (91%)	18 (6%)	7 (2%)	7	11
1	E	291/307 (95%)	267 (92%)	19 (6%)	5 (2%)	11	19
1	F	291/307 (95%)	266 (91%)	19 (6%)	6 (2%)	9	14
1	G	291/307 (95%)	268 (92%)	18 (6%)	5 (2%)	11	19
1	H	291/307 (95%)	266 (91%)	19 (6%)	6 (2%)	9	14
1	I	291/307 (95%)	268 (92%)	16 (6%)	7 (2%)	7	11
1	J	291/307 (95%)	265 (91%)	19 (6%)	7 (2%)	7	11
All	All	2910/3070 (95%)	2670 (92%)	181 (6%)	59 (2%)	9	15

5 of 59 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	THR
1	B	64	THR
1	C	64	THR
1	D	64	THR
1	D	298	ASP

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/229 (100%)	207 (90%)	23 (10%)	9	18
1	B	230/229 (100%)	206 (90%)	24 (10%)	9	16
1	C	230/229 (100%)	205 (89%)	25 (11%)	8	15
1	D	230/229 (100%)	206 (90%)	24 (10%)	9	16
1	E	230/229 (100%)	206 (90%)	24 (10%)	9	16
1	F	230/229 (100%)	206 (90%)	24 (10%)	9	16
1	G	230/229 (100%)	207 (90%)	23 (10%)	9	18
1	H	230/229 (100%)	206 (90%)	24 (10%)	9	16
1	I	230/229 (100%)	208 (90%)	22 (10%)	10	19
1	J	230/229 (100%)	208 (90%)	22 (10%)	10	19
All	All	2300/2290 (100%)	2065 (90%)	235 (10%)	9	17

5 of 235 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	114	ASN
1	F	115	ARG
1	J	57	LEU
1	E	144	PHE
1	E	299	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 109 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	204	GLN
1	F	190	ASN
1	J	68	ASN
1	E	214	ASN
1	F	68	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	282/307 (91%)	0.41	16 (5%) 27 31	10, 23, 47, 62	0
1	B	282/307 (91%)	0.28	15 (5%) 30 34	12, 22, 43, 65	0
1	C	282/307 (91%)	0.24	12 (4%) 39 44	10, 23, 43, 61	0
1	D	282/307 (91%)	0.36	18 (6%) 23 25	12, 23, 45, 67	0
1	E	282/307 (91%)	0.30	17 (6%) 25 28	11, 24, 48, 65	0
1	F	282/307 (91%)	0.27	12 (4%) 39 44	12, 25, 48, 66	0
1	G	282/307 (91%)	0.31	14 (4%) 32 37	13, 26, 45, 67	0
1	H	282/307 (91%)	0.50	24 (8%) 13 14	11, 24, 45, 70	0
1	I	282/307 (91%)	0.53	26 (9%) 11 12	13, 27, 51, 73	0
1	J	282/307 (91%)	0.40	23 (8%) 14 15	12, 29, 51, 68	0
All	All	2820/3070 (91%)	0.36	177 (6%) 23 26	10, 24, 49, 73	0

The worst 5 of 177 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	299	ILE	16.7
1	E	299	ILE	12.5
1	I	299	ILE	11.1
1	H	299	ILE	10.9
1	C	299	ILE	9.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.