



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:38 PM GMT

PDB ID : 4FOR  
Title : Crystal Structure of C-lobe of Bovine lactoferrin Complexed with Flurbiprofen at 1.58 Å Resolution  
Authors : Shukla, P.K.; Gautam, L.; Sinha, M.; Kaur, P.; Sharma, S.; Singh, T.P.  
Deposited on : 2012-06-21  
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3815 (1.60-1.56)
Clashscore	102246	4131 (1.60-1.56)
Ramachandran outliers	100387	4021 (1.60-1.56)
Sidechain outliers	100360	4018 (1.60-1.56)
RSRZ outliers	91569	3823 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	
2	B	6	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	GOL	A	712	-	-	-	X
9	FLP	A	711	-	-	-	X

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 3098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	0	0
			2560	1593	448	499	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	565	LYS	ASN	SEE REMARK 999	UNP P24627
A	608	GLU	LYS	SEE REMARK 999	UNP P24627

- Molecule 2 is a protein called C-terminal peptide from Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	6	Total	C	N	O	S	0	0	0
			45	29	6	9	1			

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is CARBONATE ION (three-letter code: CO3) (formula: CO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 5 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Zn	0	0
			2	2		

- Molecule 7 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	2	Total	C	N	O	0	0
			28	16	2	10		
8	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 9 is FLURBIPROFEN (three-letter code: FLP) (formula:  $C_{15}H_{13}FO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	F	O	0	0
			18	15	1	2		

- Molecule 10 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 11 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	385	Total 385	O 385	0	0
11	B	2	Total 2	O 2	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Lactotransferrin



#### • Molecule 2: C-terminal peptide from Lactotransferrin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.45Å 50.01Å 65.47Å 90.00° 107.14° 90.00°	Depositor
Resolution (Å)	62.56 – 1.58 31.28 – 1.58	Depositor EDS
% Data completeness (in resolution range)	99.8 (62.56-1.58) 99.6 (31.28-1.58)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.22 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.165 , 0.205 0.167 , 0.205	Depositor DCC
$R_{free}$ test set	2691 reflections (5.40%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.2	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.7	EDS
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 52547 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3098	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CO3, NAG, FLP, ZN, FE, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.40	9/2608 (0.3%)	1.35	16/3533 (0.5%)
2	B	0.93	0/45	0.91	0/58
All	All	1.39	9/2653 (0.3%)	1.35	16/3591 (0.4%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	524	TYR	CE1-CZ	-9.21	1.26	1.38
1	A	569	PHE	CG-CD1	6.29	1.48	1.38
1	A	601	SER	CB-OG	-5.85	1.34	1.42
1	A	344	ARG	C-O	5.80	1.34	1.23
1	A	444	GLU	CD-OE1	-5.66	1.19	1.25
1	A	433	TYR	CE1-CZ	5.48	1.45	1.38
1	A	400	TYR	CG-CD2	5.31	1.46	1.39
1	A	444	GLU	CD-OE2	-5.24	1.19	1.25
1	A	603	ARG	CZ-NH1	5.21	1.39	1.33

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	575	ASP	CB-CG-OD1	11.58	128.73	118.30
1	A	600	ARG	NE-CZ-NH2	-8.20	116.20	120.30
1	A	570	ARG	NE-CZ-NH1	7.91	124.26	120.30
1	A	462	ASP	CB-CG-OD2	-7.05	111.95	118.30
1	A	660	TYR	CG-CD1-CE1	6.35	126.38	121.30
1	A	641	PHE	CB-CG-CD1	-6.32	116.38	120.80
1	A	379	ASP	CB-CG-OD2	-6.24	112.69	118.30
1	A	660	TYR	CD1-CE1-CZ	-5.99	114.41	119.80
1	A	344	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	660	TYR	CZ-CE2-CD2	5.57	124.81	119.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	415	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	A	524	TYR	CG-CD1-CE1	-5.48	116.92	121.30
1	A	436	VAL	CA-CB-CG2	5.43	119.04	110.90
1	A	532	CYS	CA-CB-SG	5.38	123.69	114.00
1	A	570	ARG	NE-CZ-NH2	-5.17	117.72	120.30
1	A	524	TYR	CZ-CE2-CD2	-5.02	115.28	119.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2560	0	2480	26	0
2	B	45	0	39	1	0
3	A	1	0	0	0	0
4	A	4	0	0	0	0
5	A	5	0	0	0	0
6	A	2	0	0	0	0
7	A	14	0	13	0	0
8	A	56	0	50	1	0
9	A	18	0	12	2	0
10	A	6	0	8	0	0
11	A	385	0	0	1	0
11	B	2	0	0	0	0
All	All	3098	0	2602	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:565:LYS:HE3	1:A:567:GLU:HB2	1.69	0.74
1:A:565:LYS:HE3	1:A:567:GLU:H	1.55	0.72

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:565:LYS:CE	1:A:567:GLU:HB2	2.21	0.71
9:A:711:FLP:H8	9:A:711:FLP:O1	1.94	0.67
1:A:671:ASN:HA	1:A:674:LYS:HD3	1.78	0.65
1:A:565:LYS:HE3	1:A:567:GLU:N	2.12	0.64
1:A:395:ASP:HA	1:A:595:HIS:CD2	2.38	0.57
1:A:660:TYR:O	9:A:711:FLP:H132	2.06	0.56
1:A:342:TYR:HB2	1:A:606:HIS:HE1	1.71	0.54
1:A:669:ILE:CG2	1:A:673:LYS:HE3	2.40	0.51
1:A:565:LYS:HE3	1:A:567:GLU:CB	2.39	0.50
1:A:364:GLN:HG3	1:A:629:PHE:HB2	1.94	0.48
1:A:352:PRO:HG2	1:A:520:LYS:HD3	1.95	0.48
1:A:364:GLN:OE1	1:A:628:LYS:HE2	2.15	0.47
1:A:633:LYS:HA	1:A:633:LYS:HD3	1.83	0.44
1:A:626:PRO:HA	1:A:630:CYS:SG	2.59	0.43
1:A:668:ALA:HB1	8:A:707:NAG:H83	2.01	0.43
1:A:527:THR:CG2	1:A:560:TRP:HZ2	2.31	0.43
1:A:565:LYS:HD2	1:A:566:ARG:N	2.33	0.42
1:A:509:ASP:HA	11:A:1161:HOH:O	2.20	0.41
1:A:566:ARG:HG2	1:A:581:VAL:HG21	2.01	0.41
1:A:455:LYS:HB3	1:A:504:LEU:HD11	2.02	0.41
1:A:520:LYS:HD2	1:A:520:LYS:HA	1.71	0.41
1:A:565:LYS:CE	1:A:567:GLU:H	2.29	0.41
1:A:522:LYS:HD3	1:A:523:TYR:CE1	2.56	0.41
1:A:404:LYS:NZ	2:B:681:LEU:O	2.54	0.40
1:A:573:CYS:SG	1:A:579:LYS:HG3	2.61	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	333/335 (99%)	324 (97%)	8 (2%)	1 (0%)	46 22

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
All	All	337/341 (99%)	327 (97%)	9 (3%)	1 (0%)	46	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	432	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/278 (100%)	275 (99%)	3 (1%)	80	61
2	B	4/4 (100%)	4 (100%)	0	100	100
All	All	282/282 (100%)	279 (99%)	3 (1%)	80	61

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	419	LYS
1	A	515	CYS
1	A	621	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	355	GLN
1	A	613	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
8	NAG	A	707	1,8	14,14,15	1.10	1 (7%)	15,19,21	1.51	2 (13%)
8	NAG	A	708	8	14,14,15	0.63	0	15,19,21	2.06	5 (33%)
8	NAG	A	709	1,8	14,14,15	0.83	0	15,19,21	2.10	6 (40%)
8	NAG	A	710	8	14,14,15	0.61	0	15,19,21	2.61	9 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	A	707	1,8	-	0/6/23/26	0/1/1/1
8	NAG	A	708	8	-	0/6/23/26	0/1/1/1
8	NAG	A	709	1,8	-	0/6/23/26	0/1/1/1
8	NAG	A	710	8	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	707	NAG	C4-C3	2.09	1.57	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	709	NAG	C2-N2-C7	-3.99	117.92	123.04

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	708	NAG	O3-C3-C2	-3.85	101.49	109.11
8	A	708	NAG	O7-C7-C8	-3.06	116.44	122.06
8	A	710	NAG	C4-C3-C2	-2.94	106.65	111.23
8	A	710	NAG	C3-C4-C5	-2.92	105.11	110.20
8	A	710	NAG	O7-C7-C8	-2.48	117.50	122.06
8	A	709	NAG	O4-C4-C5	-2.41	102.86	109.24
8	A	709	NAG	C4-C3-C2	-2.23	107.76	111.23
8	A	710	NAG	C6-C5-C4	-2.21	107.56	113.02
8	A	710	NAG	O3-C3-C2	-2.12	104.90	109.11
8	A	709	NAG	O7-C7-C8	-2.02	118.35	122.06
8	A	709	NAG	C8-C7-N2	2.02	119.97	116.11
8	A	708	NAG	C1-O5-C5	2.15	114.97	112.25
8	A	707	NAG	C3-C4-C5	2.22	114.07	110.20
8	A	710	NAG	O7-C7-N2	2.71	127.40	121.86
8	A	708	NAG	C4-C3-C2	2.73	115.47	111.23
8	A	710	NAG	C1-O5-C5	3.10	116.18	112.25
8	A	707	NAG	C1-O5-C5	4.01	117.34	112.25
8	A	710	NAG	C2-N2-C7	4.08	128.28	123.04
8	A	708	NAG	C2-N2-C7	4.37	128.65	123.04
8	A	709	NAG	C1-O5-C5	4.89	118.45	112.25
8	A	710	NAG	O4-C4-C3	5.80	123.40	110.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	707	NAG	1	0

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	CO3	A	702	3	0,3,3	0.00	-	0,3,3	0.00	-
5	SO4	A	703	-	4,4,4	0.67	0	6,6,6	0.74	0
7	NAG	A	706	1	14,14,15	1.28	1 (7%)	15,19,21	2.61	7 (46%)
9	FLP	A	711	-	16,19,19	2.97	1 (6%)	21,26,26	2.29	6 (28%)
10	GOL	A	712	-	5,5,5	0.57	0	5,5,5	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CO3	A	702	3	-	0/0/0/0	0/0/0/0
5	SO4	A	703	-	-	0/0/0/0	0/0/0/0
7	NAG	A	706	1	-	0/6/23/26	0/1/1/1
9	FLP	A	711	-	-	0/8/12/12	0/2/2/2
10	GOL	A	712	-	-	0/4/4/4	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	706	NAG	O7-C7	3.70	1.31	1.23
9	A	711	FLP	C6-C11	11.09	1.53	1.38

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	711	FLP	C10-C11-C6	-5.18	117.73	124.24
7	A	706	NAG	C8-C7-N2	-4.63	107.25	116.11
9	A	711	FLP	C7-C6-C2	-2.93	113.34	118.78
7	A	706	NAG	O3-C3-C4	-2.28	105.21	110.34
9	A	711	FLP	C14-C12-C9	2.02	116.30	112.32
7	A	706	NAG	C3-C2-N2	2.04	115.44	110.56
7	A	706	NAG	O5-C5-C6	2.19	112.09	107.35
9	A	711	FLP	C7-C6-C11	2.21	118.09	115.90
7	A	706	NAG	C1-O5-C5	4.12	117.48	112.25
9	A	711	FLP	C2-C6-C11	4.28	128.53	123.05
7	A	706	NAG	O7-C7-N2	4.71	131.48	121.86
7	A	706	NAG	C2-N2-C7	4.98	129.44	123.04
9	A	711	FLP	F-C11-C6	5.53	127.03	118.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	711	FLP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/335 (100%)	0.39	27 (8%)	15 13	11, 20, 49, 90	1 (0%)
2	B	6/6 (100%)	7.09	6 (100%)	0 0	36, 43, 102, 109	0
All	All	341/341 (100%)	0.51	33 (9%)	10 9	11, 20, 50, 109	1 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	681	LEU	21.8
1	A	342	TYR	14.7
1	A	422	SER	7.5
1	A	420	HIS	7.4
1	A	418	SER	7.2
1	A	423	LEU	6.9
2	B	682	GLU	6.4
1	A	421	SER	6.4
2	B	683	ALA	5.5
1	A	419	LYS	5.2
1	A	663	THR	3.9
2	B	685	ALA	3.9
1	A	417	SER	3.5
1	A	427	LEU	3.2
1	A	416	LYS	3.2
2	B	686	PHE	3.0
1	A	509	ASP	2.9
1	A	667	THR	2.9
1	A	429	PRO	2.9
1	A	635	GLU	2.8
1	A	510	GLN	2.7
1	A	627	ASP	2.6
1	A	431	GLU	2.5
1	A	621	ASN	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	640	LEU	2.4
1	A	609	GLN	2.3
1	A	628	LYS	2.3
1	A	347	TRP	2.2
1	A	373	THR	2.2
1	A	461	VAL	2.2
1	A	641	PHE	2.0
1	A	676	SER	2.0
2	B	684	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
8	NAG	A	707	14/15	0.88	0.14	0.70	31,35,42,46	0
8	NAG	A	710	14/15	0.85	0.20	-0.05	38,44,65,69	0
8	NAG	A	709	14/15	0.93	0.08	-0.41	24,27,35,36	0
8	NAG	A	708	14/15	0.93	0.31	-	43,52,62,68	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
9	FLP	A	711	18/18	0.55	0.39	4.60	32,54,60,61	18
10	GOL	A	712	6/6	0.87	0.16	3.78	30,39,43,49	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	NAG	A	706	14/15	0.79	0.22	0.37	34,47,56,67	0
4	CO3	A	702	4/4	0.98	0.13	-0.11	11,12,12,13	0
6	ZN	A	704	1/1	0.99	0.05	-0.79	17,17,17,17	0
3	FE	A	701	1/1	1.00	0.09	-2.26	11,11,11,11	0
5	SO4	A	703	5/5	0.98	0.17	-	37,38,40,51	0
6	ZN	A	705	1/1	0.97	0.05	-	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.