



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:43 AM GMT

PDB ID : 3FRZ
Title : Crystal Structure of HCV NS5B RNA polymerase in complex with PF868554
Authors : Parge, H.E.
Deposited on : 2009-01-08
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	AG6	A	578	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	562	4331	2730	758	812	31	0	1	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	GLN	LEU	ENGINEERED	UNP P26663
A	59	ASP	VAL	ENGINEERED	UNP P26663
A	101	TYR	PHE	ENGINEERED	UNP P26663
A	114	ARG	LYS	ENGINEERED	UNP P26663
A	544	GLN	ARG	variant	UNP P26663
A	571	HIS	-	EXPRESSION TAG	UNP P26663
A	572	HIS	-	EXPRESSION TAG	UNP P26663
A	573	HIS	-	EXPRESSION TAG	UNP P26663
A	574	HIS	-	EXPRESSION TAG	UNP P26663
A	575	HIS	-	EXPRESSION TAG	UNP P26663
A	576	HIS	-	EXPRESSION TAG	UNP P26663

- Molecule 2 is (6R)-6-CYCLOPENTYL-6-[2-(2,6-DIETHYLPYRIDIN-4-YL)ETHYL]-3-[(5,7-DIMETHYL[1,2,4]TRIAZOLO[1,5-A]PYRIMIDIN-2-YL)METHYL]-4-HYDROXY-5,6-DIHYDRO-2H-PYRAN-2-ONE (three-letter code: AG0) (formula: C₂₉H₃₇N₅O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	660	Total	O	0	0
			660	660		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	83.20Å 83.20Å 180.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.86 10.00 – 1.86	Depositor EDS
% Data completeness (in resolution range)	97.5 (10.00-1.86) 97.5 (10.00-1.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.07 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.175 , 0.221 0.187 , 0.231	Depositor DCC
R_{free} test set	2662 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	27.1	Xtrriage
Anisotropy	0.009	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 57.4	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Outliers	0 of 52265 reflections	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5060	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AG6, BME, AG0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.76	1/4432 (0.0%)	0.82	6/6027 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	139	MET	CG-SD	-5.39	1.67	1.81

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	ASP	CB-CG-OD2	6.63	124.27	118.30
1	A	510	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	A	375	ASP	CB-CG-OD2	5.73	123.45	118.30
1	A	387	ASP	CB-CG-OD2	5.51	123.26	118.30
1	A	332	ASP	CB-CG-OD1	5.35	123.11	118.30
1	A	517	ARG	NE-CZ-NH2	-5.01	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4331	0	4265	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	37	0	36	0	0
3	A	28	0	24	0	0
4	A	4	0	5	0	0
5	A	660	0	0	5	0
All	All	5060	0	4330	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:THR:HG23	1:A:278[B]:ARG:HH12	1.66	0.60
1:A:160:ILE:HD11	5:A:1086:HOH:O	2.05	0.56
1:A:523:LYS:HG3	1:A:534:LEU:HD12	1.90	0.54
1:A:406:ASN:C	1:A:406:ASN:OD1	2.51	0.47
1:A:445:CYS:SG	1:A:454:ILE:HD12	2.56	0.45
1:A:278[B]:ARG:NH2	5:A:1024:HOH:O	2.49	0.44
1:A:390:THR:HB	1:A:391:PRO:HD3	1.99	0.43
1:A:268:ASN:HB3	1:A:274:CYS:SG	2.59	0.43
1:A:533:LYS:NZ	5:A:1154:HOH:O	2.51	0.43
1:A:219:TYR:HB3	1:A:320:LEU:HD23	2.00	0.43
1:A:223:CYS:HB3	5:A:641:HOH:O	2.19	0.42
1:A:34:HIS:HD2	5:A:857:HOH:O	2.02	0.42
1:A:355:GLN:HA	1:A:356:PRO:HD3	1.92	0.41
1:A:144:VAL:HB	1:A:394:ARG:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	561/576 (97%)	554 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	464/491 (94%)	457 (98%)	7 (2%)	72	60

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	SER
1	A	113	SER
1	A	313	MET
1	A	380	ARG
1	A	419	LEU
1	A	490	ARG
1	A	545	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AG0	A	577	-	34,41,41	1.81	6 (17%)	36,60,60	1.84	13 (36%)
3	AG6	A	578	1	29,29,47	2.01	5 (17%)	33,36,60	1.55	6 (18%)
4	BME	A	579	1	3,3,3	1.05	0	2,2,2	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AG0	A	577	-	-	0/20/45/45	0/5/5/5
3	AG6	A	578	1	-	1/27/27/46	0/2/2/3
4	BME	A	579	1	-	0/1/1/1	0/0/0/0

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	577	AG0	C4-N10	-5.35	1.27	1.35
2	A	577	AG0	O16-C15	-2.76	1.43	1.47
2	A	577	AG0	C9-N8	-2.76	1.31	1.33
3	A	578	AG6	C30-C23	-2.18	1.46	1.49
3	A	578	AG6	C31-C30	2.01	1.42	1.39
2	A	577	AG0	C30-C29	2.18	1.43	1.39
3	A	578	AG6	C22-C23	2.78	1.52	1.47
2	A	577	AG0	C21-C15	3.77	1.56	1.52
3	A	578	AG6	C8-N2	3.81	1.49	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	577	AG0	C1-C18	5.00	1.58	1.51
3	A	578	AG6	C22-C15	8.11	1.53	1.32

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	578	AG6	O5-C7-N2	-3.16	120.21	124.76
3	A	578	AG6	C26-C9-C8	-3.13	104.31	113.41
2	A	577	AG0	C29-C30-C31	-2.84	118.20	120.24
2	A	577	AG0	C2-N3-C4	-2.68	114.72	117.52
2	A	577	AG0	C31-N32-C33	-2.54	115.00	118.12
2	A	577	AG0	C28-C29-C30	-2.52	115.24	118.96
2	A	577	AG0	O19-C17-C18	-2.48	119.68	125.28
2	A	577	AG0	C34-C33-C28	-2.23	117.56	122.70
3	A	578	AG6	C11-C15-C22	-2.08	115.53	126.58
2	A	577	AG0	C11-C2-C7	-2.00	119.07	121.75
3	A	578	AG6	C34-C35-C33	2.03	123.49	119.93
2	A	577	AG0	C30-C31-N32	2.10	125.33	122.96
2	A	577	AG0	C23-C27-C15	2.62	116.90	112.24
2	A	577	AG0	C14-C13-C18	2.78	126.63	123.11
3	A	578	AG6	C31-C30-C32	2.83	122.81	118.60
3	A	578	AG6	O6-C10-N3	3.16	129.12	122.93
2	A	577	AG0	C15-O16-C17	3.20	127.52	121.07
2	A	577	AG0	O16-C17-O19	3.69	122.23	117.72
2	A	577	AG0	C29-C28-C33	3.71	122.90	120.24

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	578	AG6	C11-C15-C22-C23

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	562/576 (97%)	-0.07	16 (2%) 56 54	20, 28, 44, 57	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	548	SER	4.9
1	A	404	PRO	3.8
1	A	543	SER	3.6
1	A	546	ASP	3.5
1	A	402	HIS	3.4
1	A	310	ASP	2.9
1	A	95	HIS	2.9
1	A	149	PRO	2.8
1	A	247	PRO	2.7
1	A	102	GLY	2.5
1	A	532	THR	2.5
1	A	547	LEU	2.5
1	A	113	SER	2.3
1	A	377	SER	2.1
1	A	152	GLY	2.1
1	A	110	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	AG6	A	578	28/45	0.79	0.19	2.54	39,46,53,54	0
2	AG0	A	577	37/37	0.94	0.09	-0.40	22,28,38,41	0
4	BME	A	579	4/4	0.79	0.29	-	50,55,57,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.