



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:28 PM GMT

PDB ID : 1FSX
Title : THE X-RAY STRUCTURE DETERMINATION OF BOVINE CARBON-MONOXY HB AT 2.1 Å RESOLUTION AND ITS RELATIONSHIP TO THE QUATERNARY STRUCTURE OF OTHER HB CRYSTAL FORMS
Authors : Safo, M.K.; Abraham, D.J.
Deposited on : 2000-09-11
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	<div> <div>2%</div> <div>87%</div> <div>11%</div> <div>.</div> </div>
1	C	141	<div> <div>2%</div> <div>82%</div> <div>13%</div> <div>..</div> </div>
2	B	145	<div> <div>2%</div> <div>73%</div> <div>23%</div> <div>..</div> </div>
2	D	145	<div> <div>3%</div> <div>79%</div> <div>17%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4702 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HEMOGLOBIN ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	141	Total	C	N	O	S	0	0	0
			1066	683	186	196	1			
1	C	141	Total	C	N	O	S	0	0	0
			1066	683	186	196	1			

- Molecule 2 is a protein called HEMOGLOBIN BETA CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	145	Total	C	N	O	S	0	0	0
			1127	726	194	203	4			
2	D	145	Total	C	N	O	S	0	0	0
			1127	726	194	203	4			

- Molecule 3 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 2	C 1	O 1	0	0
3	B	1	Total 2	C 1	O 1	0	0
3	C	1	Total 2	C 1	O 1	0	0
3	D	1	Total 2	C 1	O 1	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
4	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
4	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
4	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	43	Total O 43 43	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	26	Total 26	O 26	0	0
5	C	38	Total 38	O 38	0	0
5	D	29	Total 29	O 29	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HEMOGLOBIN ALPHA CHAIN




- Molecule 1: HEMOGLOBIN ALPHA CHAIN



- Molecule 2: HEMOGLOBIN BETA CHAIN



- Molecule 2: HEMOGLOBIN BETA CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.41 Å 110.63 Å 65.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.50 – 2.10 64.51 – 1.94	Depositor EDS
% Data completeness (in resolution range)	100.0 (64.50-2.10) 89.6 (64.51-1.94)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.84 (at 1.94 Å)	Xtriage
Refinement program	X-PLOR 3.843	Depositor
R, R_{free}	0.205 , 0.254 0.207 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	38.3	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 52.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 72194 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4702	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/1093	0.64	1/1483 (0.1%)
1	C	0.51	0/1093	0.65	3/1483 (0.2%)
2	B	0.51	0/1152	0.63	0/1556
2	D	0.55	1/1152 (0.1%)	0.66	0/1556
All	All	0.51	1/4490 (0.0%)	0.64	4/6078 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	693	CYS	CB-SG	5.14	1.91	1.82

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	540	TYR	N-CA-C	6.00	127.21	111.00
1	A	47	ASP	N-CA-C	-5.55	96.01	111.00
1	C	539	LYS	C-N-CA	5.04	134.29	121.70
1	C	447	ASP	N-CA-C	-5.03	97.42	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	541	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1066	0	1069	9	0
1	C	1066	0	1066	17	0
2	B	1127	0	1129	23	0
2	D	1127	0	1129	21	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	43	0	30	0	0
4	B	43	0	30	0	0
4	C	43	0	30	0	0
4	D	43	0	30	2	0
5	A	43	0	0	1	0
5	B	26	0	0	0	0
5	C	38	0	0	1	0
5	D	29	0	0	2	0
All	All	4702	0	4513	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:541:ARG:HD2	5:C:809:HOH:O	1.57	1.02
1:C:541:ARG:HG3	1:C:541:ARG:NH1	1.75	0.99
1:C:541:ARG:HH11	1:C:541:ARG:HG3	1.23	0.98
2:D:693:CYS:SG	2:D:745:TYR:CE2	2.67	0.87
1:C:537:THR:O	1:C:541:ARG:O	1.97	0.82
2:D:602:MET:SD	2:D:603:LEU:N	2.55	0.80
2:D:666:LYS:HD3	4:D:748:HEM:HBA2	1.73	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:693:CYS:SG	2:D:745:TYR:CD2	2.82	0.70
1:C:539:LYS:HG3	1:C:539:LYS:O	1.93	0.68
2:B:203:LEU:HD12	2:B:207:GLU:HB3	1.80	0.64
2:D:721:GLU:HG2	5:D:849:HOH:O	1.97	0.64
2:D:621:ASP:HA	2:D:665:LYS:HD3	1.81	0.63
2:D:688:LEU:HD21	4:D:748:HEM:HMA1	1.82	0.62
1:A:141:ARG:HE	1:C:527:LYS:HD3	1.68	0.58
1:A:82:GLU:H	1:A:82:GLU:CD	2.05	0.58
2:B:290:GLU:HG2	2:B:344:ARG:HE	1.68	0.57
2:B:203:LEU:HD12	2:B:207:GLU:CB	2.35	0.56
1:C:438:THR:O	1:C:441:THR:HB	2.06	0.55
2:D:631:LEU:HD22	2:D:706:LEU:HG	1.91	0.53
1:C:503:HIS:HE1	2:D:731:GLN:OE1	1.93	0.52
1:C:541:ARG:HH11	1:C:541:ARG:CG	2.10	0.52
2:B:285:PHE:HD2	2:B:288:LEU:HD12	1.75	0.52
2:B:203:LEU:HB2	2:B:207:GLU:OE2	2.09	0.52
1:C:470:VAL:HA	1:C:473:LEU:HD13	1.91	0.51
1:A:103:HIS:HD2	5:A:838:HOH:O	1.93	0.51
1:C:489:HIS:HB2	1:C:539:LYS:HD3	1.94	0.50
1:A:139:LYS:HD3	1:A:139:LYS:O	2.11	0.50
2:D:651:ALA:O	2:D:655:MET:HG2	2.12	0.49
1:C:446:PHE:HB3	1:C:448:LEU:HD13	1.94	0.49
2:D:720:LYS:NZ	2:D:720:LYS:HA	2.27	0.49
1:C:538:SER:C	1:C:540:TYR:H	2.15	0.49
2:B:267:VAL:O	2:B:270:SER:HB3	2.13	0.49
2:D:637:TRP:HE1	2:D:702:ASN:HD21	1.61	0.49
2:B:237:TRP:HE1	2:B:302:ASN:HD21	1.60	0.48
2:B:231:LEU:HD22	2:B:306:LEU:HD23	1.94	0.48
2:B:307:GLY:HA3	2:B:334:VAL:HG13	1.95	0.48
2:D:663:HIS:O	2:D:667:VAL:HG23	2.14	0.47
2:B:220:VAL:HG13	2:B:268:LEU:CB	2.46	0.46
2:B:300:PRO:HA	2:B:303:PHE:CE2	2.50	0.46
2:D:700:PRO:HG3	2:D:745:TYR:CD2	2.50	0.46
2:D:671:PHE:O	2:D:675:MET:HG2	2.15	0.46
2:B:208:LYS:HB2	2:B:208:LYS:HE3	1.75	0.45
2:B:220:VAL:HG13	2:B:268:LEU:HB3	1.98	0.45
1:A:125:LEU:HD13	1:A:125:LEU:HA	1.78	0.45
2:D:666:LYS:HA	2:D:666:LYS:HE2	2.00	0.43
1:C:403:SER:O	1:C:407:LYS:HG3	2.18	0.43
2:B:250:THR:HG22	2:B:252:ASP:H	1.84	0.43
1:A:106:LEU:HD21	1:A:125:LEU:HB3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:LEU:O	1:A:7:LYS:HE3	2.18	0.43
1:A:94:ASP:HB3	1:A:97:ASN:ND2	2.34	0.43
2:B:278:LEU:HA	2:B:278:LEU:HD12	1.86	0.43
2:B:296:LEU:O	2:B:297:HIS:HB2	2.19	0.43
2:D:727:GLN:O	2:D:731:GLN:HG2	2.18	0.43
2:D:720:LYS:HZ2	2:D:720:LYS:HA	1.82	0.43
2:B:251:ALA:O	2:B:255:MET:HG2	2.19	0.43
2:B:204:THR:OG1	2:B:207:GLU:HG3	2.18	0.43
1:C:439:THR:HG22	1:C:497:ASN:OD1	2.18	0.42
2:B:211:VAL:HG12	2:B:212:THR:N	2.35	0.42
2:B:257:ASN:HA	2:B:258:PRO:HD3	1.91	0.42
2:D:688:LEU:HA	2:D:688:LEU:HD12	1.85	0.41
2:D:678:LEU:HD23	2:D:678:LEU:HA	1.88	0.41
1:C:512:HIS:C	1:C:514:PRO:HD3	2.40	0.41
1:C:476:LEU:HD12	1:C:476:LEU:HA	1.89	0.41
2:D:725:VAL:HG22	5:D:899:HOH:O	2.21	0.41
2:B:285:PHE:HB3	2:B:341:LEU:HD22	2.03	0.41
1:A:95:PRO:HA	1:A:98:PHE:HD2	1.85	0.41
2:B:281:LEU:HA	2:B:281:LEU:HD12	1.94	0.40
2:B:268:LEU:HA	2:B:271:PHE:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	136 (98%)	3 (2%)	0	100	100
1	C	139/141 (99%)	136 (98%)	3 (2%)	0	100	100
2	B	143/145 (99%)	140 (98%)	2 (1%)	1 (1%)	26	21
2	D	143/145 (99%)	139 (97%)	3 (2%)	1 (1%)	26	21
All	All	564/572 (99%)	551 (98%)	11 (2%)	2 (0%)	39	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	278	LEU
2	D	603	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/112 (100%)	104 (93%)	8 (7%)	18	14
1	C	112/112 (100%)	103 (92%)	9 (8%)	15	11
2	B	118/118 (100%)	107 (91%)	11 (9%)	11	7
2	D	118/118 (100%)	108 (92%)	10 (8%)	13	9
All	All	460/460 (100%)	422 (92%)	38 (8%)	14	9

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	LEU
1	A	66	LEU
1	A	73	LEU
1	A	82	GLU
1	A	109	LEU
1	A	125	LEU
1	A	140	TYR
1	A	141	ARG
2	B	202	MET
2	B	203	LEU
2	B	221	ASP
2	B	268	LEU
2	B	271	PHE
2	B	278	LEU
2	B	281	LEU
2	B	333	VAL
2	B	341	LEU
2	B	343	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	345	TYR
1	C	448	LEU
1	C	473	LEU
1	C	476	LEU
1	C	505	LEU
1	C	525	LEU
1	C	531	ASN
1	C	539	LYS
1	C	540	TYR
1	C	541	ARG
2	D	602	MET
2	D	606	GLU
2	D	643	GLU
2	D	644	SER
2	D	666	LYS
2	D	668	LEU
2	D	706	LEU
2	D	710	LEU
2	D	720	LYS
2	D	741	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	273	ASN
2	B	302	ASN
2	B	339	ASN
1	C	409	ASN
1	C	472	HIS
1	C	503	HIS
2	D	702	ASN
2	D	746	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CMO	A	142	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	A	143	1,3	30,50,50	2.72	12 (40%)	24,82,82	2.56	8 (33%)
3	CMO	B	347	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	B	348	3,2	30,50,50	2.79	12 (40%)	24,82,82	2.46	9 (37%)
3	CMO	C	542	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	C	543	1,3	30,50,50	3.37	13 (43%)	24,82,82	2.46	8 (33%)
3	CMO	D	747	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	D	748	3,2	30,50,50	2.98	12 (40%)	24,82,82	2.32	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CMO	A	142	4	-	0/0/0/0	0/0/0/0
4	HEM	A	143	1,3	-	0/10/54/54	0/0/8/8
3	CMO	B	347	4	-	0/0/0/0	0/0/0/0
4	HEM	B	348	3,2	-	0/10/54/54	0/0/8/8
3	CMO	C	542	4	-	0/0/0/0	0/0/0/0
4	HEM	C	543	1,3	-	0/10/54/54	0/0/8/8
3	CMO	D	747	4	-	0/0/0/0	0/0/0/0
4	HEM	D	748	3,2	-	0/10/54/54	0/0/8/8

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	543	HEM	C3B-C4B	-9.78	1.43	1.51
4	C	543	HEM	C3B-CAB	-7.52	1.37	1.51
4	B	348	HEM	C3D-C4D	-7.12	1.42	1.51
4	D	748	HEM	C3D-C4D	-6.82	1.42	1.51
4	C	543	HEM	C3C-CAC	-6.70	1.38	1.51
4	D	748	HEM	C2D-C3D	-6.65	1.34	1.54
4	D	748	HEM	C3C-CAC	-6.42	1.39	1.51
4	D	748	HEM	C2C-C1C	-6.21	1.40	1.52
4	B	348	HEM	C3C-CAC	-6.02	1.40	1.51
4	A	143	HEM	C2D-C3D	-5.64	1.37	1.54
4	B	348	HEM	C2D-C3D	-5.61	1.37	1.54
4	C	543	HEM	C2D-C3D	-5.47	1.38	1.54
4	B	348	HEM	C3B-CAB	-5.31	1.41	1.51
4	A	143	HEM	C3B-CAB	-5.27	1.41	1.51
4	D	748	HEM	C3B-CAB	-5.14	1.41	1.51
4	A	143	HEM	C3B-C4B	-5.05	1.47	1.51
4	C	543	HEM	C3D-C4D	-4.40	1.45	1.51
4	A	143	HEM	C3C-CAC	-4.15	1.43	1.51
4	C	543	HEM	C2C-C1C	-4.13	1.44	1.52
4	D	748	HEM	C3B-C4B	-4.09	1.48	1.51
4	B	348	HEM	C2C-C1C	-3.74	1.45	1.52
4	B	348	HEM	C3B-C4B	-3.43	1.48	1.51
4	C	543	HEM	C2B-C1B	-3.07	1.41	1.51
4	B	348	HEM	C2B-C1B	-2.62	1.43	1.51
4	A	143	HEM	C3D-C4D	-2.45	1.48	1.51
4	A	143	HEM	C2B-C1B	-2.24	1.44	1.51
4	D	748	HEM	C2B-C1B	-2.16	1.44	1.51
4	C	543	HEM	C2D-C1D	-2.14	1.44	1.51
4	D	748	HEM	C2D-C1D	-2.00	1.45	1.51
4	C	543	HEM	CBC-CAC	2.34	1.42	1.29
4	B	348	HEM	C4C-NC	2.36	1.38	1.36
4	D	748	HEM	CHD-C4C	2.42	1.42	1.36
4	D	748	HEM	C1C-NC	2.48	1.39	1.36
4	C	543	HEM	CHC-C1C	2.52	1.42	1.36
4	A	143	HEM	CBB-CAB	2.54	1.44	1.29
4	C	543	HEM	C4C-NC	2.55	1.39	1.36
4	A	143	HEM	FE-ND	2.80	2.12	1.97
4	D	748	HEM	CBC-CAC	2.81	1.45	1.29
4	C	543	HEM	FE-NB	2.85	2.12	1.97
4	A	143	HEM	CHD-C4C	2.85	1.43	1.36
4	B	348	HEM	CHD-C4C	2.87	1.43	1.36
4	B	348	HEM	FE-NC	2.87	2.07	1.95
4	B	348	HEM	CBC-CAC	2.91	1.46	1.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	748	HEM	CBB-CAB	2.92	1.46	1.29
4	B	348	HEM	CBB-CAB	2.97	1.46	1.29
4	A	143	HEM	CBC-CAC	3.00	1.46	1.29
4	C	543	HEM	FE-NC	4.75	2.14	1.95
4	A	143	HEM	C1C-NC	5.22	1.42	1.36
4	A	143	HEM	C4C-NC	6.10	1.43	1.36

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	543	HEM	CBA-CAA-C2A	-4.34	104.75	112.53
4	A	143	HEM	CAA-C2A-C1A	-2.99	123.77	127.01
4	D	748	HEM	CBA-CAA-C2A	-2.66	107.77	112.53
4	D	748	HEM	CMD-C2D-C3D	2.02	123.29	114.35
4	B	348	HEM	C3B-CAB-CBB	2.31	128.00	124.46
4	A	143	HEM	C2D-C3D-C4D	2.36	105.50	101.50
4	D	748	HEM	C3B-CAB-CBB	2.61	128.46	124.46
4	B	348	HEM	CMD-C2D-C3D	2.64	126.04	114.35
4	C	543	HEM	C4B-CHC-C1C	2.79	130.48	125.82
4	D	748	HEM	C3B-C4B-CHC	2.93	127.29	123.16
4	A	143	HEM	CMD-C2D-C3D	2.95	127.39	114.35
4	C	543	HEM	CAD-C3D-C4D	2.95	122.89	112.47
4	B	348	HEM	C2C-C1C-CHC	3.12	128.42	123.68
4	C	543	HEM	CMD-C2D-C3D	3.13	128.20	114.35
4	D	748	HEM	CMB-C2B-C3B	3.16	124.41	116.53
4	D	748	HEM	CAD-C3D-C4D	3.16	123.62	112.47
4	B	348	HEM	CAD-C3D-C4D	3.31	124.14	112.47
4	D	748	HEM	C2D-C3D-C4D	3.50	107.43	101.50
4	B	348	HEM	C3C-CAC-CBC	3.64	130.03	124.46
4	A	143	HEM	CAD-C3D-C4D	3.91	126.28	112.47
4	B	348	HEM	CMC-C2C-C3C	4.10	126.77	116.53
4	A	143	HEM	C3B-CAB-CBB	4.12	130.78	124.46
4	B	348	HEM	CAD-C3D-C2D	4.12	125.07	113.22
4	D	748	HEM	CMC-C2C-C3C	4.26	127.18	116.53
4	B	348	HEM	CMB-C2B-C3B	4.37	127.44	116.53
4	C	543	HEM	C2D-C3D-C4D	4.68	109.44	101.50
4	C	543	HEM	CMB-C2B-C3B	4.74	128.36	116.53
4	C	543	HEM	CMC-C2C-C3C	4.78	128.46	116.53
4	C	543	HEM	CAD-C3D-C2D	5.03	127.67	113.22
4	A	143	HEM	CAD-C3D-C2D	5.21	128.20	113.22
4	B	348	HEM	C2D-C3D-C4D	5.41	110.67	101.50
4	D	748	HEM	CAD-C3D-C2D	5.44	128.86	113.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	143	HEM	CMB-C2B-C3B	5.49	130.25	116.53
4	A	143	HEM	CMC-C2C-C3C	5.70	130.76	116.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	748	HEM	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	141/141 (100%)	-0.29	3 (2%) 67 72	22, 33, 51, 70	0
1	C	141/141 (100%)	-0.18	3 (2%) 67 72	20, 33, 53, 71	0
2	B	145/145 (100%)	-0.12	3 (2%) 67 72	22, 39, 58, 78	0
2	D	145/145 (100%)	-0.16	4 (2%) 56 64	22, 37, 54, 70	0
All	All	572/572 (100%)	-0.19	13 (2%) 64 70	20, 36, 56, 78	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	541	ARG	13.3
2	B	346	HIS	5.6
1	A	140	TYR	4.3
1	C	540	TYR	4.3
2	D	602	MET	4.0
2	D	746	HIS	4.0
2	D	745	TYR	3.9
2	B	202	MET	3.5
1	C	401	VAL	3.4
1	A	1	VAL	3.1
2	D	744	ARG	2.5
2	B	345	TYR	2.4
1	A	141	ARG	2.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CMO	A	142	2/2	0.98	0.12	1.24	27,27,27,33	0
3	CMO	D	747	2/2	0.98	0.13	1.13	32,32,32,40	0
4	HEM	A	143	43/43	0.97	0.10	-0.05	22,33,50,56	0
4	HEM	B	348	43/43	0.97	0.10	-0.21	24,35,56,61	0
4	HEM	C	543	43/43	0.97	0.09	-0.49	24,33,51,54	0
3	CMO	C	542	2/2	0.96	0.08	-0.64	27,27,27,31	0
4	HEM	D	748	43/43	0.97	0.09	-0.65	21,34,55,62	0
3	CMO	B	347	2/2	0.99	0.14	-	33,33,33,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.