



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 24, 2017 – 01:42 PM EST

PDB ID : 1FTX  
Title : Crystal structure of alanine racemase in complex with D-alanine phosphonate  
Authors : Stamper, G.F.; Ringe, D.  
Deposited on : 2000-09-13  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442

i



## X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	387		..
1	B	387		..

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ALANINE RACEMASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	0	0
			3024	1936	536	539	13			
1	B	380	Total	C	N	O	S	0	0	0
			3024	1936	536	539	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	129	KCX	LYS	MODIFIED RESIDUE	UNP P10724
B	129	KCX	LYS	MODIFIED RESIDUE	UNP P10724

- Molecule 2 is (1S)-1-[(1E)-{3-HYDROXY-2-METHYL-5-[(PHOSPHONOOXY)METHYL]PYRIDIN-4-YL}METHYLENE)AMINO]ETHYLPHOSPHONIC ACID (three-letter code: EPC) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>2</sub>O<sub>8</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			22	10	2	8	2		
2	B	1	Total	C	N	O	P	0	0
			22	10	2	8	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	131	Total	O	0	0
			131	131		
3	B	137	Total	O	0	0
			137	137		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ALANINE RACEMASE



#### • Molecule 1: ALANINE RACEMASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.76Å 89.93Å 85.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.52 – 2.20 29.52 – 2.20	Depositor EDS
% Data completeness (in resolution range)	90.8 (29.52-2.20) 90.6 (29.52-2.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.56 (at 2.20Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, $R_{free}$	0.183 , 0.211 0.183 , 0.210	Depositor DCC
$R_{free}$ test set	2151 reflections (6.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.6	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 69.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6360	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPC, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/3089	0.66	0/4199
1	B	0.38	0/3089	0.68	1/4199 (0.0%)
All	All	0.38	0/6178	0.67	1/8398 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	27	LEU	CA-CB-CG	5.15	127.15	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3024	0	3010	45	0
1	B	3024	0	3011	58	0
2	A	22	0	12	1	0
2	B	22	0	12	0	0
3	A	131	0	0	4	0
3	B	137	0	0	5	0
All	All	6360	0	6045	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:129:KCX:C	1:B:130:MET:N	2.06	1.17
1:A:129:KCX:C	1:A:130:MET:N	2.12	1.12
1:A:351:THR:HG21	1:A:355:GLU:OE1	1.71	0.91
1:A:163:LEU:HD21	1:A:189:LEU:HD22	1.59	0.83
1:A:163:LEU:HD21	1:A:189:LEU:CD2	2.10	0.81
1:A:198:LEU:HA	1:A:216:ASN:HD21	1.51	0.76
1:B:3:ASP:HB2	3:B:553:HOH:O	1.84	0.76
1:B:129:KCX:C	1:B:130:MET:CA	2.64	0.75
1:B:163:LEU:HD13	1:B:192:LEU:HD11	1.70	0.74
1:B:130:MET:N	1:B:164:TYR:O	2.25	0.70
1:A:163:LEU:HD22	1:A:192:LEU:CD1	2.23	0.69
1:A:97:ALA:HB2	1:A:124:ILE:HG12	1.75	0.68
1:B:129:KCX:C	1:B:130:MET:HA	2.23	0.67
1:B:38:VAL:HG13	1:B:38:VAL:O	1.93	0.67
1:B:212:ASP:OD1	1:B:213:ARG:HG3	1.95	0.66
1:A:163:LEU:HD22	1:A:192:LEU:HD11	1.78	0.66
1:B:134:MET:HG2	1:B:168:ALA:HB2	1.78	0.65
1:A:129:KCX:C	1:A:130:MET:CA	2.74	0.65
1:A:314:GLN:HG2	3:B:405:HOH:O	2.03	0.59
1:A:130:MET:N	1:A:164:TYR:O	2.36	0.59
1:B:97:ALA:HB2	1:B:124:ILE:HG12	1.84	0.59
1:A:129:KCX:C	1:A:130:MET:HA	2.33	0.59
1:B:64:VAL:HG13	1:B:69:GLU:HB2	1.85	0.58
1:B:111:LEU:HD11	1:B:148:ILE:HG23	1.85	0.58
1:B:163:LEU:HD11	1:B:189:LEU:HD23	1.86	0.57
1:A:66:PHE:HZ	1:B:363:ARG:HG2	1.69	0.56
1:A:9:TRP:CD1	1:A:367:ILE:HD12	2.40	0.56
1:B:151:LEU:CD2	1:B:154:ARG:HH21	2.18	0.56
1:B:6:ARG:HD3	1:B:8:THR:HB	1.88	0.54
1:A:39:LYS:HE2	1:B:313:ASP:OD2	2.08	0.53
1:A:296:HIS:NE2	1:A:303:LYS:HE3	2.24	0.52
1:A:134:MET:HB3	3:A:419:HOH:O	2.08	0.51
1:A:351:THR:HG22	1:B:290:ARG:HE	1.74	0.51
1:A:314:GLN:HG3	1:B:136:ARG:HD2	1.92	0.51
1:A:351:THR:HG23	3:A:496:HOH:O	2.11	0.51
1:B:282:ILE:HG21	1:B:331:LEU:HD21	1.92	0.51
1:B:151:LEU:HD23	1:B:154:ARG:HH21	1.75	0.51

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:38:VAL:CG1	1:B:38:VAL:O	2.57	0.50
1:A:130:MET:SD	1:A:163:LEU:HD11	2.52	0.50
1:B:293:GLN:HG3	1:B:308:GLY:O	2.11	0.49
1:B:339:VAL:CG2	3:B:618:HOH:O	2.60	0.49
1:B:48:VAL:HG11	1:B:76:LYS:HB3	1.94	0.49
1:B:331:LEU:HD12	1:B:331:LEU:N	2.28	0.49
1:B:161:GLU:O	1:B:197:PRO:HG2	2.14	0.48
1:A:363:ARG:HG2	1:B:66:PHE:HZ	1.78	0.48
1:A:296:HIS:CD2	1:A:303:LYS:HE3	2.48	0.48
1:A:98:GLN:HG2	1:A:122:PHE:CE2	2.50	0.47
1:A:17:ILE:HD13	1:A:50:VAL:HG22	1.96	0.47
1:B:222:ILE:HG13	1:B:227:LEU:HB2	1.96	0.47
1:A:331:LEU:HD22	1:A:331:LEU:N	2.30	0.47
1:B:293:GLN:O	1:B:306:ILE:O	2.33	0.46
1:B:64:VAL:CG1	1:B:69:GLU:HB2	2.45	0.46
1:A:229:PRO:HB2	1:A:233:ILE:HD11	1.97	0.46
1:A:14:LEU:HD23	1:A:17:ILE:HD12	1.97	0.46
1:A:93:ALA:HB1	1:A:124:ILE:HD13	1.96	0.46
1:B:291:ARG:HH11	1:B:291:ARG:HB2	1.81	0.46
1:B:164:TYR:HB2	1:B:200:HIS:O	2.16	0.45
1:B:174:ASN:ND2	1:B:176:ASP:OD1	2.49	0.45
1:B:251:LEU:HD21	1:B:317:ILE:HD11	1.97	0.45
1:B:155:HIS:HB3	1:B:158:PHE:HB2	1.97	0.45
1:A:350:GLU:HB2	1:B:291:ARG:HH22	1.81	0.45
1:B:229:PRO:HD2	1:B:233:ILE:HG21	2.00	0.44
1:A:127:HIS:ND1	1:A:162:GLY:HA3	2.32	0.44
1:A:344:ASP:OD1	1:A:347:ARG:NH2	2.51	0.44
1:B:198:LEU:HA	1:B:216:ASN:HD21	1.82	0.43
1:B:187:HIS:HB3	3:B:619:HOH:O	2.18	0.43
1:B:29:PRO:HB2	1:B:31:ASP:OD1	2.18	0.43
1:B:222:ILE:HG23	1:B:223:ALA:N	2.33	0.43
1:B:252:VAL:HG22	1:B:279:THR:O	2.18	0.43
1:B:35:MET:SD	1:B:63:ALA:HB2	2.58	0.43
1:A:253:HIS:HD2	1:B:107:ARG:HH11	1.67	0.43
1:B:183:THR:O	1:B:187:HIS:HD2	2.01	0.43
1:A:129:KCX:OQ2	1:A:136:ARG:HB3	2.18	0.43
1:B:200:HIS:CD2	1:B:200:HIS:N	2.87	0.43
1:B:4:PHE:N	1:B:4:PHE:CD1	2.87	0.43
1:B:195:ARG:HE	1:B:199:VAL:HG21	1.84	0.42
1:B:199:VAL:H	1:B:216:ASN:HD21	1.67	0.42
1:B:6:ARG:HD2	3:B:508:HOH:O	2.17	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:GLU:O	1:B:209:ARG:NH2	2.52	0.42
1:B:269:TYR:CG	1:B:307:VAL:HG13	2.54	0.42
1:B:293:GLN:HG2	1:B:293:GLN:O	2.19	0.42
1:A:234:LYS:N	1:A:235:PRO:HD2	2.35	0.42
1:A:219:ARG:HB2	2:A:1001:EPC:HC61	2.02	0.42
1:B:186:LEU:HA	1:B:189:LEU:HD12	2.00	0.42
1:A:290:ARG:NH1	3:A:582:HOH:O	2.48	0.42
1:A:331:LEU:N	1:A:331:LEU:CD2	2.83	0.41
1:B:38:VAL:HG22	1:B:42:ALA:HB2	2.01	0.41
1:A:268:THR:HB	3:A:491:HOH:O	2.20	0.41
1:B:288:TRP:CE2	1:B:331:LEU:HD23	2.55	0.41
1:A:173:VAL:HG23	1:A:174:ASN:N	2.36	0.41
1:A:47:ASP:OD1	1:A:48:VAL:N	2.53	0.41
1:A:272:GLN:HB2	1:A:272:GLN:HE21	1.60	0.40
1:A:269:TYR:CE1	1:A:307:VAL:HG22	2.57	0.40
1:A:231:PRO:HA	1:A:234:LYS:HD2	2.03	0.40
1:A:209:ARG:C	1:A:211:PRO:HD3	2.42	0.40
1:B:139:VAL:CG2	1:B:144:GLU:HB3	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/387 (97%)	364 (97%)	12 (3%)	0	100	100
1	B	376/387 (97%)	361 (96%)	14 (4%)	1 (0%)	46	50
All	All	752/774 (97%)	725 (96%)	26 (4%)	1 (0%)	56	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	133	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/321 (99%)	302 (95%)	15 (5%)	32	39
1	B	317/321 (99%)	295 (93%)	22 (7%)	19	20
All	All	634/642 (99%)	597 (94%)	37 (6%)	25	28

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	26	ARG
1	A	37	VAL
1	A	39	LYS
1	A	98	GLN
1	A	107	ARG
1	A	172	GLU
1	A	184	ARG
1	A	195	ARG
1	A	272	GLN
1	A	273	THR
1	A	331	LEU
1	A	335	GLN
1	A	348	HIS
1	A	351	THR
1	B	6	ARG
1	B	27	LEU
1	B	38	VAL
1	B	48	VAL
1	B	64	VAL
1	B	134	MET
1	B	141	ASP
1	B	143	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	147	ARG
1	B	153	GLU
1	B	163	LEU
1	B	184	ARG
1	B	195	ARG
1	B	216	ASN
1	B	237	LEU
1	B	262	LYS
1	B	273	THR
1	B	302	GLN
1	B	307	VAL
1	B	335	GLN
1	B	337	ASP
1	B	379	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	216	ASN
1	A	253	HIS
1	A	272	GLN
1	A	293	GLN
1	A	314	GLN
1	B	23	ASN
1	B	187	HIS
1	B	216	ASN
1	B	294	HIS
1	B	348	HIS
1	B	379	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	KCX	A	129	1	6,11,12	0.78	0	7,12,14	1.75	1 (14%)
1	KCX	B	129	1	6,11,12	0.97	0	7,12,14	2.90	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	129	1	-	0/6/10/12	0/0/0/0
1	KCX	B	129	1	-	0/6/10/12	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	KCX	CE-NZ-CX	-4.27	119.04	123.53
1	B	129	KCX	CG-CD-CE	-2.20	104.25	113.69
1	B	129	KCX	CE-NZ-CX	6.97	130.84	123.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	129	KCX	4	0
1	B	129	KCX	3	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EPC	A	1001	-	21,22,22	4.04	9 (42%)	26,33,33	1.20	1 (3%)
2	EPC	B	1002	-	21,22,22	4.71	11 (52%)	26,33,33	1.37	4 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EPC	A	1001	-	-	0/15/17/17	0/1/1/1
2	EPC	B	1002	-	-	0/15/17/17	0/1/1/1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1002	EPC	P2-C	-6.60	1.76	1.84
2	B	1002	EPC	P2-O6	-6.22	1.45	1.54
2	A	1001	EPC	P2-C	-4.90	1.78	1.84
2	A	1001	EPC	P2-O6	-4.64	1.47	1.54
2	A	1001	EPC	C2A-C2	2.02	1.54	1.50
2	B	1002	EPC	C6-C5	2.12	1.42	1.37
2	B	1002	EPC	C4-C5	2.24	1.45	1.42
2	B	1002	EPC	C-N2	2.94	1.49	1.46
2	B	1002	EPC	P2-O8	4.10	1.61	1.54
2	A	1001	EPC	C3-C2	4.36	1.43	1.40
2	A	1001	EPC	P2-O8	4.52	1.61	1.54
2	B	1002	EPC	P2-O7	6.74	1.62	1.49
2	B	1002	EPC	P1-O5	6.80	1.78	1.54
2	A	1001	EPC	P1-O4	6.85	1.78	1.54
2	B	1002	EPC	P1-O4	7.16	1.79	1.54
2	A	1001	EPC	P1-O5	7.33	1.80	1.54

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	EPC	P2-O7	7.42	1.63	1.49
2	A	1001	EPC	P1-O3	9.01	1.79	1.50
2	B	1002	EPC	P1-O3	9.58	1.81	1.50
2	B	1002	EPC	C3-C2	10.41	1.48	1.40

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1002	EPC	O7-P2-C	-2.12	104.62	114.08
2	B	1002	EPC	C6-N1-C2	2.01	123.29	119.26
2	B	1002	EPC	C2A-C2-C3	3.09	124.03	120.90
2	A	1001	EPC	C-N2-C4A	3.41	125.47	118.00
2	B	1002	EPC	C-N2-C4A	3.52	125.70	118.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	EPC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	379/387 (97%)	-0.49	2 (0%) 91 91	7, 20, 40, 54	0
1	B	379/387 (97%)	-0.46	3 (0%) 87 87	10, 20, 43, 69	0
All	All	758/774 (97%)	-0.48	5 (0%) 89 88	7, 20, 41, 69	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	213	ARG	3.0
1	B	2	ASN	2.9
1	A	2	ASN	2.8
1	A	30	ASP	2.2
1	B	174	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	KCX	B	129	12/13	0.92	0.22	-	8,20,34,36	0
1	KCX	A	129	12/13	0.94	0.19	-	10,16,19,20	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EPC	B	1002	22/22	0.97	0.17	1.65	7,16,21,21	0
2	EPC	A	1001	22/22	0.98	0.15	0.66	7,12,18,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.