wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 08:31 PM BST

PDB ID	:	2FV4
Title	:	NMR solution structure of the yeast kinetochore Spc24/Spc25 globular domain
Authors	:	Schnell, J.R.; Chou, J.J.
Deposited on	:	2006-01-29

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	98	31% 37% 8% 15% 9%
2	B	77	39% 26% • 9% 25%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:134-A:152, A:158-A:173, A:183-A:221, B:156-B:184, B:188-B:194, B:199-B:213 (125)	0.45	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 6 single-model clusters were found.

Cluster number	Models
1	1, 2, 6, 8, 10, 13, 15
2	7, 9, 12, 16, 19
3	5, 20
Single-model clusters	3; 4; 11; 14; 17; 18

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2347 atoms, of which 1176 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Hypothetical 25.2 kDa protein in AFG3-SEB2 intergenic region.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	89	1363	432	677	123	128	3	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	GLY	-	CLONING ARTIFACT	UNP P40014
A	125	SER	-	CLONING ARTIFACT	UNP P40014
A	126	HIS	-	CLONING ARTIFACT	UNP P40014
A	127	MET	-	CLONING ARTIFACT	UNP P40014

- Molecule 2 is a protein called Hypothetical 24.6 kDa protein in ILV2-ADE17 intergenic region.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
2	B	58	984	310	499	83	92	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	137	MET	-	CLONING ARTIFACT	UNP Q04477

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Hypothetical 25.2 kDa protein in AFG3-SEB2 intergenic region

Chain A: 31% 37% 8% 15% 9%



• Molecule 2: Hypothetical 24.6 kDa protein in ILV2-ADE17 intergenic region

Chain B: 39% 26% • 9% 25%



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

• Molecule 1: Hypothetical 25.2 kDa protein in AFG3-SEB2 intergenic region

Chain A: 32% 31% 11% • 15% 9%

• Molecule 2: Hypothetical 24.6 kDa protein in ILV2-ADE17 intergenic region

Chain B: 34% 27% • • 9% 25%


MET
ALA
LYS
VAL
ILE
GLU
PRO
GLU
LEU
GLU
GLU
GLN
SER
ALA
VAL
THR
PRO
GLU
ALA
N156
E157
N158
I159
L160
V170
I171
I172
D173
L174
E175
N176
D177
Q178
I181
N182
R183
K184
N185
D186
G187
N188
I189
L192
D195
N196
N197
L198
F201
Y202
K203
T204
W208
E209
R210
L211
G212
K213

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *Simulated annealing.*.

Of the 30 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Xplor-NIH	refinement	2.11

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.04±0.00	0±0/588 (0.0±0.0%)	0.92±0.01	0±0/799 (0.0±0.1%)
2	B	0.98±0.00	0±0/439 (0.0±0.0%)	0.94±0.01	0±0/589 (0.0±0.0%)
All	All	1.01	0/20540 (0.0%)	0.93	6/27760 (0.0%)

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	141	TYR	CB-CG-CD2	-5.36	117.79	121.00	16	6

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	578	583	583	35±4
2	B	433	459	458	16±4
All	All	20220	20840	20820	947

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

5 of 362 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:217:LEU:HD23	1:A:218:LEU:N	0.84	1.87	9	4
1:A:206:LEU:HD23	1:A:206:LEU:O	0.81	1.76	17	3
1:A:188:LEU:HD13	1:A:188:LEU:N	0.75	1.97	11	1
1:A:134:ASP:O	1:A:138:VAL:HG23	0.74	1.83	18	16
2:B:191:ILE:H	2:B:191:ILE:HD13	0.74	1.41	16	2

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/98 (74%)	62±2 (85±2%)	6±2 (8±2%)	5±1 (7±2%)	3	19
2	B	49/77 (64%)	42±2 (86±3%)	5±1 (9±3%)	2±1 (4±1%)	6	30
All	All	2440/3500 (70%)	2087 (86%)	212 (9%)	141 (6%)	4	22

5 of 20 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	186	PRO	20
2	B	176	ASN	20
1	A	187	ALA	19
1	A	168	ARG	18
2	B	157	GLU	9

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	62/78 (79%)	51±2 (82±3%)	11±2 (18±3%)	5	39
2	B	49/71 (69%)	41±2 (83±4%)	8±2 (17±4%)	6	42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	2220/2980 (74%)	1828 (82%)	392 (18%)	6 41

5 of 71 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	209	PHE	20
2	B	171	ILE	20
2	B	181	ILE	20
1	A	189	ASP	19
1	A	200	LEU	18

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided