



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:42 PM GMT

PDB ID : 4FWE
Title : Native structure of LSD2 /AOF1/KDM1b in spacegroup of C2221 at 2.13Å
Authors : Zhang, Q.; Chen, Z.
Deposited on : 2012-07-01
Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1693 (2.16-2.12)
Clashscore	102246	1824 (2.16-2.12)
Ramachandran outliers	100387	1798 (2.16-2.12)
Sidechain outliers	100360	1798 (2.16-2.12)
RSRZ outliers	91569	1699 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	796	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6155 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific histone demethylase 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	750	Total	C	N	O	S	0	1	0
			5850	3741	996	1072	41			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	GLY	-	EXPRESSION TAG	UNP Q8NB78
A	28	HIS	-	EXPRESSION TAG	UNP Q8NB78
A	29	MET	-	EXPRESSION TAG	UNP Q8NB78

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is CITRATE ANION (three-letter code: FLC) (formula: C₆H₅O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	236	Total 236	O 236	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysine-specific histone demethylase 1B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	93.69 Å 96.27 Å 182.80 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.13 29.95 – 2.11	Depositor EDS
% Data completeness (in resolution range)	94.0 (50.00-2.13) 93.0 (29.95-2.11)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.12 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.203 , 0.246 0.203 , 0.201	Depositor DCC
R_{free} test set	2228 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	44.2	Xtriage
Anisotropy	0.423	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.4	EDS
Estimated twinning fraction	0.031 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 44372 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6155	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FLC, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	6/6000 (0.1%)	0.63	2/8145 (0.0%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	201	TRP	CD2-CE2	5.38	1.47	1.41
1	A	422	TRP	CD2-CE2	5.26	1.47	1.41
1	A	559	TRP	CD2-CE2	5.25	1.47	1.41
1	A	200	TRP	CD2-CE2	5.21	1.47	1.41
1	A	497	TRP	CD2-CE2	5.08	1.47	1.41
1	A	139	TRP	CD2-CE2	5.07	1.47	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	631	LEU	CA-CB-CG	-6.82	99.61	115.30
1	A	664	LEU	CA-CB-CG	5.83	128.70	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5850	0	5681	86	0
2	A	3	0	0	0	0
3	A	53	0	31	1	0
4	A	13	0	5	0	0
5	A	236	0	0	7	0
All	All	6155	0	5717	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:THR:HG21	1:A:92:CYS:HB2	1.70	0.74
1:A:473:ARG:HH11	1:A:473:ARG:HG3	1.54	0.73
1:A:502:THR:H	1:A:505:GLN:HE21	1.38	0.71
1:A:738:ARG:HD2	5:A:1175:HOH:O	1.92	0.70
1:A:260:VAL:HG12	1:A:261:HIS:H	1.59	0.68
1:A:375:LEU:HD22	1:A:403:ASN:HB3	1.77	0.66
1:A:158:GLN:HG3	5:A:1028:HOH:O	1.95	0.66
1:A:698:TYR:HB2	1:A:710[A]:MET:HG3	1.79	0.65
1:A:700:MET:CE	1:A:700:MET:HA	2.28	0.64
1:A:502:THR:H	1:A:505:GLN:NE2	1.96	0.62
1:A:172:LYS:CB	1:A:173:PRO:HD2	2.29	0.62
1:A:261:HIS:CE1	1:A:267:ARG:HG2	2.37	0.60
1:A:134:GLN:HG2	5:A:1224:HOH:O	2.02	0.59
1:A:470:GLU:OE1	1:A:683:HIS:HE1	1.85	0.59
1:A:544:GLU:HG2	1:A:551:LEU:HG	1.86	0.57
1:A:229:GLY:HA2	1:A:305:THR:HG23	1.86	0.57
1:A:700:MET:HA	1:A:700:MET:HE2	1.85	0.57
1:A:220:LEU:HD13	1:A:228:VAL:HG11	1.87	0.56
1:A:698:TYR:HB3	1:A:700:MET:HE1	1.88	0.56
1:A:700:MET:SD	1:A:710[A]:MET:HG2	2.47	0.55
1:A:285:ARG:NH1	1:A:290:GLU:OE2	2.41	0.54
1:A:454:LEU:HD21	1:A:585:LYS:HG2	1.89	0.54
1:A:413:ALA:O	1:A:595:LYS:HA	2.09	0.53
1:A:139:TRP:CE2	1:A:340:LEU:HD13	2.44	0.53
1:A:501:LYS:HA	1:A:505:GLN:NE2	2.24	0.52
1:A:698:TYR:HB3	1:A:700:MET:CE	2.39	0.52
1:A:316:ALA:O	1:A:320:THR:HG23	2.09	0.52
1:A:79:THR:CG2	1:A:92:CYS:HB2	2.39	0.51
1:A:782:ILE:HG22	1:A:797:THR:HG22	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:795:GLU:HG2	1:A:803:GLN:CA	2.40	0.51
1:A:543:LEU:HD21	1:A:559:TRP:CZ3	2.46	0.51
1:A:217:ALA:HB3	1:A:218:PRO:HD3	1.91	0.51
1:A:383:LYS:HA	1:A:621:GLN:OE1	2.10	0.51
1:A:74:ALA:HB2	1:A:79:THR:HG23	1.92	0.50
1:A:81:ARG:HD2	1:A:83:TYR:CZ	2.46	0.50
1:A:332:ILE:HD11	1:A:346:VAL:HG13	1.92	0.50
1:A:561:HIS:HE1	5:A:1143:HOH:O	1.94	0.49
1:A:205:LEU:HD11	1:A:341:VAL:HG23	1.95	0.49
1:A:274:GLN:HB3	1:A:275:PRO:HD2	1.94	0.49
1:A:666:PHE:O	1:A:707:SER:OG	2.31	0.48
1:A:293:GLU:OE2	1:A:311:ARG:HD3	2.13	0.48
1:A:501:LYS:HD2	1:A:505:GLN:NE2	2.29	0.47
1:A:220:LEU:HD13	1:A:228:VAL:CG1	2.44	0.47
1:A:138:TYR:CG	1:A:159:LEU:HB2	2.50	0.47
1:A:800:HIS:HD2	5:A:1030:HOH:O	1.98	0.47
1:A:473:ARG:HG3	1:A:473:ARG:NH1	2.26	0.46
1:A:229:GLY:HA3	1:A:309:ALA:HB2	1.96	0.46
1:A:81:ARG:HD2	1:A:83:TYR:CE1	2.49	0.46
1:A:261:HIS:HB3	1:A:270:GLN:HG2	1.98	0.46
1:A:703:GLN:HA	1:A:703:GLN:HE21	1.79	0.46
1:A:280:LYS:HE3	1:A:283:CYS:SG	2.55	0.46
1:A:693:LEU:HD22	1:A:694:PHE:CE2	2.51	0.45
1:A:722:ARG:HH11	1:A:722:ARG:HB3	1.81	0.45
1:A:401:LEU:HB2	1:A:408:VAL:HG11	1.98	0.45
1:A:102:SER:HA	1:A:107:TYR:CG	2.51	0.45
1:A:374:TYR:CD1	1:A:405:GLY:HA2	2.52	0.45
1:A:205:LEU:HD11	1:A:341:VAL:CG2	2.47	0.45
1:A:140:VAL:HG21	1:A:153:LEU:HD11	1.99	0.45
1:A:388:ILE:CD1	1:A:601:ILE:HD11	2.47	0.44
1:A:498:ARG:HD2	1:A:557:ARG:HA	1.98	0.44
1:A:640:ASN:HA	1:A:641:PRO:C	2.38	0.44
1:A:126:SER:HB2	5:A:1185:HOH:O	2.18	0.44
1:A:79:THR:HG21	1:A:93:ASN:H	1.83	0.44
1:A:700:MET:SD	1:A:710[B]:MET:HE2	2.59	0.43
1:A:795:GLU:HG2	1:A:803:GLN:N	2.32	0.43
1:A:144:LYS:HA	1:A:145:PRO:HD3	1.93	0.43
1:A:685:PRO:HB3	1:A:691:ARG:HA	2.00	0.43
1:A:482:ARG:NH1	1:A:532:GLU:OE2	2.51	0.42
1:A:628:PRO:HG2	1:A:763:ILE:HA	2.01	0.42
1:A:66:PHE:HZ	1:A:131:MET:HB3	1.84	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:357:THR:HG21	1:A:375:LEU:HB3	2.02	0.42
1:A:626:THR:HG22	1:A:793:ALA:HB3	2.02	0.42
1:A:486:HIS:HB2	5:A:1205:HOH:O	2.19	0.42
1:A:576:THR:HB	1:A:577:PRO:HD3	2.02	0.42
1:A:647:LYS:NZ	1:A:781:ILE:O	2.53	0.41
1:A:228:VAL:CG1	1:A:228:VAL:O	2.67	0.41
1:A:595:LYS:C	1:A:597:PRO:HD3	2.39	0.41
1:A:229:GLY:HA2	1:A:305:THR:CG2	2.50	0.41
1:A:473:ARG:HH11	1:A:473:ARG:CG	2.30	0.41
1:A:803:GLN:O	3:A:904:FAD:H1'2	2.20	0.41
1:A:464:ARG:HH12	1:A:676:GLN:HB2	1.85	0.41
1:A:755:THR:C	1:A:756:ARG:HG2	2.41	0.41
1:A:114:LYS:HG3	1:A:125:PRO:HB2	2.03	0.41
1:A:502:THR:N	1:A:505:GLN:HE21	2.10	0.41
1:A:555:SER:O	1:A:559:TRP:HB3	2.20	0.40
1:A:456:ILE:HB	1:A:577:PRO:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	745/796 (94%)	707 (95%)	36 (5%)	2 (0%)	46 41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	180	GLU
1	A	173	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	619/678 (91%)	577 (93%)	42 (7%)	20	13

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LYS
1	A	75	LYS
1	A	79	THR
1	A	101	ARG
1	A	148	ARG
1	A	158	GLN
1	A	159	LEU
1	A	174	ASN
1	A	192	ARG
1	A	194	LEU
1	A	204	MET
1	A	290	GLU
1	A	294	LEU
1	A	330	LYS
1	A	332	ILE
1	A	340	LEU
1	A	344	ARG
1	A	361	LEU
1	A	367	LEU
1	A	369	VAL
1	A	375	LEU
1	A	376	LEU
1	A	431	THR
1	A	432	VAL
1	A	536	LEU
1	A	540	LEU
1	A	543	LEU
1	A	576	THR
1	A	583	ILE
1	A	616	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	631	LEU
1	A	680	PHE
1	A	683	HIS
1	A	689	SER
1	A	693	LEU
1	A	703	GLN
1	A	710[A]	MET
1	A	710[B]	MET
1	A	732	GLN
1	A	756	ARG
1	A	789	THR
1	A	795	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
1	A	135	GLN
1	A	141	GLN
1	A	198	ASN
1	A	334	HIS
1	A	459	HIS
1	A	488	ASN
1	A	503	GLN
1	A	505	GLN
1	A	550	ASN
1	A	552	HIS
1	A	640	ASN
1	A	683	HIS
1	A	703	GLN
1	A	800	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FAD	A	904	-	48,58,58	1.17	3 (6%)	54,89,89	2.03	10 (18%)
4	FLC	A	905	-	3,12,12	0.79	0	3,17,17	2.08	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	904	-	-	0/30/50/50	0/6/6/6
4	FLC	A	905	-	-	0/6/16/16	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	904	FAD	C5X-N5	2.66	1.39	1.35
3	A	904	FAD	C4X-N5	3.99	1.39	1.33
3	A	904	FAD	C4-N3	4.18	1.40	1.33

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	904	FAD	N3A-C2A-N1A	-9.88	121.33	128.89
3	A	904	FAD	C4B-O4B-C1B	-4.74	104.50	109.72
3	A	904	FAD	P-O3P-PA	-3.57	122.71	132.73
3	A	904	FAD	C1B-N9A-C4A	-2.63	122.98	126.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	904	FAD	C4A-C5A-N7A	-2.51	107.17	109.48
3	A	904	FAD	C4X-C4-N3	-2.34	120.39	123.59
3	A	904	FAD	C4X-N5-C5X	2.25	119.36	116.76
3	A	904	FAD	C1'-N10-C9A	2.31	121.46	118.86
3	A	904	FAD	C4-C4X-N5	2.89	122.22	118.72
4	A	905	FLC	CG-CB-CA	3.41	117.96	109.81
3	A	904	FAD	C4-N3-C2	4.49	119.13	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	904	FAD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	750/796 (94%)	0.25	44 (5%) 26 34	32, 48, 70, 114	1 (0%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	179	PRO	5.9
1	A	822	PHE	5.3
1	A	504	LEU	4.8
1	A	181	THR	4.8
1	A	180	GLU	4.6
1	A	370	GLY	4.4
1	A	50	TYR	4.3
1	A	56	ALA	4.3
1	A	200	TRP	4.2
1	A	374	TYR	4.1
1	A	182	SER	4.0
1	A	373	GLN	3.8
1	A	522	LYS	3.7
1	A	173	PRO	3.7
1	A	371	ALA	3.7
1	A	265	MET	3.5
1	A	382	ASN	3.4
1	A	183	ASP	3.2
1	A	168	ARG	3.0
1	A	260	VAL	3.0
1	A	341	VAL	2.9
1	A	606	ASP	2.9
1	A	810	LEU	2.9
1	A	813	VAL	2.8
1	A	368	SER	2.6
1	A	805	VAL	2.6
1	A	174	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	793	ALA	2.4
1	A	428	LYS	2.4
1	A	276	ASN	2.4
1	A	806	THR	2.3
1	A	101	ARG	2.3
1	A	372	ASP	2.2
1	A	145	PRO	2.2
1	A	346	VAL	2.1
1	A	108	ASP	2.1
1	A	224	TYR	2.1
1	A	808	ALA	2.1
1	A	809	TYR	2.0
1	A	103	HIS	2.0
1	A	737	LEU	2.0
1	A	380	TYR	2.0
1	A	278	CYS	2.0
1	A	605	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	FLC	A	905	13/13	0.91	0.15	0.19	55,60,64,65	0
3	FAD	A	904	53/53	0.97	0.13	-0.22	29,34,38,39	0
2	ZN	A	902	1/1	0.99	0.06	-1.21	49,49,49,49	0
2	ZN	A	901	1/1	0.99	0.07	-1.72	55,55,55,55	0
2	ZN	A	903	1/1	0.99	0.04	-1.93	48,48,48,48	0

6.5 Other polymers

There are no such residues in this entry.