



wwPDB EM Map/Model Validation Report ⓘ

Jun 9, 2016 – 03:29 PM EDT

PDB ID : 5FXG
EMDB ID: : 3352
Title : GLUN1B-GLUN2B NMDA RECEPTOR IN ACTIVE CONFORMATION
Authors : Tajima, N.; Karakas, E.; Grant, T.; Simorowski, N.; Diaz-Avalos, R.; Grigorieff, N.; Furukawa, H.
Deposited on : 2016-03-02
Resolution : 6.80 Å(reported)

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/EMValidationReportHelp>

MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 6.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
Ramachandran outliers	111179	726

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	846	53% 24% 23%
1	C	846	52% 24% 23%
2	B	827	52% 24% 23%
2	D	827	52% 24% 23%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-METHYL-D-ASPARTATE RECEPTOR GLUN1.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	A	648	Total	C	N	O	0	0
			3199	1903	648	648		
1	C	648	Total	C	N	O	0	0
			3199	1903	648	648		

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	GLN	ASN	ENGINEERED MUTATION	UNP P35439
A	260	ASP	ASN	ENGINEERED MUTATION	UNP P35439
A	371	GLN	ASN	ENGINEERED MUTATION	UNP P35439
A	492	GLN	ASN	ENGINEERED MUTATION	UNP P35439
A	512	GLN	ASN	ENGINEERED MUTATION	UNP P35439
A	615	GLN	GLU	ENGINEERED MUTATION	UNP P35439
A	616	SER	GLU	ENGINEERED MUTATION	UNP P35439
A	618	SER	GLU	ENGINEERED MUTATION	UNP P35439
A	619	THR	GLU	ENGINEERED MUTATION	UNP P35439
A	792	GLN	ASN	ENGINEERED MUTATION	UNP P35439
A	831	CYS	PHE	ENGINEERED MUTATION	UNP P35439
A	865	ASN	ARG	ENGINEERED MUTATION	UNP P35439
A	866	GLY	ARG	ENGINEERED MUTATION	UNP P35439
A	867	ALA	LYS	ENGINEERED MUTATION	UNP P35439
C	61	GLN	ASN	ENGINEERED MUTATION	UNP P35439
C	260	ASP	ASN	ENGINEERED MUTATION	UNP P35439
C	371	GLN	ASN	ENGINEERED MUTATION	UNP P35439
C	492	GLN	ASN	ENGINEERED MUTATION	UNP P35439
C	512	GLN	ASN	ENGINEERED MUTATION	UNP P35439
C	615	GLN	GLU	ENGINEERED MUTATION	UNP P35439
C	616	SER	GLU	ENGINEERED MUTATION	UNP P35439
C	618	SER	GLU	ENGINEERED MUTATION	UNP P35439
C	619	THR	GLU	ENGINEERED MUTATION	UNP P35439
C	792	GLN	ASN	ENGINEERED MUTATION	UNP P35439
C	831	CYS	PHE	ENGINEERED MUTATION	UNP P35439
C	865	ASN	ARG	ENGINEERED MUTATION	UNP P35439

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	866	GLY	ARG	ENGINEERED MUTATION	UNP P35439
C	867	ALA	LYS	ENGINEERED MUTATION	UNP P35439

- Molecule 2 is a protein called N-METHYL-D-ASPARTATE RECEPTOR GLUN2B.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	B	634	Total	C	N	O	0	0
			3125	1857	634	634		
2	D	634	Total	C	N	O	0	0
			3125	1857	634	634		

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	26	GLY	-	EXPRESSION TAG	UNP Q00960
B	348	ASP	ASN	ENGINEERED MUTATION	UNP Q00960
B	557	CYS	ASP	ENGINEERED MUTATION	UNP Q00960
B	588	SER	CYS	ENGINEERED MUTATION	UNP Q00960
B	838	SER	CYS	ENGINEERED MUTATION	UNP Q00960
B	849	SER	CYS	ENGINEERED MUTATION	UNP Q00960
D	26	GLY	-	EXPRESSION TAG	UNP Q00960
D	348	ASP	ASN	ENGINEERED MUTATION	UNP Q00960
D	557	CYS	ASP	ENGINEERED MUTATION	UNP Q00960
D	588	SER	CYS	ENGINEERED MUTATION	UNP Q00960
D	838	SER	CYS	ENGINEERED MUTATION	UNP Q00960
D	849	SER	CYS	ENGINEERED MUTATION	UNP Q00960



Category	Percentage
Used	52%
Not used	24%
Don't know	23%



● Molecule 2: N-METHYL-D-ASPARTATE RECEPTOR GLUN2B



4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of particles used	Not provided	Depositor
Resolution determination method	Not provided	Depositor
CTF correction method	Not provided	Depositor
Microscope	OTHER	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	100	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	22500	Depositor
Image detector	GATAN K2 SUMMIT (4K X 4K)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 2$	RMSZ	$\# Z > 2$
1	A	0.27	0/3194	0.45	0/4439
1	C	0.28	0/3194	0.45	0/4439
2	B	0.27	0/3121	0.45	0/4336
2	D	0.27	0/3121	0.45	0/4336
All	All	0.27	0/12630	0.45	0/17550

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3199	0	1427	124	0
1	C	3199	0	1427	127	0
2	B	3125	0	1379	123	0
2	D	3125	0	1379	121	0
All	All	12648	0	5612	493	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 493 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:736:LEU:O	2:B:740:ALA:HB2	1.55	1.06
2:D:736:LEU:O	2:D:740:ALA:HB2	1.55	1.04
1:A:388:TYR:HA	1:A:393:VAL:HA	1.57	0.87
1:C:388:TYR:HA	1:C:393:VAL:HA	1.57	0.86
2:B:343:THR:HA	2:B:349:LEU:H	1.42	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	638/846 (75%)	563 (88%)	75 (12%)	0	100	100
1	C	638/846 (75%)	563 (88%)	75 (12%)	0	100	100
2	B	626/827 (76%)	554 (88%)	69 (11%)	3 (0%)	34	77
2	D	626/827 (76%)	554 (88%)	69 (11%)	3 (0%)	34	77
All	All	2528/3346 (76%)	2234 (88%)	288 (11%)	6 (0%)	56	86

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	660	VAL
2	D	660	VAL
2	B	523	VAL
2	D	523	VAL
2	B	468	ILE

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.