



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:49 PM GMT

PDB ID : 4GBY
Title : The structure of the MFS (major facilitator superfamily) proton:xylose symporter XylE bound to D-xylose
Authors : Sun, L.F.; Zeng, X.; Yan, C.Y.; Yan, N.
Deposited on : 2012-07-28
Resolution : 2.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	491	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3687 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-xylose-proton symporter.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	475	3587	2370	568	625	24	0	0	0

- Molecule 2 is SUGAR (BETA-D-XYLOPYRANOSE) (three-letter code: XYP) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	10	5	5	0	0

- Molecule 3 is SUGAR (B-NONYLGLUCOSIDE) (three-letter code: BNG) (formula: C₁₅H₃₀O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			21	15	6		
3	A	1	Total	C	O	0	0
			21	15	6		
3	A	1	Total	C	O	0	0
			21	15	6		
3	A	1	Total	C	O	0	0
			21	15	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	O	0	0
			6	6		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-xylose-proton symporter



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	95.29 Å 95.29 Å 168.18 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.71 – 2.81 29.71 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.71-2.81) 99.7 (29.71-2.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.04 (at 2.80 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.228 , 0.274 0.231 , 0.279	Depositor DCC
R_{free} test set	1004 reflections (5.42%)	DCC
Wilson B-factor (Å ²)	77.8	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 70.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 19517 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3687	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, BNG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/3680	0.60	1/5019 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	30	VAL	CB-CA-C	-5.60	100.76	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3587	0	3591	94	0
2	A	10	0	10	2	0
3	A	84	0	120	9	0
4	A	6	0	0	0	0
All	All	3687	0	3721	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:307:LEU:HD13	3:A:502:BNG:O2	1.65	0.96
1:A:305:LYS:O	1:A:309:ALA:HA	1.88	0.73
1:A:470:THR:HG22	1:A:473:GLU:HB2	1.72	0.72
1:A:168:GLN:NE2	2:A:501:XYP:O4A	2.28	0.67
1:A:31:ILE:HD12	1:A:133:ARG:HD2	1.76	0.66
1:A:168:GLN:HG3	1:A:389:PRO:CA	2.26	0.66
1:A:168:GLN:HG3	1:A:389:PRO:HA	1.79	0.64
1:A:251:GLN:O	1:A:254:GLN:N	2.30	0.63
1:A:225:ARG:HG2	1:A:401:ASN:HD22	1.64	0.62
1:A:221:PRO:HB2	1:A:239:ILE:HD13	1.81	0.61
1:A:225:ARG:HG2	1:A:401:ASN:ND2	2.16	0.60
1:A:243:ILE:HG22	1:A:244:MET:HG2	1.84	0.60
1:A:145:MET:HG3	1:A:408:LEU:HD13	1.83	0.59
1:A:85:ARG:HD3	1:A:89:LYS:HE3	1.85	0.59
1:A:249:ALA:O	1:A:253:VAL:HG23	2.02	0.58
1:A:19:LEU:HB2	1:A:213:PHE:CD2	2.40	0.56
1:A:316:LEU:O	1:A:320:ILE:HG12	2.05	0.56
1:A:306:THR:HG22	1:A:307:LEU:N	2.20	0.56
1:A:332:ALA:O	1:A:336:VAL:HG22	2.05	0.56
1:A:393:VAL:O	1:A:396:SER:HB2	2.06	0.56
1:A:122:LEU:HD11	1:A:200:ARG:NH2	2.21	0.55
1:A:300:ALA:HB2	3:A:502:BNG:H7'2	1.88	0.55
1:A:421:PHE:HA	1:A:425:THR:HG23	1.88	0.55
1:A:247:THR:OG1	1:A:248:LEU:N	2.40	0.54
1:A:372:ALA:HB2	3:A:502:BNG:O6	2.08	0.54
1:A:26:TYR:CZ	1:A:178:VAL:HG13	2.43	0.54
1:A:284:LEU:HD13	1:A:458:PHE:HE1	1.75	0.52
1:A:168:GLN:HG3	1:A:389:PRO:N	2.24	0.52
1:A:31:ILE:HD13	1:A:61:VAL:HA	1.92	0.52
1:A:30:VAL:HG13	1:A:178:VAL:HG11	1.92	0.51
1:A:68:CYS:HB3	1:A:416:TRP:CD2	2.46	0.51
1:A:164:VAL:HB	1:A:393:VAL:HG22	1.93	0.51
1:A:26:TYR:O	1:A:30:VAL:HG22	2.12	0.50
1:A:309:ALA:O	1:A:310:SER:O	2.30	0.50
1:A:159:ILE:HD12	1:A:163:LEU:HD11	1.92	0.50
1:A:385:MET:HE2	1:A:386:SER:HB2	1.94	0.49
1:A:441:ASN:HB3	1:A:443:PHE:H	1.77	0.49
1:A:41:VAL:HG13	1:A:122:LEU:HB2	1.93	0.49
1:A:307:LEU:HD13	3:A:502:BNG:HO2	1.72	0.49
1:A:352:MET:HB2	1:A:379:TYR:CD2	2.48	0.49
1:A:251:GLN:O	1:A:252:ALA:C	2.52	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:GLY:HA3	1:A:132:TYR:CD1	2.49	0.48
1:A:88:LEU:HD21	1:A:146:LEU:HB2	1.96	0.48
1:A:13:ILE:HG21	1:A:154:LEU:HD22	1.95	0.47
1:A:113:ASN:OD1	1:A:119:PRO:HG3	2.15	0.47
1:A:341:ARG:N	1:A:465:GLU:OE2	2.40	0.46
3:A:504:BNG:H8'2	3:A:504:BNG:H5'1	1.57	0.46
1:A:240:LEU:O	1:A:244:MET:HG3	2.17	0.45
1:A:356:MET:HG2	1:A:379:TYR:CD2	2.51	0.45
1:A:101:GLY:HA3	1:A:132:TYR:CG	2.52	0.45
1:A:18:THR:HG21	1:A:166:PHE:HD1	1.82	0.45
1:A:314:ALA:O	1:A:315:LEU:C	2.55	0.45
1:A:352:MET:HE2	1:A:379:TYR:CE2	2.52	0.45
1:A:194:LEU:HD12	1:A:198:GLY:HA3	1.99	0.45
1:A:317:GLN:O	1:A:321:VAL:HG23	2.17	0.45
1:A:249:ALA:O	1:A:250:THR:C	2.55	0.44
1:A:228:MET:SD	1:A:233:GLN:HG2	2.57	0.44
1:A:463:VAL:HA	1:A:464:PRO:HD3	1.86	0.44
1:A:140:VAL:O	1:A:144:SER:HB3	2.16	0.44
1:A:311:THR:O	1:A:312:ASP:C	2.56	0.44
1:A:280:ILE:O	1:A:284:LEU:HB2	2.18	0.44
1:A:118:VAL:HA	1:A:119:PRO:HD3	1.75	0.44
1:A:24:PHE:CD1	1:A:24:PHE:C	2.92	0.44
1:A:441:ASN:ND2	3:A:503:BNG:O4	2.50	0.43
1:A:13:ILE:HG21	1:A:154:LEU:CD2	2.48	0.43
1:A:168:GLN:HE22	2:A:501:XYP:C1B	2.32	0.43
1:A:20:GLY:HA3	1:A:147:SER:OG	2.19	0.43
1:A:248:LEU:O	1:A:249:ALA:C	2.57	0.43
1:A:313:ILE:O	1:A:314:ALA:C	2.57	0.43
1:A:311:THR:O	1:A:314:ALA:HB3	2.19	0.43
1:A:230:ARG:HD3	1:A:230:ARG:HA	1.80	0.43
1:A:470:THR:CG2	1:A:473:GLU:HB2	2.46	0.43
1:A:152:ALA:HB2	1:A:396:SER:HB3	2.01	0.43
1:A:24:PHE:HD1	1:A:24:PHE:C	2.21	0.43
1:A:285:SER:OG	1:A:415:GLN:HB2	2.19	0.42
1:A:309:ALA:O	1:A:310:SER:C	2.57	0.42
1:A:332:ALA:HB2	1:A:385:MET:HA	2.02	0.42
1:A:324:ILE:HG13	1:A:377:LEU:HD22	2.00	0.42
1:A:465:GLU:HG3	1:A:467:LYS:HG2	2.01	0.42
1:A:107:LEU:HD22	3:A:504:BNG:H2'1	2.02	0.41
1:A:382:ALA:O	1:A:385:MET:HG3	2.21	0.41
1:A:225:ARG:NH2	1:A:472:GLU:OE1	2.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:MET:HG3	1:A:256:ILE:HG21	2.02	0.41
1:A:372:ALA:HB1	3:A:502:BNG:H3'1	2.02	0.41
1:A:470:THR:HG23	1:A:473:GLU:H	1.85	0.41
1:A:243:ILE:C	1:A:244:MET:HG2	2.41	0.41
1:A:243:ILE:HG22	1:A:243:ILE:O	2.19	0.41
1:A:299:TYR:O	1:A:302:GLU:N	2.53	0.41
1:A:193:TRP:HA	1:A:196:THR:HG22	2.03	0.41
1:A:377:LEU:HD23	1:A:377:LEU:HA	1.86	0.41
1:A:352:MET:CE	1:A:383:PHE:HB2	2.51	0.40
1:A:293:ILE:HG23	1:A:294:ASN:N	2.37	0.40
1:A:75:GLY:HA3	1:A:145:MET:SD	2.62	0.40
1:A:314:ALA:O	1:A:317:GLN:N	2.55	0.40
1:A:110:THR:O	3:A:504:BNG:O6	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	473/491 (96%)	417 (88%)	49 (10%)	7 (2%)	13	40

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	107	LEU
1	A	156	PRO
1	A	310	SER
1	A	314	ALA
1	A	7	SER
1	A	430	ASP
1	A	190	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	367/396 (93%)	329 (90%)	38 (10%)	9 25

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	PHE
1	A	30	VAL
1	A	49	SER
1	A	81	ARG
1	A	106	GLU
1	A	107	LEU
1	A	116	ASN
1	A	133	ARG
1	A	144	SER
1	A	154	LEU
1	A	177	LEU
1	A	178	VAL
1	A	196	THR
1	A	244	MET
1	A	251	GLN
1	A	261	ASP
1	A	272	MET
1	A	275	VAL
1	A	278	ILE
1	A	283	MET
1	A	285	SER
1	A	291	VAL
1	A	307	LEU
1	A	317	GLN
1	A	336	VAL
1	A	339	PHE
1	A	366	GLN
1	A	373	LEU
1	A	376	MET
1	A	385	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	401	ASN
1	A	403	ILE
1	A	408	LEU
1	A	417	LEU
1	A	425	THR
1	A	472	GLU
1	A	475	GLU
1	A	477	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	GLN
1	A	415	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XYP	A	501	-	10,10,10	0.60	0	12,14,14	1.57	3 (25%)
3	BNG	A	502	-	21,21,21	0.42	0	26,26,26	0.91	1 (3%)
3	BNG	A	503	-	21,21,21	0.45	0	26,26,26	1.05	1 (3%)
3	BNG	A	504	-	21,21,21	0.45	0	26,26,26	0.78	0
3	BNG	A	505	-	21,21,21	0.41	0	26,26,26	0.92	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	A	501	-	-	0/0/17/17	0/1/1/1
3	BNG	A	502	-	-	0/12/32/32	0/1/1/1
3	BNG	A	503	-	-	0/12/32/32	0/1/1/1
3	BNG	A	504	-	-	0/12/32/32	0/1/1/1
3	BNG	A	505	-	-	0/12/32/32	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	XYP	O4B-C4B-C5B	-2.45	104.29	109.21
3	A	505	BNG	O1-C1-C2	-2.42	104.99	108.04
3	A	503	BNG	C1-O5-C5	-2.39	109.11	113.75
3	A	502	BNG	C1'-O1-C1	-2.38	109.78	113.94
2	A	501	XYP	O3B-C3B-C4B	2.40	114.34	110.00
2	A	501	XYP	C5B-C4B-C3B	2.58	112.59	109.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	XYP	2	0
3	A	502	BNG	5	0
3	A	503	BNG	1	0
3	A	504	BNG	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	475/491 (96%)	0.15	48 (10%) 9 4	40, 69, 147, 221	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	TYR	5.0
1	A	251	GLN	4.5
1	A	115	ASP	4.3
1	A	7	SER	4.2
1	A	310	SER	4.1
1	A	274	GLY	4.0
1	A	245	GLY	4.0
1	A	183	TYR	3.9
1	A	311	THR	3.8
1	A	272	MET	3.7
1	A	308	GLY	3.6
1	A	307	LEU	3.4
1	A	250	THR	3.4
1	A	313	ILE	3.3
1	A	192	SER	3.3
1	A	315	LEU	3.2
1	A	167	ASN	3.2
1	A	392	TRP	3.2
1	A	429	MET	3.2
1	A	309	ALA	3.1
1	A	191	ALA	3.1
1	A	186	ALA	3.0
1	A	6	ASN	3.0
1	A	28	THR	2.9
1	A	460	TRP	2.9
1	A	187	ARG	2.9
1	A	275	VAL	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	184	PHE	2.8
1	A	298	TYR	2.7
1	A	334	MET	2.7
1	A	188	SER	2.6
1	A	244	MET	2.6
1	A	338	LYS	2.5
1	A	11	PHE	2.5
1	A	304	PHE	2.5
1	A	396	SER	2.5
1	A	271	LEU	2.4
1	A	386	SER	2.4
1	A	218	TYR	2.3
1	A	8	SER	2.3
1	A	50	GLU	2.3
1	A	312	ASP	2.3
1	A	165	SER	2.1
1	A	164	VAL	2.1
1	A	395	LEU	2.1
1	A	148	PRO	2.1
1	A	152	ALA	2.1
1	A	389	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	BNG	A	504	21/21	0.87	0.34	1.67	117,123,127,128	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	BNG	A	503	21/21	0.78	0.33	1.48	137,140,142,144	0
3	BNG	A	505	21/21	0.92	0.27	1.42	105,110,116,117	0
3	BNG	A	502	21/21	0.94	0.32	1.18	83,94,96,97	0
2	XYP	A	501	10/10	0.96	0.34	1.07	47,52,56,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.