



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 20, 2016 – 06:58 PM EST

PDB ID : 4GDD  
Title : Crystal structure of oxidized Aspergillus fumigatus UDP-galactopyranose mutase complexed with NADH  
Authors : Tanner, J.J.; Dhatwalia, R.D.; Singh, H.  
Deposited on : 2012-07-31  
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	513	
1	B	513	
1	C	513	
1	D	513	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	B	605	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15540 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-galactopyranose mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	494	Total	C	N	O	S	0	0	0
			3776	2409	634	712	21			
1	B	494	Total	C	N	O	S	0	0	0
			3777	2406	633	718	20			
1	C	499	Total	C	N	O	S	0	0	0
			3798	2422	641	716	19			
1	D	502	Total	C	N	O	S	0	0	0
			3813	2434	640	719	20			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	ALA	-	EXPRESSION TAG	UNP Q4W1X2
A	-1	ILE	-	EXPRESSION TAG	UNP Q4W1X2
A	0	ALA	-	EXPRESSION TAG	UNP Q4W1X2
A	344	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
A	345	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
A	429	THR	ALA	SEE REMARK 999	UNP Q4W1X2
B	-2	ALA	-	EXPRESSION TAG	UNP Q4W1X2
B	-1	ILE	-	EXPRESSION TAG	UNP Q4W1X2
B	0	ALA	-	EXPRESSION TAG	UNP Q4W1X2
B	344	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
B	345	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
B	429	THR	ALA	SEE REMARK 999	UNP Q4W1X2
C	-2	ALA	-	EXPRESSION TAG	UNP Q4W1X2
C	-1	ILE	-	EXPRESSION TAG	UNP Q4W1X2
C	0	ALA	-	EXPRESSION TAG	UNP Q4W1X2
C	344	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
C	345	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
C	429	THR	ALA	SEE REMARK 999	UNP Q4W1X2
D	-2	ALA	-	EXPRESSION TAG	UNP Q4W1X2
D	-1	ILE	-	EXPRESSION TAG	UNP Q4W1X2
D	0	ALA	-	EXPRESSION TAG	UNP Q4W1X2

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
D	344	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
D	345	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
D	429	THR	ALA	SEE REMARK 999	UNP Q4W1X2

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula:  $C_{21}H_{29}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	O	0	0
			4	4		
5	B	7	Total	O	0	0
			7	7		
5	C	7	Total	O	0	0
			7	7		
5	D	8	Total	O	0	0
			8	8		







● Molecule 1: UDP-galactopyranose mutase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	218.25Å 218.25Å 318.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	162.48 – 2.75 162.48 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.4 (162.48-2.75) 99.3 (162.48-2.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.16	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 2.73Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.2_432)	Depositor
R, $R_{free}$	0.212 , 0.245 0.191 , 0.227	Depositor DCC
$R_{free}$ test set	5786 reflections (5.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	51.2	Xtriage
Anisotropy	0.318	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	15540	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAI, SO4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/3869	0.60	0/5276
1	B	0.45	0/3870	0.61	1/5279 (0.0%)
1	C	0.41	0/3894	0.56	0/5314
1	D	0.42	0/3909	0.57	0/5333
All	All	0.43	0/15542	0.59	1/21202 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	372	LEU	CA-CB-CG	6.95	131.28	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3776	0	3587	53	0
1	B	3777	0	3583	64	0
1	C	3798	0	3583	46	0
1	D	3813	0	3602	52	0
2	A	53	0	30	3	0
2	B	53	0	30	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	53	0	31	5	0
2	D	53	0	31	3	0
3	A	44	0	27	2	0
3	B	44	0	27	2	0
4	A	10	0	0	0	0
4	B	15	0	0	2	0
4	C	10	0	0	1	0
4	D	15	0	0	0	0
5	A	4	0	0	0	0
5	B	7	0	0	0	0
5	C	7	0	0	0	0
5	D	8	0	0	0	0
All	All	15540	0	14531	214	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 214 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:453:TYR:CE1	3:B:602:NAI:H42N	2.04	0.92
1:B:325:PHE:CD1	1:B:372:LEU:HD12	2.07	0.90
1:D:38:ASP:OD1	2:D:601:FAD:H1B	1.83	0.78
1:D:359:SER:OG	1:D:361:GLU:HG2	1.85	0.77
1:A:453:TYR:CE1	3:A:602:NAI:H42N	2.22	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	488/513 (95%)	469 (96%)	19 (4%)	0	100	100
1	B	488/513 (95%)	468 (96%)	20 (4%)	0	100	100
1	C	495/513 (96%)	478 (97%)	17 (3%)	0	100	100
1	D	498/513 (97%)	482 (97%)	16 (3%)	0	100	100
All	All	1969/2052 (96%)	1897 (96%)	72 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	386/432 (89%)	376 (97%)	10 (3%)	54	84
1	B	389/432 (90%)	378 (97%)	11 (3%)	51	82
1	C	383/432 (89%)	374 (98%)	9 (2%)	58	87
1	D	384/432 (89%)	374 (97%)	10 (3%)	54	84
All	All	1542/1728 (89%)	1502 (97%)	40 (3%)	54	84

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	430	LEU
1	C	27	ASN
1	D	430	LEU
1	B	485	ASP
1	C	175	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	GLN
1	A	458	GLN
1	B	98	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	458	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	601	-	52,58,58	2.01	19 (36%)	52,89,89	2.76	21 (40%)
3	NAI	A	602	-	41,48,48	1.76	15 (36%)	46,73,73	2.36	7 (15%)
4	SO4	A	603	-	4,4,4	0.11	0	6,6,6	0.12	0
4	SO4	A	604	-	4,4,4	0.23	0	6,6,6	0.24	0
2	FAD	B	601	-	52,58,58	2.06	19 (36%)	52,89,89	2.62	20 (38%)
3	NAI	B	602	-	41,48,48	1.82	14 (34%)	46,73,73	2.27	9 (19%)
4	SO4	B	603	-	4,4,4	0.17	0	6,6,6	0.29	0
4	SO4	B	604	-	4,4,4	0.18	0	6,6,6	0.27	0
4	SO4	B	605	-	4,4,4	0.29	0	6,6,6	0.17	0
2	FAD	C	601	-	52,58,58	1.92	17 (32%)	52,89,89	2.48	15 (28%)
4	SO4	C	602	-	4,4,4	0.14	0	6,6,6	0.23	0
4	SO4	C	603	-	4,4,4	0.26	0	6,6,6	0.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	D	601	-	52,58,58	2.03	19 (36%)	52,89,89	2.60	18 (34%)
4	SO4	D	602	-	4,4,4	0.32	0	6,6,6	0.31	0
4	SO4	D	603	-	4,4,4	0.17	0	6,6,6	0.23	0
4	SO4	D	604	-	4,4,4	0.25	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	0/30/50/50	0/6/6/6
3	NAI	A	602	-	-	0/25/72/72	0/5/5/5
4	SO4	A	603	-	-	0/0/0/0	0/0/0/0
4	SO4	A	604	-	-	0/0/0/0	0/0/0/0
2	FAD	B	601	-	-	0/30/50/50	0/6/6/6
3	NAI	B	602	-	-	0/25/72/72	0/5/5/5
4	SO4	B	603	-	-	0/0/0/0	0/0/0/0
4	SO4	B	604	-	-	0/0/0/0	0/0/0/0
4	SO4	B	605	-	-	0/0/0/0	0/0/0/0
2	FAD	C	601	-	-	0/30/50/50	0/6/6/6
4	SO4	C	602	-	-	0/0/0/0	0/0/0/0
4	SO4	C	603	-	-	0/0/0/0	0/0/0/0
2	FAD	D	601	-	-	0/30/50/50	0/6/6/6
4	SO4	D	602	-	-	0/0/0/0	0/0/0/0
4	SO4	D	603	-	-	0/0/0/0	0/0/0/0
4	SO4	D	604	-	-	0/0/0/0	0/0/0/0

The worst 5 of 103 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	FAD	C2B-C1B	-3.80	1.47	1.53
2	A	601	FAD	O2'-C2'	-3.42	1.35	1.43
3	B	602	NAI	C4N-C5N	-3.38	1.41	1.49
2	A	601	FAD	C2B-C1B	-3.16	1.48	1.53
3	A	602	NAI	C4N-C5N	-3.15	1.42	1.49

The worst 5 of 90 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	NAI	N3A-C2A-N1A	-12.02	119.43	128.87
3	B	602	NAI	N3A-C2A-N1A	-12.00	119.45	128.87

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	N3A-C2A-N1A	-10.75	120.43	128.87
2	B	601	FAD	N3A-C2A-N1A	-10.42	120.69	128.87
2	D	601	FAD	N3A-C2A-N1A	-10.22	120.84	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	FAD	3	0
3	A	602	NAI	2	0
2	B	601	FAD	6	0
3	B	602	NAI	2	0
4	B	605	SO4	2	0
2	C	601	FAD	5	0
4	C	603	SO4	1	0
2	D	601	FAD	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	494/513 (96%)	0.29	2 (0%) 93 92	25, 46, 76, 103	0
1	B	494/513 (96%)	0.26	3 (0%) 90 88	24, 47, 77, 121	0
1	C	499/513 (97%)	0.37	9 (1%) 71 66	24, 53, 84, 106	0
1	D	502/513 (97%)	0.38	11 (2%) 65 59	23, 54, 86, 104	0
All	All	1989/2052 (96%)	0.32	25 (1%) 79 75	23, 50, 81, 121	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	236	PHE	3.3
1	C	245	VAL	2.9
1	C	273	PHE	2.7
1	B	184	ALA	2.6
1	C	242	VAL	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	SO4	C	603	5/5	0.96	0.27	2.00	46,49,61,87	5
4	SO4	B	605	5/5	0.95	0.23	0.63	70,74,84,89	0
3	NAI	B	602	44/44	0.97	0.21	0.19	29,48,63,71	0
4	SO4	D	602	5/5	0.99	0.24	0.15	30,31,55,82	5
2	FAD	B	601	53/53	0.98	0.20	-0.04	18,43,62,66	0
2	FAD	D	601	53/53	0.95	0.21	-0.26	26,57,91,107	0
2	FAD	A	601	53/53	0.98	0.20	-0.37	25,44,63,69	0
2	FAD	C	601	53/53	0.97	0.20	-0.60	24,60,86,96	0
3	NAI	A	602	44/44	0.96	0.19	-1.10	22,54,71,78	0
4	SO4	A	603	5/5	0.98	0.17	-	29,37,66,74	5
4	SO4	C	602	5/5	0.94	0.16	-	63,74,90,102	0
4	SO4	D	604	5/5	0.95	0.22	-	51,102,114,116	0
4	SO4	A	604	5/5	0.93	0.23	-	53,68,89,90	0
4	SO4	B	604	5/5	0.97	0.21	-	45,66,73,76	5
4	SO4	B	603	5/5	0.93	0.12	-	68,94,101,101	0
4	SO4	D	603	5/5	0.94	0.21	-	80,89,94,94	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.