



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:54 AM GMT

PDB ID : 3GFB
Title : L-Threonine Dehydrogenase (TkTDH) from the hyperthermophilic archaeon
Thermococcus kodakaraensis
Authors : Bowyer, A.
Deposited on : 2009-02-26
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	<div> <div>3%</div> <div>64% 27% 7% ..</div> </div>
1	B	350	<div> <div>6%</div> <div>56% 36% 6% ..</div> </div>
1	C	350	<div> <div>8%</div> <div>57% 32% 9% ..</div> </div>
1	D	350	<div> <div>5%</div> <div>62% 29% 7% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAD	A	500	X	-	-	X
2	NAD	B	501	X	-	-	X
2	NAD	C	502	X	-	-	X
2	NAD	D	503	X	-	-	X
3	SO4	B	351	-	X	-	X
3	SO4	B	352	-	X	-	X
3	SO4	B	353	-	-	-	X
3	SO4	B	354	-	-	-	X
3	SO4	C	352	-	-	-	X
3	SO4	C	353	-	-	-	X
3	SO4	D	351	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11389 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-threonine 3-dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	347	Total	C	N	O	S	0	0	0
			2668	1726	444	486	12			
1	B	347	Total	C	N	O	S	0	0	0
			2668	1726	444	486	12			
1	C	347	Total	C	N	O	S	0	0	0
			2668	1726	444	486	12			
1	D	347	Total	C	N	O	S	0	0	0
			2668	1726	444	486	12			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	126	Total	O		0	0
			126	126			
4	B	119	Total	O		0	0
			119	119			
4	C	127	Total	O		0	0
			127	127			
4	D	119	Total	O		0	0
			119	119			


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-threonine 3-dehydrogenase




- Molecule 1: L-threonine 3-dehydrogenase




- Molecule 1: L-threonine 3-dehydrogenase





• Molecule 1: L-threonine 3-dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	124.48Å 124.48Å 271.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	73.92 – 2.40 73.83 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (73.92-2.40) 100.0 (73.83-2.40)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 2.40Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.217 , 0.277 0.217 , 0.275	Depositor DCC
R_{free} test set	4215 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 58.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	17 of 83945 reflections (0.020%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11389	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.44 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5213e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.52	15/2736 (0.5%)	1.25	15/3716 (0.4%)
1	B	1.54	23/2736 (0.8%)	1.29	13/3716 (0.3%)
1	C	1.57	24/2736 (0.9%)	1.24	13/3716 (0.3%)
1	D	1.58	17/2736 (0.6%)	1.32	17/3716 (0.5%)
All	All	1.55	79/10944 (0.7%)	1.28	58/14864 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
1	C	0	1
1	D	0	2
All	All	0	7

The worst 5 of 79 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	238	GLU	CB-CG	-11.32	1.30	1.52
1	B	257	VAL	CB-CG2	-10.54	1.30	1.52
1	B	238	GLU	CB-CG	-9.64	1.33	1.52
1	D	238	GLU	CG-CD	-9.55	1.37	1.51
1	B	238	GLU	CG-CD	-8.92	1.38	1.51

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	271	ARG	NE-CZ-NH2	-14.27	113.17	120.30
1	B	238	GLU	OE1-CD-OE2	11.38	136.95	123.30
1	D	179	LEU	CB-CG-CD2	-10.14	93.76	111.00
1	C	154	LEU	CA-CB-CG	9.21	136.49	115.30
1	C	230	ASP	CB-CG-OD2	-8.50	110.65	118.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	112	GLN	Peptide
1	A	113	ASN	Peptide
1	A	348	HIS	Peptide
1	B	348	HIS	Peptide
1	C	12	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2668	0	2683	89	0
1	B	2668	0	2683	121	0
1	C	2668	0	2684	116	0
1	D	2668	0	2683	101	0
2	A	44	0	26	6	0
2	B	44	0	26	11	0
2	C	44	0	26	4	0
2	D	44	0	26	13	0
3	B	25	0	0	2	0
3	C	15	0	0	1	0
3	D	10	0	0	0	0
4	A	126	0	0	7	0
4	B	119	0	0	14	0
4	C	127	0	0	6	0
4	D	119	0	0	14	0
All	All	11389	0	10837	414	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 414 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:311:LYS:CD	1:B:311:LYS:CG	1.76	1.60
1:C:259:PRO:CG	1:C:259:PRO:CB	1.75	1.43
1:B:178:PRO:CB	1:B:178:PRO:CG	1.75	1.42
1:A:110:VAL:HG11	1:B:285:ALA:CB	1.80	1.11
1:D:67:HIS:HB2	1:D:93:THR:HG21	1.22	1.11

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/350 (99%)	313 (91%)	21 (6%)	11 (3%)	5	4
1	B	345/350 (99%)	308 (89%)	31 (9%)	6 (2%)	11	14
1	C	345/350 (99%)	311 (90%)	26 (8%)	8 (2%)	8	8
1	D	345/350 (99%)	312 (90%)	27 (8%)	6 (2%)	11	14
All	All	1380/1400 (99%)	1244 (90%)	105 (8%)	31 (2%)	8	9

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	14	ALA
1	A	97	CYS
1	B	97	CYS
1	B	271	ARG
1	C	13	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/284 (100%)	262 (93%)	21 (7%)	17	26
1	B	283/284 (100%)	258 (91%)	25 (9%)	12	18
1	C	283/284 (100%)	258 (91%)	25 (9%)	12	18
1	D	283/284 (100%)	257 (91%)	26 (9%)	11	16
All	All	1132/1136 (100%)	1035 (91%)	97 (9%)	13	19

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	335	MET
1	C	134	LYS
1	D	206	LYS
1	B	349	LYS
1	C	41	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	320	HIS
1	C	139	ASN
1	D	308	GLN
1	B	348	HIS
1	C	94	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	A	500	-	38,48,48	1.52	5 (13%)	47,73,73	3.01	15 (31%)
3	SO4	B	351	-	4,4,4	3.72	3 (75%)	6,6,6	2.13	1 (16%)
3	SO4	B	352	-	4,4,4	3.02	3 (75%)	6,6,6	1.40	2 (33%)
3	SO4	B	353	-	4,4,4	5.34	3 (75%)	6,6,6	0.56	0
3	SO4	B	354	-	4,4,4	1.20	0	6,6,6	0.70	0
3	SO4	B	355	-	4,4,4	1.04	0	6,6,6	1.18	1 (16%)
2	NAD	B	501	-	38,48,48	3.07	6 (15%)	47,73,73	4.18	21 (44%)
3	SO4	C	351	-	4,4,4	0.30	0	6,6,6	0.65	0
3	SO4	C	352	-	4,4,4	0.26	0	6,6,6	0.13	0
3	SO4	C	353	-	4,4,4	0.64	0	6,6,6	1.05	0
2	NAD	C	502	-	38,48,48	2.43	6 (15%)	47,73,73	4.00	23 (48%)
3	SO4	D	351	-	4,4,4	0.13	0	6,6,6	0.58	0
3	SO4	D	352	-	4,4,4	0.88	0	6,6,6	0.96	0
2	NAD	D	503	-	38,48,48	2.26	8 (21%)	47,73,73	3.41	24 (51%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	500	-	2/2/11/11	0/22/62/62	0/5/5/5
3	SO4	B	351	-	-	0/0/0/0	0/0/0/0
3	SO4	B	352	-	-	0/0/0/0	0/0/0/0
3	SO4	B	353	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	B	354	-	-	0/0/0/0	0/0/0/0
3	SO4	B	355	-	-	0/0/0/0	0/0/0/0
2	NAD	B	501	-	2/2/11/11	0/22/62/62	0/5/5/5
3	SO4	C	351	-	-	0/0/0/0	0/0/0/0
3	SO4	C	352	-	-	0/0/0/0	0/0/0/0
3	SO4	C	353	-	-	0/0/0/0	0/0/0/0
2	NAD	C	502	-	2/2/11/11	0/22/62/62	0/5/5/5
3	SO4	D	351	-	-	0/0/0/0	0/0/0/0
3	SO4	D	352	-	-	0/0/0/0	0/0/0/0
2	NAD	D	503	-	2/2/11/11	0/22/62/62	0/5/5/5

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	502	NAD	O4D-C4D	-3.83	1.36	1.45
2	D	503	NAD	C3N-C7N	-3.11	1.45	1.50
2	B	501	NAD	O4B-C1B	-3.06	1.37	1.41
2	A	500	NAD	O4D-C4D	-2.94	1.38	1.45
2	C	502	NAD	PN-O5D	-2.87	1.45	1.59

The worst 5 of 87 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	502	NAD	N3A-C2A-N1A	-15.54	116.99	128.89
2	B	501	NAD	N3A-C2A-N1A	-12.79	119.10	128.89
2	C	502	NAD	C3N-C7N-N7N	-11.65	105.07	117.82
2	A	500	NAD	N3A-C2A-N1A	-10.91	120.54	128.89
2	D	503	NAD	C4D-O4D-C1D	-10.82	97.83	109.72

5 of 8 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	500	NAD	C4B
2	A	500	NAD	C4D
2	B	501	NAD	C4B
2	B	501	NAD	C4D
2	D	503	NAD	C4B

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 37 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	NAD	6	0
3	B	353	SO4	1	0
3	B	355	SO4	1	0
2	B	501	NAD	11	0
3	C	351	SO4	1	0
2	C	502	NAD	4	0
2	D	503	NAD	13	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/350 (99%)	0.15	12 (3%) 48 48	15, 33, 63, 88	1 (0%)
1	B	347/350 (99%)	0.27	22 (6%) 23 24	17, 35, 76, 100	1 (0%)
1	C	347/350 (99%)	0.33	29 (8%) 14 13	17, 36, 67, 88	1 (0%)
1	D	347/350 (99%)	0.18	19 (5%) 29 29	15, 33, 72, 97	1 (0%)
All	All	1388/1400 (99%)	0.23	82 (5%) 26 26	15, 34, 69, 100	4 (0%)

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	97	CYS	13.6
1	B	96	VAL	10.9
1	A	97	CYS	8.6
1	A	113	ASN	8.4
1	B	111	CYS	8.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SO4	B	351	5/5	0.56	0.47	21.92	116,117,118,118	0
3	SO4	B	352	5/5	0.76	0.31	13.57	105,106,108,109	0
3	SO4	D	351	5/5	0.67	0.48	11.82	127,128,129,131	0
3	SO4	C	352	5/5	0.84	0.34	7.43	101,101,102,102	0
2	NAD	D	503	44/44	0.92	0.27	5.98	8,23,27,29	44
3	SO4	C	353	5/5	0.78	0.32	5.69	87,88,90,90	0
2	NAD	A	500	44/44	0.86	0.33	5.42	9,28,32,33	44
3	SO4	B	353	5/5	0.50	0.35	4.99	114,115,117,117	0
2	NAD	C	502	44/44	0.89	0.32	4.88	6,26,31,35	44
2	NAD	B	501	44/44	0.82	0.31	4.74	6,26,34,37	44
3	SO4	B	354	5/5	0.78	0.23	3.55	95,96,98,99	0
3	SO4	B	355	5/5	0.96	0.15	-0.16	38,47,50,53	0
3	SO4	D	352	5/5	0.99	0.10	-1.67	30,44,46,49	0
3	SO4	C	351	5/5	0.80	0.41	-	112,112,112,113	0

6.5 Other polymers

There are no such residues in this entry.