



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:54 AM GMT

PDB ID : 3GHU  
Title : Human aldose reductase in complex with NADP<sup>+</sup> and the inhibitor IDD594. Investigation of global effects of radiation damage on protein structure. Forth stage of radiation damage.  
Authors : Petrova, T.; Ginell, S.; Hazemann, I.; Mitschler, A.; Podjarny, A.; Joachimiak, A.  
Deposited on : 2009-03-04  
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1495 (1.26-1.14)
Clashscore	102246	1607 (1.26-1.14)
Ramachandran outliers	100387	1540 (1.26-1.14)
Sidechain outliers	100360	1538 (1.26-1.14)
RSRZ outliers	91569	1500 (1.26-1.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div>2%</div> <div>82%</div> <div>16%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CIT	A	450[A]	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	316	3043	1958	514	555	16	5	105	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	4	ILE	LEU	SEE REMARK 999	UNP P15121

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is IDD594 (three-letter code: LDT) (formula:  $C_{16}H_{12}BrF_2NO_3S$ ).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	Br	C	F	N	O	S	0	0
			24	1	16	2	1	3	1		

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula:  $C_6H_8O_7$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			13	6	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	630	Total 671	O 671	0	295

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aldose reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.55Å 66.83Å 47.36Å 90.00° 92.22° 90.00°	Depositor
Resolution (Å)	50.00 – 1.20 23.21 – 1.20	Depositor EDS
% Data completeness (in resolution range)	84.9 (50.00-1.20) 96.3 (23.21-1.20)	Depositor EDS
$R_{merge}$	0.02	Depositor
$R_{sym}$	0.03	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.00 (at 1.20Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.095 , 0.137 0.104 , 0.106	Depositor DCC
$R_{free}$ test set	4618 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.3	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , 58.3	EDS
Estimated twinning fraction	0.006 for -l,k,h 0.028 for -h,-k,l 0.024 for l,-k,h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 92142 reflections	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3799	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, CIT, LDT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	2.73	24/3330 (0.7%)	1.76	42/4508 (0.9%)

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	26[A]	GLN	CD-OE1	98.69	3.41	1.24
1	A	26[B]	GLN	CD-OE1	98.69	3.41	1.24
1	A	26[A]	GLN	CD-NE2	33.02	2.15	1.32
1	A	26[B]	GLN	CD-NE2	33.02	2.15	1.32
1	A	315[A]	PHE	C-OXT	13.53	1.49	1.23
1	A	98[A]	ASP	CG-OD1	13.41	1.56	1.25
1	A	98[B]	ASP	CG-OD1	13.41	1.56	1.25
1	A	53	GLU	CD-OE1	11.92	1.38	1.25
1	A	139	ASP	CG-OD2	-10.64	1.00	1.25
1	A	98[A]	ASP	CG-OD2	9.23	1.46	1.25
1	A	98[B]	ASP	CG-OD2	9.23	1.46	1.25
1	A	216	ASP	CG-OD1	8.67	1.45	1.25
1	A	284[A]	ASP	CG-OD2	8.25	1.44	1.25
1	A	125[A]	ASP	CG-OD2	8.18	1.44	1.25
1	A	139	ASP	CG-OD1	7.96	1.43	1.25
1	A	51[A]	GLU	CD-OE1	6.88	1.33	1.25
1	A	149	ASP	CG-OD1	6.77	1.41	1.25
1	A	145	GLU	CD-OE1	6.65	1.32	1.25
1	A	94[A]	LYS	CB-CG	6.64	1.70	1.52
1	A	94[B]	LYS	CB-CG	6.64	1.70	1.52
1	A	224	ASP	CG-OD1	5.87	1.38	1.25
1	A	150	GLU	CD-OE2	-5.61	1.19	1.25
1	A	216	ASP	CG-OD2	5.60	1.38	1.25
1	A	145	GLU	CD-OE2	-5.05	1.20	1.25

All (42) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	26[A]	GLN	CG-CD-OE1	-36.43	48.74	121.60
1	A	26[B]	GLN	CG-CD-OE1	-36.43	48.74	121.60
1	A	26[A]	GLN	OE1-CD-NE2	-35.63	39.94	121.90
1	A	26[B]	GLN	OE1-CD-NE2	-35.63	39.94	121.90
1	A	98[A]	ASP	CB-CG-OD1	19.45	135.80	118.30
1	A	98[B]	ASP	CB-CG-OD1	19.45	135.80	118.30
1	A	167	GLU	OE1-CD-OE2	-18.85	100.68	123.30
1	A	26[A]	GLN	CG-CD-NE2	-14.68	81.47	116.70
1	A	26[B]	GLN	CG-CD-NE2	-14.68	81.47	116.70
1	A	98[A]	ASP	OD1-CG-OD2	-12.42	99.71	123.30
1	A	98[B]	ASP	OD1-CG-OD2	-12.42	99.71	123.30
1	A	149	ASP	CB-CG-OD2	12.31	129.38	118.30
1	A	139	ASP	CB-CG-OD2	12.21	129.29	118.30
1	A	167	GLU	CG-CD-OE2	10.09	138.49	118.30
1	A	139	ASP	CB-CG-OD1	-9.15	110.06	118.30
1	A	125[A]	ASP	CB-CG-OD2	-9.09	110.12	118.30
1	A	125[A]	ASP	CB-CG-OD1	8.74	126.16	118.30
1	A	224	ASP	CB-CG-OD2	7.97	125.47	118.30
1	A	115	PHE	CB-CG-CD1	7.58	126.10	120.80
1	A	284[A]	ASP	CB-CG-OD1	-7.54	111.51	118.30
1	A	293[A]	ARG	NE-CZ-NH1	-7.39	116.60	120.30
1	A	293[B]	ARG	NE-CZ-NH1	-7.39	116.60	120.30
1	A	293[C]	ARG	NE-CZ-NH1	-7.39	116.60	120.30
1	A	64[A]	GLU	OE1-CD-OE2	-7.17	114.70	123.30
1	A	64[B]	GLU	OE1-CD-OE2	-7.17	114.70	123.30
1	A	268	ARG	NE-CZ-NH2	-7.15	116.72	120.30
1	A	60[A]	GLU	OE1-CD-OE2	-6.90	115.02	123.30
1	A	98[A]	ASP	CB-CG-OD2	6.88	124.49	118.30
1	A	98[B]	ASP	CB-CG-OD2	6.88	124.49	118.30
1	A	60[A]	GLU	CG-CD-OE1	6.55	131.40	118.30
1	A	209	TYR	CA-CB-CG	6.50	125.75	113.40
1	A	284[A]	ASP	CB-CG-OD2	6.43	124.08	118.30
1	A	217[A]	ARG	CD-NE-CZ	6.41	132.57	123.60
1	A	217[B]	ARG	CD-NE-CZ	6.41	132.57	123.60
1	A	149	ASP	CB-CG-OD1	-6.37	112.57	118.30
1	A	189[A]	TYR	CB-CG-CD1	6.11	124.67	121.00
1	A	189[B]	TYR	CB-CG-CD1	6.11	124.67	121.00
1	A	284[A]	ASP	CA-CB-CG	5.84	126.24	113.40
1	A	53	GLU	CG-CD-OE1	5.36	129.01	118.30
1	A	189[A]	TYR	CB-CG-CD2	-5.04	117.97	121.00
1	A	189[B]	TYR	CB-CG-CD2	-5.04	117.97	121.00
1	A	115	PHE	CB-CG-CD2	-5.01	117.29	120.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3043	0	2939	59	0
2	A	48	0	26	4	0
3	A	24	0	11	0	0
4	A	13	0	5	1	0
5	A	671	0	0	22	0
All	All	3799	0	2981	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:PRO:HG3	1:A:293[A]:ARG:NH1	1.76	0.99
1:A:94[A]:LYS:HE2	5:A:460[A]:HOH:O	1.76	0.84
1:A:136[B]:ASN:C	1:A:136[B]:ASN:OD1	2.24	0.74
1:A:293[C]:ARG:HG2	1:A:293[C]:ARG:HH11	1.54	0.72
1:A:26[A]:GLN:HG3	5:A:3009[A]:HOH:O	1.88	0.72
1:A:194[A]:LYS:HD2	5:A:2411:HOH:O	1.91	0.70
1:A:225:PRO:HD2	1:A:295[B]:TRP:CZ2	2.28	0.68
1:A:222[A]:PRO:HB2	1:A:223[A]:GLU:OE2	1.94	0.68
1:A:154[B]:LYS:HD3	5:A:2444:HOH:O	1.96	0.65
1:A:94[A]:LYS:HE3	1:A:98[A]:ASP:OD2	1.98	0.64
1:A:221[A]:LYS:CB	1:A:222[A]:PRO:HD2	2.26	0.64
1:A:209:TYR:CE1	2:A:318:NDP:H41N	2.34	0.63
1:A:137[B]:ILE:HG12	5:A:3113[B]:HOH:O	1.98	0.63
1:A:94[A]:LYS:NZ	1:A:98[A]:ASP:OD1	2.25	0.62
1:A:209:TYR:CZ	2:A:318:NDP:H41N	2.35	0.62
1:A:315[A]:PHE:N	1:A:315[A]:PHE:CD1	2.70	0.60
1:A:221[A]:LYS:HB3	1:A:222[A]:PRO:HD2	1.86	0.58
1:A:221[A]:LYS:HB2	1:A:221[A]:LYS:HZ2	1.71	0.55
1:A:293[C]:ARG:CG	1:A:293[C]:ARG:HH11	2.16	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:293[C]:ARG:HB3	1:A:293[C]:ARG:CZ	2.38	0.53
1:A:136[A]:ASN:HB2	5:A:2085:HOH:O	2.09	0.53
1:A:244[B]:THR:HG22	5:A:2140:HOH:O	2.08	0.53
1:A:138:LEU:HG	5:A:2319[B]:HOH:O	2.09	0.52
1:A:94[A]:LYS:HD2	1:A:98[A]:ASP:OD1	2.09	0.52
1:A:20:TRP:CE3	1:A:21:LYS:HE3	2.44	0.52
1:A:225:PRO:HG3	1:A:293[A]:ARG:HH12	1.67	0.52
1:A:164[A]:LEU:C	1:A:164[A]:LEU:HD23	2.30	0.52
1:A:190[A]:LEU:HB2	1:A:295[A]:TRP:O	2.12	0.49
1:A:221[A]:LYS:HB2	1:A:221[A]:LYS:NZ	2.27	0.49
1:A:221[A]:LYS:O	1:A:222[A]:PRO:C	2.51	0.48
1:A:181:VAL:HA	1:A:205:VAL:O	2.14	0.48
1:A:225:PRO:O	5:A:2145[A]:HOH:O	2.20	0.48
1:A:305[B]:SER:HB3	5:A:3585[B]:HOH:O	2.14	0.47
1:A:70[A]:GLU:HG2	5:A:8100:HOH:O	2.15	0.47
1:A:244[B]:THR:HG21	5:A:6087[B]:HOH:O	2.15	0.46
1:A:94[A]:LYS:CE	5:A:460[A]:HOH:O	2.48	0.46
1:A:103:TYR:CE2	1:A:154[B]:LYS:HE3	2.51	0.46
1:A:293[C]:ARG:NH1	1:A:293[C]:ARG:CG	2.74	0.46
1:A:15[A]:LEU:HD23	1:A:273:PHE:CD1	2.51	0.46
1:A:234[B]:LYS:HE3	5:A:2595:HOH:O	2.15	0.45
1:A:194[A]:LYS:CE	5:A:2411:HOH:O	2.65	0.44
1:A:136[B]:ASN:O	1:A:136[B]:ASN:OD1	2.35	0.44
1:A:225:PRO:HD2	1:A:295[B]:TRP:CE2	2.53	0.43
1:A:81:THR:O	5:A:3760[B]:HOH:O	2.21	0.43
1:A:234[B]:LYS:CE	5:A:2595:HOH:O	2.66	0.43
1:A:221[A]:LYS:CB	1:A:221[A]:LYS:NZ	2.82	0.43
1:A:262:LYS:O	2:A:318:NDP:H8A	2.20	0.42
1:A:23:PRO:HA	1:A:24[B]:PRO:HD3	1.94	0.42
1:A:277[B]:ASP:CG	5:A:3551[B]:HOH:O	2.58	0.42
1:A:1:ALA:HB3	5:A:2462:HOH:O	2.18	0.42
1:A:94[A]:LYS:CE	1:A:98[A]:ASP:OD2	2.67	0.41
1:A:256[B]:ASN:ND2	5:A:3630[B]:HOH:O	2.53	0.41
1:A:183:GLN:OE1	2:A:318:NDP:H2N	2.20	0.41
1:A:163[A]:HIS:O	1:A:164[A]:LEU:C	2.58	0.41
1:A:53:GLU:OE1	4:A:450[A]:CIT:O2	2.38	0.41
1:A:82:TYR:OH	5:A:2236:HOH:O	2.22	0.41
1:A:194[A]:LYS:HE2	5:A:2411:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/316 (127%)	395 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/281 (129%)	358 (99%)	4 (1%)	80	48

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26[A]	GLN
1	A	26[B]	GLN
1	A	119[A]	LYS
1	A	126[A]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NDP	A	318	-	42,52,52	1.29	5 (11%)	55,80,80	1.32	6 (10%)
3	LDT	A	320	-	22,25,25	1.91	5 (22%)	29,34,34	1.56	6 (20%)
4	CIT	A	450[A]	-	3,12,12	0.85	0	3,17,17	2.63	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	318	-	-	0/30/77/77	0/5/5/5
3	LDT	A	320	-	-	0/12/14/14	0/2/2/2
4	CIT	A	450[A]	-	-	0/6/16/16	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	320	LDT	BR8-C25	-6.25	1.76	1.90
2	A	318	NDP	C4N-C5N	-4.68	1.39	1.49
3	A	320	LDT	C11-N17	-2.92	1.31	1.33
2	A	318	NDP	PA-O2A	-2.29	1.45	1.54

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	320	LDT	F9-C5	-2.02	1.31	1.36
2	A	318	NDP	C2N-C3N	2.03	1.39	1.34
3	A	320	LDT	C6-C5	2.05	1.41	1.37
2	A	318	NDP	C6N-C5N	2.30	1.37	1.33
3	A	320	LDT	C2-C5	2.96	1.42	1.37
2	A	318	NDP	C2A-N3A	3.03	1.37	1.32

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	318	NDP	N3A-C2A-N1A	-5.70	124.53	128.89
4	A	450[A]	CIT	C3-C2-C1	-4.19	108.26	114.96
3	A	320	LDT	BR8-C25-C28	-4.05	113.38	119.28
3	A	320	LDT	C26-C29-C25	-3.23	114.48	119.17
3	A	320	LDT	C7-C11-S16	-3.19	117.59	121.75
2	A	318	NDP	C3N-C2N-N1N	-2.60	119.41	123.14
3	A	320	LDT	C28-C27-C24	-2.57	120.86	124.13
2	A	318	NDP	O3-PA-O5B	-2.20	97.11	102.94
2	A	318	NDP	C1D-N1N-C2N	-2.19	117.09	120.91
2	A	318	NDP	PN-O3-PA	2.32	139.24	132.73
3	A	320	LDT	C7-C11-N17	2.34	117.36	115.47
3	A	320	LDT	C29-C25-C28	2.49	125.22	121.44
2	A	318	NDP	C5N-C4N-C3N	2.57	119.61	112.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	318	NDP	4	0
4	A	450[A]	CIT	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	316/316 (100%)	-0.37	7 (2%) 65 62	7, 14, 26, 50	34 (10%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	315[A]	PHE	6.4
1	A	314[A]	GLU	4.7
1	A	222[A]	PRO	4.6
1	A	313[A]	GLU	3.4
1	A	312[A]	HIS	3.2
1	A	223[A]	GLU	2.7
1	A	0[A]	MET	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CIT	A	450[A]	13/13	0.86	0.31	9.97	13,31,38,43	13
3	LDT	A	320	24/24	0.98	0.05	-0.70	11,14,16,19	2
2	NDP	A	318	48/48	0.99	0.03	-1.34	7,9,13,15	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.