



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:18 AM GMT

PDB ID : 2GK9
Title : Human Phosphatidylinositol-4-phosphate 5-kinase, type II, gamma
Authors : Uppenberg, J.; Hogbom, M.; Ogg, D.; Arrowsmith, C.; Berglund, H.; Collins, R.; Ehn, M.; Flodin, S.; Flores, A.; Graslund, S.; Holmberg-Schiavone, L.; Edwards, A.; Hammarstrom, M.; Kotenyova, T.; Nilsson-Ehle, P.; Nordlund, P.; Nyman, T.; Persson, C.; Sagemark, J.; Stenmark, P.; Sundstrom, M.; Thorsell, A.G.; Van Den Berg, S.; Weigelt, J.; Hallberg, B.M.; Structural Genomics Consortium (SGC)
Deposited on : 2006-03-31
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	<div> <div>14%</div> <div>36%</div> <div>12%</div> <div>38%</div> </div>
1	B	392	<div> <div>3%</div> <div>15%</div> <div>39%</div> <div>10%</div> <div>36%</div> </div>
1	C	392	<div> <div>3%</div> <div>14%</div> <div>39%</div> <div>10%</div> <div>38%</div> </div>
1	D	392	<div> <div>2%</div> <div>15%</div> <div>36%</div> <div>13%</div> <div>36%</div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 8233 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called phosphatidylinositol-4-phosphate 5-kinase, type II, gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	243	Total	C	N	O	S	0	0	0
			2023	1307	343	362	11			
1	B	250	Total	C	N	O	S	0	0	0
			2084	1345	353	375	11			
1	C	245	Total	C	N	O	S	0	0	0
			2042	1317	347	367	11			
1	D	250	Total	C	N	O	S	0	0	0
			2084	1345	353	375	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	SER	-	cloning artifact	UNP Q8TBX8
A	31	MET	-	cloning artifact	UNP Q8TBX8
B	30	SER	-	cloning artifact	UNP Q8TBX8
B	31	MET	-	cloning artifact	UNP Q8TBX8
C	30	SER	-	cloning artifact	UNP Q8TBX8
C	31	MET	-	cloning artifact	UNP Q8TBX8
D	30	SER	-	cloning artifact	UNP Q8TBX8
D	31	MET	-	cloning artifact	UNP Q8TBX8

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: ■ ■ ■ ■

14% 36% 12% 38%

Sequence (Top to Bottom):

THR VAL LEU
 HA04 PRO GLY
 P405 GLY
 P406 GLY
 Q407 PHE
 Y408 GLY
 A409 SER
 K410 PHE
 R411 ILE
 R412 ASP
 F412 VAL
 L413 TYR
 D414 ALA
 I415 ALA
 I416 ILE
 T417 ILE
 ASN SER
 ILE ALA
 ALA


Sequence (Left to Right):

M93 L94
 P95 SER
 S96 LEU
 H97 VAL
 F98 VAL
 K99 SER
 F100 ARG
 S166 GLU
 A164 ALA
 M165 ILE
 L166 ASP
 S167 GLY
 M168 SER
 H169 GLY
 W170 PRO
 Q171 LYS
 Q172 VAL
 Y173 GLY
 I174 GLY
 V174 LEU
 T239 PRO
 K175 ASP
 C176 LYS
 H177 VAL
 G178 ILE
 M179 ASP
 D242 GLY
 M243 GLY
 D244 SER
 L181 GLY
 L182 VAL
 P183 VAL
 Q184 ASP
 K248 GLY
 Y253 CYS
 L254 SER
 G190 LEU
 V191 THR
 S192 GLY
 D193 PRO
 K260 ALA
 L261 LEU
 F262 VAL
 L263 GLY
 K264 SER
 K265 TYR
 L266 GLY
 K267 THR
 R268 ALA
 D269 ALA
 V270 LYS
 E271 THR
 I272 VAL
 L273 LYS
 V274 HIS
 Q275 GLY
 L276 ALA
 I277 ILE
 K278 ALA
 R279 GLY
 D280 ALA
 V281 SER


Sequence (Right to Left):

M70 L71
 L72 L73
 P73 D74
 D75 F76
 K77 A78
 S79 S80
 R81 I82
 K83 V84
 M85 M86
 H87 L88
 F89 H90
 E151 K152
 E153 E154
 V154 S155
 F156


- Chain B:
-
- 3% 15% 39% 10% 36%
- SER MET HIS PHE VAL GLN GLY LYS VAL LYS VAL PHE ARG ALA ALA D45 P46 L47 V48 G49 V50 F51 L52 L53 G54 V55 A56 B57 S58 I59 L62 S63 Q64 V65 P66 P67 P68 V69 M70 L71 L72 P73 D74 D75 F76 F77 A78 S79 S80 K81 K82 I83 K83 N86 H87 L88 L89 F89 R90 R91 R92 N93 L94 P95 F98 F99 F100 K101 L102 Y103 C104 P105 Q106 V107 V108 R109 H110 L111 R112 N117 R113 R114 I117 Q120 D121 A122 L123 V124 S125 L126 T127 R128 S132 E133 E134 S135 GLY SER ASP GLY R140 R141 F141 L142 L143 S144 Y145 D146 R147 T148 L149 V150 I151 K152 D153 V154 F157



• Molecule 1: phosphatidylinositol-4-phosphate 5-kinase, type II, gamma



• Molecule 1: phosphatidylinositol-4-phosphate 5-kinase, type II, gamma



H163	L163	H288	GLU	K409
S164	VAL	D289	SER	K410
N165	SER	I290	PHE	R411
L166	ARG	ILE	ILE	F412
S167	GLU	ARG	ASP	L413
N168	ALA	GLY	VAL	D414
Y169	SER	SER	TYR	F415
H170	ASP	GLU	ALA	I416
Q171	LYS	PRO	ILE	T417
Y172	GLU	GLU	ARG	ASN
I173	GLU	GLU	SER	ILE
V174	VAL	ALA	ALA	PHE
K175	LYS	GLU	GLU	ALA
C176	GLU	PRO	GLY	
H177	LEU	VAL	ALA	
G178	PRO	ARG	PRO	
N179	T239	GLN	GLN	
T180	ASP	LYS	LYS	
L181	GLU	GLU	E366	
L182	L181	SER	Y367	
P183	GLU	SER	Y368	
Q184	VAL	VAL	F369	
F185	ASP	ASP	M370	
L186	GLY	GLY	G371	
G187	ASP	ASP	L372	
M188	Y252	CYS	I373	
Y189	G255	SER	D374	
R190	E256	LEU	I375	
V191	GLY	THR	I376	
S192	E257	THR	THR	
V193	E258	PRO	GLN	
D194	K259	PRO	TYR	
N195	K260	ALA	ASP	
E196	I281	LEU	ALA	
D197	F262	VAL	LYS	
S198	GLY	VAL	LYS	
Y199	E264	SER	LYS	
M200	K265	TYR	ALA	
L201	L266	GLY	ALA	
V202	K267	THR	HIS	
M203	R268	SER	ALA	
R204	D269	PRO	ALA	
N205	V270	GLU	LYS	
M206	E271	THR	THR	
F207	F272	ILE	VAL	
S208	L273	GLY	LYS	
H209	V274	GLY	HIS	
R210	Q275	TYR	GLY	
L211	L276	ILE	ALA	
P212	K277	HIS	GLY	
V213	I278	SER	ALA	
H214	M279	HIS	GLU	
R215	D280	ARG	ILE	
K216	Y281	PRO	SER	
Y217	S282	LEU	THR	
D218	L283	GLY	VAL	
L219	I284	PRO	H404	
K220	L285	GLY	Q407	
G221	G286	GLU	Y408	
SER	I287	PHE		

4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	95.39Å 95.39Å 189.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80 29.74 – 2.79	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-2.80) 95.5 (29.74-2.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.80Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.254 , 0.297 0.256 , 0.263	Depositor DCC
R_{free} test set	1894 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	65.5	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 85.7	EDS
Estimated twinning fraction	0.469 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	1 of 40811 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	8233	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2068	0.58	4/2784 (0.1%)
1	B	0.47	0/2132	0.52	2/2872 (0.1%)
1	C	0.50	0/2088	0.58	5/2813 (0.2%)
1	D	0.45	0/2132	0.55	1/2872 (0.0%)
All	All	0.47	0/8420	0.56	12/11341 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	D	0	1
All	All	0	3

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	114	ARG	NE-CZ-NH2	-5.55	117.52	120.30
1	C	253	TYR	CB-CG-CD2	-5.31	117.81	121.00
1	A	244	ASP	CB-CG-OD2	5.27	123.05	118.30
1	B	242	ASP	CB-CG-OD2	5.26	123.03	118.30
1	C	414	ASP	CB-CG-OD2	5.26	123.03	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	279	MET	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	91	ARG	Peptide
1	D	128	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2023	0	2012	301	0
1	B	2084	0	2068	279	0
1	C	2042	0	2028	277	0
1	D	2084	0	2068	265	0
All	All	8233	0	8176	1077	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 66.

The worst 5 of 1077 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:266:LEU:CD2	1:C:413:LEU:HD23	1.52	1.39
1:C:176:CYS:SG	1:C:180:THR:HG22	1.65	1.34
1:A:72:LEU:HD21	1:B:95:PRO:CA	1.64	1.25
1:A:72:LEU:CD2	1:B:95:PRO:HA	1.67	1.24
1:D:285:LEU:CD2	1:D:370:MET:HG2	1.66	1.24

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/392 (58%)	189 (82%)	39 (17%)	1 (0%)	39	74
1	B	240/392 (61%)	203 (85%)	37 (15%)	0	100	100
1	C	233/392 (59%)	197 (84%)	35 (15%)	1 (0%)	39	74
1	D	240/392 (61%)	194 (81%)	45 (19%)	1 (0%)	39	74
All	All	942/1568 (60%)	783 (83%)	156 (17%)	3 (0%)	46	79

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	375	ILE
1	C	241	ARG
1	A	95	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/348 (66%)	175 (76%)	56 (24%)	1	2
1	B	237/348 (68%)	182 (77%)	55 (23%)	1	3
1	C	233/348 (67%)	181 (78%)	52 (22%)	1	3
1	D	237/348 (68%)	177 (75%)	60 (25%)	1	2
All	All	938/1392 (67%)	715 (76%)	223 (24%)	1	2

5 of 223 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	289	ASP
1	C	126	LEU
1	D	252	VAL
1	B	370	MET
1	C	62	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	86	ASN
1	C	87	HIS
1	D	214	HIS
1	B	195	ASN
1	B	247	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/392 (61%)	-0.09	3 (1%) 81 73	24, 58, 114, 158	0
1	B	250/392 (63%)	0.03	12 (4%) 34 23	24, 58, 138, 152	0
1	C	245/392 (62%)	-0.06	10 (4%) 41 29	25, 59, 128, 158	0
1	D	250/392 (63%)	-0.05	9 (3%) 46 34	24, 60, 144, 158	0
All	All	988/1568 (63%)	-0.04	34 (3%) 49 36	24, 59, 136, 158	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	93	ASN	5.3
1	B	416	ILE	4.8
1	A	405	PRO	4.2
1	A	253	TYR	4.2
1	B	253	TYR	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.