



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:19 AM GMT

PDB ID : 2GMY  
Title : Crystal Structure of a Protein of Unknown Function ATU0492 from *Agrobacterium tumefaciens*, Putative Antioxidant Defence Protein AhpD  
Authors : Zhang, R.; Xu, X.; Gu, J.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2006-04-07  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	<div> <div>82%</div> <div>14%</div> <div>• •</div> </div>
1	B	153	<div> <div>2%</div> <div>84%</div> <div>12%</div> <div>5%</div> </div>
1	C	153	<div> <div>86%</div> <div>9%</div> <div>• 5%</div> </div>
1	D	153	<div> <div>88%</div> <div>7%</div> <div>5%</div> </div>
1	E	153	<div> <div>89%</div> <div>6%</div> <div>5%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	153	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: a small red segment at the beginning labeled '2%', a large green segment labeled '88%', a small yellow segment labeled '8%', and a small grey segment at the end. A small black dot is located at the far right end of the bar.

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7730 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein Atu0492.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	147	Total	C	N	O	S	0	0	0
			1167	740	207	215	5			
1	B	146	Total	C	N	O	S	0	0	0
			1161	737	206	212	6			
1	C	146	Total	C	N	O	S	0	0	0
			1162	737	206	214	5			
1	D	145	Total	C	N	O	S	0	0	0
			1153	732	205	211	5			
1	E	145	Total	C	N	O	S	0	0	0
			1153	732	205	211	5			
1	F	147	Total	C	N	O	S	0	0	0
			1167	740	207	215	5			



- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	141	Total	O	0	0
			141	141		
2	B	120	Total	O	0	0
			120	120		
2	C	122	Total	O	0	0
			122	122		
2	D	138	Total	O	0	0
			138	138		
2	E	112	Total	O	0	0
			112	112		
2	F	134	Total	O	0	0
			134	134		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Hypothetical protein Atu0492




- Molecule 1: Hypothetical protein Atu0492




- Molecule 1: Hypothetical protein Atu0492




- Molecule 1: Hypothetical protein Atu0492



- Molecule 1: Hypothetical protein Atu0492



- Molecule 1: Hypothetical protein Atu0492



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.01Å 75.01Å 148.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	64.96 – 1.60 39.41 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (64.96-1.60) 99.7 (39.41-1.60)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.167 , 0.191 0.166 , 0.190	Depositor DCC
$R_{free}$ test set	6180 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.3	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 54.1	EDS
Estimated twinning fraction	0.015 for -h,-k,l 0.057 for h,-h-k,-l 0.026 for -k,-h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 123121 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7730	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/1193	0.59	1/1617 (0.1%)
1	B	0.44	0/1187	0.52	0/1608
1	C	0.50	0/1188	0.60	1/1610 (0.1%)
1	D	0.47	0/1179	0.55	0/1598
1	E	0.43	0/1179	0.52	0/1598
1	F	0.47	0/1193	0.56	0/1617
All	All	0.46	0/7119	0.56	2/9648 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	72	MET	CG-SD-CE	-8.46	86.66	100.20
1	A	72	MET	CG-SD-CE	-6.93	89.10	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1167	0	1149	24	0
1	B	1161	0	1150	17	0
1	C	1162	0	1144	15	0
1	D	1153	0	1138	11	0
1	E	1153	0	1138	9	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1167	0	1149	9	0
2	A	141	0	0	8	1
2	B	120	0	0	3	0
2	C	122	0	0	2	0
2	D	138	0	0	4	1
2	E	112	0	0	0	0
2	F	134	0	0	1	0
All	All	7730	0	6868	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:ASN:HD22	1:A:134:ASN:HD22	1.09	1.00
1:D:97:LYS:HE3	2:D:290:HOH:O	1.63	0.99
1:D:46:ASN:HD22	1:D:134:ASN:HD22	1.16	0.94
1:A:72:MET:CE	2:A:269:HOH:O	2.15	0.94
1:A:72:MET:SD	2:A:269:HOH:O	2.30	0.89
1:B:46:ASN:HD22	1:B:134:ASN:HD22	1.21	0.88
1:F:46:ASN:HD22	1:F:134:ASN:HD22	1.19	0.87
1:A:33:ARG:HD3	2:A:206:HOH:O	1.85	0.75
1:A:70:ASN:HD21	1:E:70:ASN:HD22	1.35	0.75
1:A:72:MET:HE1	2:A:269:HOH:O	1.79	0.72
1:C:33:ARG:HD2	2:C:190:HOH:O	1.90	0.70
1:D:36:HIS:HE1	1:D:62:ASP:OD2	1.78	0.67
1:F:129:ALA:HB1	2:F:282:HOH:O	1.95	0.66
1:D:60:ARG:HH22	1:F:70:ASN:HD21	1.44	0.64
1:A:108:PHE:CE2	1:A:112:ARG:HD2	2.32	0.64
1:D:36:HIS:HD2	2:D:192:HOH:O	1.83	0.61
1:C:147:GLU:HG3	2:C:239:HOH:O	1.99	0.61
1:A:72:MET:O	1:A:72:MET:HG3	2.03	0.58
1:A:100:GLU:HG3	2:A:218:HOH:O	2.05	0.56
1:A:147:GLU:H	1:E:67:GLN:HE22	1.53	0.55
1:A:2:LYS:N	2:A:288:HOH:O	2.40	0.54
1:D:100:GLU:HG2	2:D:290:HOH:O	2.07	0.54
1:B:60:ARG:HH22	1:C:70:ASN:HD21	1.54	0.53
1:A:147:GLU:H	1:E:67:GLN:NE2	2.08	0.52
1:A:13:GLU:HG2	2:A:202:HOH:O	2.11	0.49
1:B:15:PHE:CZ	1:B:19:MET:HE3	2.48	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:LYS:HD3	1:A:55:HIS:CE1	2.47	0.49
1:B:71:LEU:HD23	1:C:144:HIS:NE2	2.27	0.49
1:B:114:HIS:HE1	2:B:200:HOH:O	1.96	0.49
1:C:35:ILE:CG2	1:C:39:LYS:HE2	2.43	0.48
1:A:70:ASN:ND2	1:E:70:ASN:HD22	2.09	0.48
1:B:16:LYS:HD2	2:B:245:HOH:O	2.14	0.48
1:B:26:GLN:NE2	1:B:39:LYS:HZ2	2.12	0.48
1:C:78:SER:OG	1:C:80:VAL:HG12	2.14	0.48
1:C:26:GLN:NE2	1:C:39:LYS:HE3	2.28	0.47
1:D:39:LYS:HD3	1:D:55:HIS:CE1	2.49	0.47
1:B:71:LEU:HD21	1:C:144:HIS:CE1	2.49	0.47
1:C:72:MET:O	1:C:72:MET:HG3	2.15	0.46
1:A:67:GLN:HG2	1:A:71:LEU:HD12	1.96	0.46
1:F:35:ILE:CG2	1:F:39:LYS:HE2	2.45	0.46
1:B:71:LEU:CD2	1:C:144:HIS:CE1	2.98	0.46
1:A:18:VAL:HG12	1:A:133:TRP:CZ2	2.51	0.46
1:B:114:HIS:HD2	2:B:228:HOH:O	1.98	0.46
1:D:35:ILE:CG2	1:D:39:LYS:HE2	2.46	0.45
1:B:47:GLY:HA2	1:C:74:VAL:HG12	1.99	0.45
1:C:39:LYS:HD3	1:C:55:HIS:CE1	2.51	0.45
1:A:71:LEU:HD13	1:A:78:SER:OG	2.17	0.45
1:B:56:VAL:HG13	1:B:69:ILE:HG21	1.99	0.44
1:D:97:LYS:HD2	2:D:288:HOH:O	2.17	0.44
1:F:18:VAL:HG12	1:F:133:TRP:CZ2	2.53	0.44
1:E:22:GLU:OE2	1:E:39:LYS:NZ	2.51	0.43
1:A:70:ASN:HD21	1:E:70:ASN:ND2	2.10	0.43
1:B:71:LEU:CD2	1:C:144:HIS:NE2	2.81	0.43
1:D:47:GLY:HA2	1:F:74:VAL:HG12	2.00	0.43
1:E:39:LYS:HD3	1:E:55:HIS:CE1	2.53	0.43
1:B:39:LYS:HD3	1:B:55:HIS:CE1	2.52	0.43
1:A:55:HIS:ND1	1:A:58:GLU:OE1	2.41	0.43
1:B:37:LEU:HD23	1:B:37:LEU:C	2.39	0.43
1:E:91:TRP:CG	1:E:111:LEU:HD22	2.54	0.42
1:A:31:GLU:OE2	1:A:33:ARG:NH1	2.53	0.41
1:B:71:LEU:HD23	1:B:78:SER:OG	2.20	0.41
1:D:60:ARG:HH22	1:F:70:ASN:ND2	2.15	0.41
1:C:91:TRP:CG	1:C:111:LEU:HD22	2.55	0.41
1:A:13:GLU:CG	2:A:202:HOH:O	2.68	0.41
1:F:37:LEU:HD23	1:F:37:LEU:C	2.41	0.41
1:A:74:VAL:HG12	1:E:47:GLY:HA2	2.02	0.41
1:F:33:ARG:HG3	1:F:64:LEU:HD21	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:SER:OG	1:A:80:VAL:HG12	2.21	0.40
1:B:74:VAL:HG12	1:C:47:GLY:HA2	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:270:HOH:O	2:D:165:HOH:O[1_665]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/153 (95%)	142 (98%)	3 (2%)	0	100	100
1	B	144/153 (94%)	141 (98%)	3 (2%)	0	100	100
1	C	144/153 (94%)	141 (98%)	3 (2%)	0	100	100
1	D	143/153 (94%)	140 (98%)	3 (2%)	0	100	100
1	E	143/153 (94%)	140 (98%)	3 (2%)	0	100	100
1	F	145/153 (95%)	142 (98%)	3 (2%)	0	100	100
All	All	864/918 (94%)	846 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/125 (98%)	123 (100%)	0	100	100
1	B	123/125 (98%)	122 (99%)	1 (1%)	86	75
1	C	123/125 (98%)	123 (100%)	0	100	100
1	D	122/125 (98%)	122 (100%)	0	100	100
1	E	122/125 (98%)	121 (99%)	1 (1%)	86	75
1	F	123/125 (98%)	123 (100%)	0	100	100
All	All	736/750 (98%)	734 (100%)	2 (0%)	94	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1	MET
1	E	2	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	32	HIS
1	A	70	ASN
1	A	134	ASN
1	B	26	GLN
1	B	70	ASN
1	B	114	HIS
1	B	134	ASN
1	C	26	GLN
1	C	46	ASN
1	C	70	ASN
1	C	134	ASN
1	C	143	GLN
1	D	36	HIS
1	D	70	ASN
1	D	134	ASN
1	E	26	GLN
1	E	46	ASN
1	E	55	HIS
1	E	67	GLN
1	E	134	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	70	ASN
1	F	134	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	147/153 (96%)	-0.20	2 (1%) 78 77	10, 16, 29, 39	0
1	B	146/153 (95%)	-0.25	3 (2%) 67 65	12, 18, 26, 39	0
1	C	146/153 (95%)	-0.26	1 (0%) 89 89	11, 17, 26, 33	0
1	D	145/153 (94%)	-0.32	2 (1%) 78 77	10, 16, 26, 34	0
1	E	145/153 (94%)	-0.13	2 (1%) 78 77	12, 20, 30, 35	0
1	F	147/153 (96%)	-0.16	3 (2%) 68 67	11, 17, 30, 36	0
All	All	876/918 (95%)	-0.22	13 (1%) 76 75	10, 17, 29, 39	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	4.9
1	A	148	ALA	4.1
1	D	2	LYS	3.5
1	C	147	GLU	2.8
1	E	7	TYR	2.6
1	A	2	LYS	2.5
1	B	7	TYR	2.3
1	F	2	LYS	2.3
1	E	9	LYS	2.3
1	F	9	LYS	2.2
1	F	148	ALA	2.2
1	B	32	HIS	2.2
1	D	9	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.