



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:57 AM GMT

PDB ID : 3GOV
Title : Crystal structure of the catalytic region of human MASP-1
Authors : Harmat, V.; Dobo, J.; Beinrohr, L.; Sebestyen, E.; Zavodszky, P.; Gal, P.
Deposited on : 2009-03-20
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4549 (2.58-2.50)
Clashscore	102246	5292 (2.58-2.50)
Ramachandran outliers	100387	5194 (2.58-2.50)
Sidechain outliers	100360	5196 (2.58-2.50)
RSRZ outliers	91569	4561 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	155	
2	B	251	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3196 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MASP-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	147	Total	C	N	O	S	0	4	0
			1163	744	189	218	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	294	ALA	-	EXPRESSION TAG	UNP P48740
A	295	SER	-	EXPRESSION TAG	UNP P48740
A	296	MET	-	EXPRESSION TAG	UNP P48740
A	297	THR	-	EXPRESSION TAG	UNP P48740

- Molecule 2 is a protein called MASP-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	251	Total	C	N	O	S	0	1	0
			1931	1226	332	360	13			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	499	LYS	GLU	SEE REMARK 999	UNP P48740

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	33	Total	O	0	0
			33	33		
4	B	57	Total	O	0	0
			57	57		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: MASP-1



• Molecule 2: MASP-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.42Å 70.41Å 121.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.98 – 2.55 45.41 – 2.55	Depositor EDS
% Data completeness (in resolution range)	97.9 (45.98-2.55) 97.9 (45.41-2.55)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.229 , 0.278 0.246 , 0.290	Depositor DCC
R_{free} test set	989 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	33.6	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 32.1	EDS
Estimated twinning fraction	0.036 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	1 of 19340 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3196	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1202	0.57	0/1628
2	B	0.35	0/1986	0.57	2/2701 (0.1%)
All	All	0.36	0/3188	0.57	2/4329 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	2
All	All	0	4

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	621	LEU	N-CA-C	-5.39	96.45	111.00
2	B	528	GLU	N-CA-C	5.12	124.82	111.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	382	THR	Peptide
1	A	406	ASN	Peptide
2	B	527	ASP	Peptide
2	B	620	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1163	0	1141	11	0
2	B	1931	0	1837	38	0
3	A	6	0	8	2	0
3	B	6	0	8	0	0
4	A	33	0	0	0	0
4	B	57	0	0	0	0
All	All	3196	0	2994	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:LEU:HD22	1:A:359:ILE:HD11	1.21	1.10
2:B:695:THR:CG2	2:B:697:VAL:HG23	1.95	0.96
2:B:695:THR:HG22	2:B:697:VAL:HG23	1.50	0.94
2:B:485:ILE:CD1	2:B:516:ILE:HG21	2.01	0.91
2:B:619:ALA:HA	2:B:620:PRO:C	1.99	0.83
2:B:584:VAL:HG12	2:B:652:THR:CG2	2.10	0.81
2:B:485:ILE:HD13	2:B:516:ILE:HG21	1.62	0.80
2:B:527:ASP:HB3	2:B:529:ASN:H	1.51	0.74
2:B:620:PRO:HB2	2:B:621:LEU:HG	1.76	0.67
2:B:484:TRP:CD1	2:B:695:THR:HG23	2.31	0.65
2:B:485:ILE:HD11	2:B:516:ILE:HG21	1.79	0.64
2:B:608:ILE:HD13	2:B:630:ILE:HD11	1.80	0.64
2:B:465:MET:CE	2:B:517:ILE:HD12	2.27	0.64
2:B:695:THR:HG21	2:B:697:VAL:HG23	1.78	0.63
2:B:527:ASP:HB3	2:B:529:ASN:N	2.13	0.63
2:B:695:THR:CG2	2:B:697:VAL:CG2	2.75	0.63
2:B:695:THR:HG21	2:B:697:VAL:CG2	2.30	0.61
2:B:465:MET:HE2	2:B:517:ILE:HD12	1.81	0.61
1:A:304:LEU:CD2	1:A:359:ILE:HD11	2.15	0.61
2:B:584:VAL:HG12	2:B:652:THR:HG23	1.82	0.61
2:B:465:MET:HE1	2:B:520:LYS:HE2	1.82	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:654:ASN:HD22	2:B:655:ARG:N	2.00	0.59
2:B:619:ALA:HA	2:B:621:LEU:N	2.18	0.57
1:A:304:LEU:HD22	1:A:359:ILE:CD1	2.15	0.54
2:B:578:GLN:HG3	2:B:663:VAL:CG1	2.38	0.53
2:B:665:THR:HG23	2:B:684:ILE:HD11	1.91	0.52
2:B:578:GLN:HG3	2:B:663:VAL:HG11	1.92	0.51
2:B:579:GLN:CD	2:B:579:GLN:H	2.15	0.50
2:B:485:ILE:HD13	2:B:516:ILE:CG2	2.39	0.49
2:B:620:PRO:HB2	2:B:621:LEU:CG	2.42	0.49
1:A:313:GLU:HB2	3:A:1:GOL:H32	1.94	0.49
2:B:485:ILE:HD12	2:B:558:LEU:HD21	1.94	0.48
2:B:540:LEU:N	2:B:540:LEU:HD23	2.27	0.48
1:A:441:PHE:O	2:B:459:THR:HG22	2.15	0.47
2:B:606:ILE:HD12	2:B:632:ALA:HB1	1.98	0.46
2:B:478:SER:OG	2:B:649:PRO:HB3	2.17	0.45
1:A:406:ASN:ND2	1:A:408:ASN:O	2.51	0.44
2:B:620:PRO:HB2	2:B:621:LEU:CA	2.48	0.44
2:B:584:VAL:HA	2:B:652:THR:HG22	2.01	0.43
2:B:600:THR:HG22	2:B:601:LEU:O	2.19	0.43
1:A:326:LEU:HB2	3:A:1:GOL:H31	2.02	0.41
1:A:347:ILE:HG22	1:A:357:ASN:HB2	2.01	0.41
1:A:382:THR:HG22	1:A:383:ARG:H	1.84	0.41
2:B:465:MET:HE1	2:B:517:ILE:HD12	2.02	0.41
1:A:382:THR:HG22	1:A:383:ARG:N	2.36	0.41
1:A:308:VAL:HG12	1:A:309:HIS:CD2	2.55	0.40
2:B:619:ALA:CA	2:B:620:PRO:C	2.77	0.40
2:B:600:THR:HG22	2:B:601:LEU:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/155 (96%)	142 (95%)	6 (4%)	1 (1%)	26	44
2	B	250/251 (100%)	237 (95%)	10 (4%)	3 (1%)	16	27
All	All	399/406 (98%)	379 (95%)	16 (4%)	4 (1%)	19	33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	407	ASN
2	B	528	GLU
2	B	620	PRO
2	B	670	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/140 (94%)	128 (98%)	3 (2%)	58	82
2	B	203/215 (94%)	198 (98%)	5 (2%)	55	80
All	All	334/355 (94%)	326 (98%)	8 (2%)	57	81

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	338	ASP
1	A	385	ASN
1	A	408	ASN
2	B	502	THR
2	B	509	LEU
2	B	592	GLN
2	B	593	PHE
2	B	654	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	309	HIS
1	A	385	ASN
2	B	543	GLN
2	B	592	GLN
2	B	654	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	1	-	5,5,5	0.38	0	5,5,5	0.58	0
3	GOL	B	2	-	5,5,5	0.40	0	5,5,5	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1	-	-	0/4/4/4	0/0/0/0
3	GOL	B	2	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	147/155 (94%)	0.43	5 (3%)	49	55	41, 48, 58, 65	0
2	B	251/251 (100%)	0.53	18 (7%)	18	21	36, 49, 58, 62	0
All	All	398/406 (98%)	0.49	23 (5%)	26	31	36, 48, 58, 65	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	656	GLU	5.7
2	B	673	GLY	3.8
2	B	676	ASP	3.7
2	B	620	PRO	3.5
2	B	621	LEU	3.4
2	B	593	PHE	3.3
2	B	594	LEU	3.2
1	A	407	ASN	3.1
2	B	678	TYR	2.9
2	B	622	LYS	2.8
1	A	382	THR	2.8
2	B	464	ALA	2.8
2	B	619	ALA	2.7
2	B	579	GLN	2.6
2	B	655	ARG	2.5
1	A	405	LEU	2.4
1	A	406	ASN	2.4
2	B	614	CYS	2.4
2	B	635	LYS	2.2
2	B	670	ASP	2.1
2	B	638	GLY	2.1
2	B	636	GLU	2.1
1	A	383	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	B	2	6/6	0.89	0.18	0.77	59,60,60,60	0
3	GOL	A	1	6/6	0.89	0.15	-1.82	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.