



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:58 PM GMT

PDB ID : 4GT4  
Title : Structure of unliganded, inactive Ror2 kinase domain  
Authors : Mendrola, J.M.; Lemmon, M.A.  
Deposited on : 2012-08-28  
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	
1	B	308	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NO3	A	801	-	X	-	-
2	NO3	A	802	-	X	-	-
2	NO3	B	801	-	X	-	X
2	NO3	B	802	-	X	-	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4518 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase transmembrane receptor ROR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	1	0
			2146	1395	360	376	15			
1	B	273	Total	C	N	O	S	0	0	0
			2128	1386	359	368	15			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	446	HIS	-	EXPRESSION TAG	UNP Q01974
A	447	HIS	-	EXPRESSION TAG	UNP Q01974
A	448	HIS	-	EXPRESSION TAG	UNP Q01974
A	449	HIS	-	EXPRESSION TAG	UNP Q01974
A	450	HIS	-	EXPRESSION TAG	UNP Q01974
A	451	HIS	-	EXPRESSION TAG	UNP Q01974
B	446	HIS	-	EXPRESSION TAG	UNP Q01974
B	447	HIS	-	EXPRESSION TAG	UNP Q01974
B	448	HIS	-	EXPRESSION TAG	UNP Q01974
B	449	HIS	-	EXPRESSION TAG	UNP Q01974
B	450	HIS	-	EXPRESSION TAG	UNP Q01974
B	451	HIS	-	EXPRESSION TAG	UNP Q01974

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	N	O	0	0
			4	1	3		
2	A	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	123	Total	O	0	0
			123	123		
3	B	105	Total	O	0	0
			105	105		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.83Å 112.92Å 114.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.80 – 2.41 46.92 – 2.41	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.80-2.41) 99.9 (46.92-2.41)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.26 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1108)	Depositor
R, $R_{free}$	0.177 , 0.205 0.177 , 0.206	Depositor DCC
$R_{free}$ test set	1328 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.9	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 60.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 26257 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4518	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.21	0/2201	0.37	0/2988
1	B	0.21	0/2184	0.37	0/2968
All	All	0.21	0/4385	0.37	0/5956

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2146	0	2048	8	0
1	B	2128	0	2037	12	0
2	A	8	0	0	0	0
2	B	8	0	0	0	0
3	A	123	0	0	2	0
3	B	105	0	0	0	0
All	All	4518	0	4085	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:672:ASP:OD1	1:B:738:ARG:NH2	2.24	0.71
1:B:736:ARG:O	1:B:738:ARG:NH1	2.33	0.62
1:B:528:ARG:NH2	1:B:537:VAL:O	2.36	0.56
1:A:582:THR:HG22	1:B:670:SER:HB3	1.88	0.54
1:B:645:TYR:HB3	1:B:653:LEU:HB3	1.91	0.52
1:A:645:TYR:HB3	1:A:653:LEU:HB3	1.90	0.52
1:A:709:GLN:NE2	3:A:937:HOH:O	2.42	0.51
1:B:478:GLU:HA	1:B:488:TYR:HA	1.93	0.50
1:A:465:LEU:HD13	1:A:542:VAL:HG21	1.94	0.49
1:A:638:ARG:NH1	3:A:959:HOH:O	2.36	0.49
1:B:510:LYS:HB2	1:B:513:ALA:HB2	1.95	0.47
1:A:514:GLU:O	1:A:517:LEU:N	2.49	0.45
1:B:465:LEU:HD23	1:B:542:VAL:HG21	1.99	0.45
1:B:517:LEU:HD23	1:B:517:LEU:HA	1.81	0.43
1:B:648:LEU:HD21	1:B:664:ILE:HG23	2.01	0.42
1:A:465:LEU:HD21	1:A:529:ALA:HB2	2.02	0.41
1:A:533:HIS:CG	1:A:534:PRO:HD2	2.56	0.41
1:B:738:ARG:HD2	1:B:738:ARG:N	2.35	0.40
1:B:544:THR:HA	1:B:549:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/308 (86%)	257 (97%)	8 (3%)	0	100	100
1	B	265/308 (86%)	260 (98%)	5 (2%)	0	100	100
All	All	530/616 (86%)	517 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/271 (80%)	212 (97%)	6 (3%)	51	72
1	B	215/271 (79%)	208 (97%)	7 (3%)	45	66
All	All	433/542 (80%)	420 (97%)	13 (3%)	48	70

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	487	VAL
1	A	627	LEU
1	A	703	GLU
1	A	710	VAL
1	A	732	PHE
1	A	749	TRP
1	B	466	LYS
1	B	487	VAL
1	B	504	VAL
1	B	517	LEU
1	B	709	GLN
1	B	710	VAL
1	B	738	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NO3	A	801	-	3,3,3	3.29	3 (100%)	3,3,3	0.13	0
2	NO3	A	802	-	3,3,3	3.29	3 (100%)	3,3,3	0.11	0
2	NO3	B	801	-	3,3,3	3.30	3 (100%)	3,3,3	0.12	0
2	NO3	B	802	-	3,3,3	3.30	3 (100%)	3,3,3	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NO3	A	801	-	-	0/0/0/0	0/0/0/0
2	NO3	A	802	-	-	0/0/0/0	0/0/0/0
2	NO3	B	801	-	-	0/0/0/0	0/0/0/0
2	NO3	B	802	-	-	0/0/0/0	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	NO3	O3-N	2.91	1.40	1.25
2	A	802	NO3	O3-N	2.92	1.40	1.25
2	B	802	NO3	O2-N	2.92	1.40	1.25
2	B	801	NO3	O3-N	2.92	1.40	1.25
2	B	802	NO3	O3-N	2.92	1.40	1.25
2	B	801	NO3	O2-N	2.92	1.40	1.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	802	NO3	O2-N	2.93	1.40	1.25
2	A	801	NO3	O2-N	2.93	1.40	1.25
2	A	801	NO3	O1-N	3.92	1.40	1.24
2	A	802	NO3	O1-N	3.93	1.40	1.24
2	B	802	NO3	O1-N	3.94	1.40	1.24
2	B	801	NO3	O1-N	3.94	1.40	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	274/308 (88%)	0.03	17 (6%) 24 25	13, 33, 79, 97	0
1	B	273/308 (88%)	-0.03	13 (4%) 34 35	16, 36, 76, 89	0
All	All	547/616 (88%)	0.00	30 (5%) 29 29	13, 34, 78, 97	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	471	SER	4.1
1	A	749	TRP	3.8
1	A	465	LEU	3.7
1	A	464	LYS	3.7
1	A	751	ASN	3.3
1	A	468	ILE	3.2
1	A	543	VAL	3.0
1	B	465	LEU	2.9
1	B	469	SER	2.8
1	A	470	LEU	2.7
1	A	473	VAL	2.7
1	B	572	SER	2.6
1	B	749	TRP	2.6
1	A	549	LEU	2.6
1	A	548	PRO	2.6
1	A	545	LYS	2.5
1	B	473	VAL	2.4
1	A	750	GLY	2.4
1	B	482	ASP	2.4
1	A	748	ALA	2.3
1	B	752	LEU	2.3
1	A	475	PHE	2.2
1	A	493	PHE	2.2
1	A	472	ALA	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	467	GLU	2.1
1	A	501	THR	2.1
1	B	470	LEU	2.1
1	B	543	VAL	2.0
1	B	571	HIS	2.0
1	B	493	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NO3	B	801	4/4	0.85	0.18	2.99	39,44,44,50	0
2	NO3	A	802	4/4	0.80	0.18	1.68	55,61,61,63	0
2	NO3	A	801	4/4	0.99	0.15	0.26	23,26,27,35	0
2	NO3	B	802	4/4	0.93	0.11	-0.09	62,64,65,67	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.