



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:04 AM GMT

PDB ID : 3H3R  
Title : Crystal structure of the CERT START domain in complex with HPA-14  
Authors : Kudo, N.; Wakatsuki, S.; Kato, R.  
Deposited on : 2009-04-17  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	<div> <div>4%</div> <div>75%</div> <div>16%</div> <div>7%</div> </div>
1	B	255	<div> <div>3%</div> <div>81%</div> <div>10%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	14H	A	600	-	-	-	X
2	14H	B	600	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Goodpasture antigen binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	236	Total	C	N	O	S	0	1	0
			1901	1205	334	352	10			
1	B	235	Total	C	N	O	S	0	0	0
			1889	1198	333	349	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	344	SER	-	EXPRESSION TAG	UNP A8K7S2
A	345	ASN	-	EXPRESSION TAG	UNP A8K7S2
A	346	SER	-	EXPRESSION TAG	UNP A8K7S2
B	344	SER	-	EXPRESSION TAG	UNP A8K7S2
B	345	ASN	-	EXPRESSION TAG	UNP A8K7S2
B	346	SER	-	EXPRESSION TAG	UNP A8K7S2

- Molecule 2 is N-[(1R,3R)-3-HYDROXY-1-(HYDROXYMETHYL)-3-PHENYLPROPYL]TETRADECANAMIDE (three-letter code: 14H) (formula: C<sub>24</sub>H<sub>41</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			28	24	1	3		
2	B	1	Total	C	N	O	0	0
			28	24	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	85	Total	O	0	0
			85	85		
3	B	85	Total	O	0	0
			85	85		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Goodpasture antigen binding protein



- Molecule 1: Goodpasture antigen binding protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.42Å 74.72Å 77.71Å 90.00° 102.86° 90.00°	Depositor
Resolution (Å)	20.00 – 1.85 19.93 – 1.85	Depositor EDS
% Data completeness (in resolution range)	91.5 (20.00-1.85) 91.5 (19.93-1.85)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.90 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.4.0069	Depositor
R, $R_{free}$	0.203 , 0.253 0.202 , 0.252	Depositor DCC
$R_{free}$ test set	1848 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.1	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 51.6	EDS
Estimated twinning fraction	0.007 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 36941 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4016	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.21 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.3541e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 14H

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.70	1/1949 (0.1%)	0.73	1/2652 (0.0%)
1	B	0.71	0/1933	0.74	1/2632 (0.0%)
All	All	0.71	1/3882 (0.0%)	0.74	2/5284 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	503	CYS	CB-SG	-5.07	1.73	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	471	ARG	NE-CZ-NH1	-5.56	117.52	120.30
1	A	471	ARG	NE-CZ-NH1	-5.30	117.65	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1901	0	1875	34	0
1	B	1889	0	1863	24	0
2	A	28	0	40	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	28	0	40	2	0
3	A	85	0	0	2	0
3	B	85	0	0	1	0
All	All	4016	0	3818	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:406:ARG:HH11	1:A:406:ARG:CG	1.91	0.83
1:A:533:VAL:HG12	1:A:547:ILE:HG22	1.58	0.83
1:A:406:ARG:HH11	1:A:406:ARG:HG2	1.45	0.80
1:A:446:GLU:HG2	1:A:449:ILE:HD12	1.70	0.74
1:B:543:SER:H	1:B:546:ASN:HD22	1.40	0.70
1:A:384:GLN:O	1:A:419:LYS:NZ	2.26	0.68
1:A:377:ASN:HB3	3:A:168:HOH:O	1.94	0.67
1:A:398:GLU:OE1	1:A:581:LYS:NZ	2.27	0.67
1:B:471:ARG:HH11	1:B:471:ARG:HG2	1.60	0.66
1:B:471:ARG:NH1	1:B:471:ARG:HG2	2.11	0.66
1:B:469:HIS:O	1:B:477:GLN:HG3	1.97	0.65
1:A:386:VAL:HG13	1:A:386:VAL:O	1.96	0.65
1:A:376:GLN:O	1:A:380:THR:HG23	1.97	0.64
1:B:460:ASP:HB2	3:B:161:HOH:O	1.96	0.64
1:B:543:SER:H	1:B:546:ASN:ND2	1.97	0.62
1:A:386:VAL:CG1	1:A:386:VAL:O	2.47	0.62
1:B:446:GLU:CG	1:B:449:ILE:HG12	2.31	0.61
1:A:468:THR:HG22	1:A:479:ASP:HB3	1.83	0.61
1:B:471:ARG:NH1	1:B:475:ALA:O	2.36	0.59
1:B:449:ILE:HD11	1:B:467:GLN:NE2	2.18	0.57
1:A:398:GLU:OE2	1:A:581:LYS:NZ	2.35	0.57
1:B:446:GLU:HG2	1:B:449:ILE:HG13	1.87	0.56
1:B:534:SER:O	1:B:546:ASN:HB3	2.07	0.55
1:A:406:ARG:CG	1:A:406:ARG:NH1	2.58	0.55
1:A:398:GLU:CD	1:A:581:LYS:NZ	2.61	0.54
1:B:446:GLU:HG2	1:B:449:ILE:CG1	2.37	0.54
1:B:385:ASP:C	1:B:385:ASP:OD1	2.45	0.54
1:B:491:ALA:H	1:B:498:GLU:HG2	1.72	0.53
1:B:467:GLN:HG2	1:B:480:VAL:CG1	2.39	0.52
1:B:446:GLU:CG	1:B:449:ILE:CG1	2.88	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:585:SER:O	1:B:589:GLU:HG3	2.11	0.51
2:A:600:14H:C4	2:A:600:14H:C21	2.90	0.50
1:A:474:PRO:HD3	1:A:567:VAL:HG11	1.94	0.50
1:A:409:GLU:C	1:A:409:GLU:OE2	2.50	0.50
1:A:471:ARG:HH11	1:A:471:ARG:HG3	1.77	0.49
1:A:473:TRP:CZ2	2:A:600:14H:H30	2.48	0.49
1:A:534:SER:O	1:A:546:ASN:HB3	2.12	0.49
1:B:409:GLU:HA	1:B:413:ILE:O	2.13	0.49
1:A:452:PHE:HB2	1:A:467:GLN:HB3	1.96	0.48
1:A:488:LYS:HD3	1:A:500:TRP:CZ2	2.49	0.48
2:B:600:14H:H34B	2:B:600:14H:H31A	1.54	0.47
1:A:568:LEU:HD23	1:A:568:LEU:N	2.30	0.46
1:B:567:VAL:O	1:B:571:VAL:HG13	2.15	0.46
1:A:474:PRO:HG3	1:A:567:VAL:HG11	1.99	0.45
1:B:567:VAL:O	1:B:571:VAL:CG1	2.64	0.45
1:A:376:GLN:NE2	3:A:67:HOH:O	2.50	0.44
1:A:484:SER:OG	1:A:504:ASN:ND2	2.51	0.44
1:A:406:ARG:HH11	1:A:406:ARG:HG3	1.79	0.44
1:A:414:VAL:HG21	1:A:565:ALA:CB	2.48	0.44
1:B:467:GLN:HG2	1:B:480:VAL:HG12	2.00	0.43
2:B:600:14H:C21	2:B:600:14H:C4	2.96	0.43
1:B:385:ASP:OD1	1:B:385:ASP:O	2.36	0.43
1:B:493:THR:OG1	1:B:496:ASP:HB2	2.18	0.43
1:A:383:LEU:HD23	1:A:383:LEU:HA	1.75	0.42
1:A:406:ARG:HG2	1:A:406:ARG:NH1	2.24	0.42
1:B:415:LEU:O	1:B:416:ASP:C	2.58	0.42
1:A:573:LYS:O	1:A:577:PRO:HG2	2.20	0.41
1:A:419:LYS:HE3	1:A:552:THR:HG23	2.03	0.41
1:A:471:ARG:NH1	1:A:471:ARG:HG3	2.35	0.41
1:A:383:LEU:HD21	1:A:528:ILE:HG21	2.03	0.41
1:A:444:ASP:HB3	1:A:586:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/255 (92%)	231 (98%)	4 (2%)	0	100	100
1	B	233/255 (91%)	228 (98%)	3 (1%)	2 (1%)	21	7
All	All	468/510 (92%)	459 (98%)	7 (2%)	2 (0%)	39	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	536	PRO
1	B	416	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/226 (93%)	202 (96%)	8 (4%)	40	19
1	B	208/226 (92%)	203 (98%)	5 (2%)	57	39
All	All	418/452 (92%)	405 (97%)	13 (3%)	47	28

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	406	ARG
1	A	409	GLU
1	A	418	LEU
1	A	422	HIS
1	A	540	GLN
1	A	568	LEU
1	A	579	PHE
1	A	597	LEU
1	B	422	HIS
1	B	471	ARG
1	B	476	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	534	SER
1	B	579	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	411	ASN
1	A	434	ASN
1	A	443	ASN
1	A	495	ASN
1	A	504	ASN
1	A	539	ASN
1	A	556	ASN
1	B	369	GLN
1	B	434	ASN
1	B	443	ASN
1	B	477	GLN
1	B	504	ASN
1	B	546	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	14H	A	600	-	28,28,28	0.57	0	32,32,32	2.69	8 (25%)
2	14H	B	600	-	28,28,28	0.58	1 (3%)	32,32,32	2.35	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	14H	A	600	-	-	0/27/27/27	0/1/1/1
2	14H	B	600	-	-	0/27/27/27	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	600	14H	C5-C4	-2.05	1.48	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	14H	C4-C3-C2	-10.17	99.18	114.11
2	B	600	14H	C4-C3-C2	-7.91	102.50	114.11
2	A	600	14H	C3-C2-C1	-2.80	108.02	111.75
2	A	600	14H	C22-C21-N2	-2.64	111.53	115.83
2	B	600	14H	O1-C1-C2	-2.15	106.42	112.19
2	B	600	14H	C10-C5-C4	-2.06	116.37	120.75
2	B	600	14H	O21-C21-N2	2.71	127.60	123.01
2	A	600	14H	O21-C21-N2	3.41	128.78	123.01
2	A	600	14H	O4-C4-C3	3.42	115.72	108.94
2	B	600	14H	O4-C4-C5	3.94	120.03	111.20
2	A	600	14H	C3-C4-C5	4.09	118.92	111.36
2	B	600	14H	O4-C4-C3	4.16	117.17	108.94
2	B	600	14H	C2-N2-C21	4.17	128.94	122.79
2	A	600	14H	O4-C4-C5	4.60	121.51	111.20
2	B	600	14H	C3-C4-C5	4.72	120.10	111.36
2	A	600	14H	C2-N2-C21	5.29	130.60	122.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	14H	2	0
2	B	600	14H	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	236/255 (92%)	0.10	10 (4%) 40 37	18, 29, 46, 56	0
1	B	235/255 (92%)	0.03	7 (2%) 54 51	17, 27, 41, 54	0
All	All	471/510 (92%)	0.06	17 (3%) 46 44	17, 27, 44, 56	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	537	GLU	4.7
1	A	364	THR	4.4
1	A	539	ASN	4.2
1	B	536	PRO	3.9
1	B	537	GLU	3.6
1	B	411	ASN	3.2
1	B	540	GLN	2.8
1	A	517	ARG	2.8
1	A	516	ASN	2.7
1	A	510	ASP	2.6
1	A	536	PRO	2.6
1	B	516	ASN	2.6
1	A	387	GLY	2.5
1	B	364	THR	2.4
1	A	492	LEU	2.2
1	A	540	GLN	2.1
1	B	535	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	14H	B	600	28/28	0.94	0.15	2.21	17,27,44,45	0
2	14H	A	600	28/28	0.93	0.14	2.02	19,26,35,35	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.