



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:26 AM GMT

PDB ID : 2H6F  
Title : Protein Farnesyltransferase Complexed with a Farnesylated DDPTASACVLS Peptide Product at 1.5A Resolution  
Authors : Terry, K.L.; Beese, L.S.  
Deposited on : 2006-05-31  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	382	<div> <div>7%</div> <div>76%</div> <div>7%</div> <div>18%</div> </div>
2	B	437	<div> <div>7%</div> <div>84%</div> <div>10%</div> <div>6%</div> </div>
3	P	11	<div> <div>27%</div> <div>27%</div> <div>9%</div> <div>64%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	FAR	P	2010	-	-	X	X
7	ACY	A	3001	-	-	X	-
7	ACY	A	3002	-	-	X	X
7	ACY	B	3003	-	-	X	-
7	ACY	B	3005	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein farnesyltransferase/geranylgeranyltransferase type I alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	0	0
			2683	1711	467	500	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	380	GLU	-	cloning artifact	UNP P49354
A	381	GLU	-	cloning artifact	UNP P49354
A	382	PHE	-	cloning artifact	UNP P49354


- Molecule 2 is a protein called Protein farnesyltransferase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	410	Total	C	N	O	S	0	0	0
			3228	2065	552	589	22			

- Molecule 3 is a protein called farnesylated peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	4	Total	C	N	O	S	0	0	0
			28	17	4	6	1			

- Molecule 4 is SUGAR (SUCROSE) (three-letter code: SUC) (formula: C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			23	12	11		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Zn	0	0
			1	1		

- Molecule 6 is FARNESYL (three-letter code: FAR) (formula: C<sub>15</sub>H<sub>26</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	P	1	Total C 15 15	0	0

- Molecule 7 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	330	Total O 330 330	0	0
8	B	381	Total O 381 381	0	0
8	P	8	Total O 8 8	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein farnesyltransferase/geranylgeranyltransferase type I alpha subunit



- Molecule 2: Protein farnesyltransferase beta subunit



- Molecule 3: farnesylated peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	178.57Å 178.57Å 64.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.66 – 1.50 38.66 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.7 (38.66-1.50) 98.9 (38.66-1.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 1.50Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.190 , 0.203 0.189 , 0.202	Depositor DCC
$R_{free}$ test set	9297 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.7	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 45.9	EDS
Estimated twinning fraction	0.019 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 184691 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6717	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAR, ZN, SUC, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.26	0/2750	0.49	0/3735
2	B	0.29	0/3317	0.56	0/4508
3	P	0.57	0/27	1.15	1/34 (2.9%)
All	All	0.28	0/6094	0.54	1/8277 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	P	2007	VAL	N-CA-CB	5.67	123.97	111.50

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2683	0	2601	18	0
2	B	3228	0	3147	25	0
3	P	28	0	28	0	0
4	B	23	0	22	0	0
5	B	1	0	0	0	0
6	P	15	0	24	10	0
7	A	12	0	9	15	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	8	0	6	2	0
8	A	330	0	0	6	0
8	B	381	0	0	4	0
8	P	8	0	0	3	0
All	All	6717	0	5837	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:3002:ACY:H2	8:P:1691:HOH:O	1.11	1.29
7:B:3003:ACY:H1	8:B:1551:HOH:O	1.10	1.27
7:B:3003:ACY:CH3	8:B:1551:HOH:O	1.75	0.93
7:A:3002:ACY:H3	6:P:2010:FAR:H102	1.50	0.91
2:B:627:CYS:HB3	2:B:671:ILE:HD11	1.52	0.88
7:A:3002:ACY:H3	6:P:2010:FAR:C10	2.04	0.87
7:A:3002:ACY:CH3	8:P:1691:HOH:O	1.88	0.80
2:B:845:CYS:HB2	2:B:851:LEU:HD21	1.66	0.76
7:A:3001:ACY:H3	6:P:2010:FAR:H42	1.69	0.74
1:A:316:VAL:O	1:A:320:GLU:HG3	1.89	0.72
1:A:221:GLN:HG3	8:A:1180:HOH:O	1.90	0.72
7:A:3002:ACY:CH3	6:P:2010:FAR:C10	2.69	0.71
2:B:766:ARG:HH21	2:B:818:GLN:HG2	1.55	0.70
7:A:3002:ACY:CH3	6:P:2010:FAR:H103	2.26	0.66
7:A:3002:ACY:O	6:P:2010:FAR:C4	2.47	0.63
2:B:881:MET:C	2:B:882:LEU:HD12	2.20	0.62
2:B:577:LYS:HD3	2:B:846:PRO:O	2.00	0.62
7:A:3002:ACY:O	6:P:2010:FAR:H41	1.99	0.61
7:A:3001:ACY:H3	6:P:2010:FAR:C4	2.30	0.61
1:A:298:GLN:O	1:A:302:LEU:HD23	2.03	0.59
1:A:303:GLN:N	1:A:304:PRO:HD2	2.20	0.56
2:B:739:ILE:HB	2:B:752:THR:HA	1.87	0.56
2:B:801:SER:O	2:B:805:ALA:HB3	2.05	0.56
7:A:3001:ACY:H1	8:A:1438:HOH:O	2.06	0.55
1:A:304:PRO:HG2	8:A:1998:HOH:O	2.10	0.52
1:A:342:GLU:CD	1:A:357:ARG:HH12	2.13	0.51
2:B:696:GLY:HA2	8:B:1901:HOH:O	2.10	0.51
2:B:763:LYS:HB2	2:B:763:LYS:NZ	2.26	0.51
1:A:335:ASN:O	1:A:339:GLU:HG3	2.12	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:LEU:HD11	1:A:126:LEU:HG	1.94	0.50
2:B:692:LEU:HD23	2:B:699:VAL:CG2	2.42	0.49
7:A:3002:ACY:CH3	6:P:2010:FAR:H102	2.30	0.48
2:B:806:GLY:HA2	2:B:872:ILE:HD13	1.96	0.48
2:B:526:LEU:HD13	2:B:563:LYS:HB2	1.95	0.47
2:B:635:SER:OG	2:B:637:GLU:HG2	2.15	0.47
1:A:354:GLU:HB2	8:A:1697:HOH:O	2.14	0.47
2:B:753:PHE:HA	2:B:807:LEU:HD21	1.97	0.47
7:A:3001:ACY:OXT	8:A:1360:HOH:O	2.19	0.47
2:B:624:THR:O	2:B:628:GLN:HG3	2.14	0.47
2:B:908:VAL:O	2:B:912:THR:HG23	2.14	0.46
2:B:627:CYS:CB	2:B:671:ILE:HD11	2.34	0.46
1:A:96:PHE:HA	1:A:126:LEU:HD13	1.99	0.45
2:B:687:PRO:HG2	8:B:1434:HOH:O	2.16	0.45
2:B:718:THR:HB	2:B:719:PRO:HD2	1.99	0.44
7:A:3002:ACY:C	8:P:1691:HOH:O	2.46	0.44
1:A:109:ARG:HG3	8:A:1256:HOH:O	2.16	0.44
2:B:651:ALA:HB3	2:B:652:PRO:CD	2.47	0.44
7:A:3002:ACY:O	6:P:2010:FAR:H42	2.15	0.44
1:A:109:ARG:HH11	1:A:109:ARG:CB	2.32	0.43
1:A:84:PRO:C	1:A:86:PRO:HD3	2.39	0.43
2:B:797:ASP:HB3	2:B:800:TYR:HD1	1.84	0.42
1:A:353:LYS:O	1:A:357:ARG:HG2	2.19	0.42
1:A:58:LEU:HD23	1:A:63:TYR:CZ	2.54	0.42
1:A:264:LEU:O	1:A:268:LYS:HG3	2.19	0.42
2:B:870:LEU:O	2:B:874:GLN:HG3	2.20	0.41
2:B:576:GLU:HA	2:B:576:GLU:OE1	2.20	0.41
1:A:58:LEU:HB2	1:A:125:GLU:OE1	2.21	0.41
2:B:850:LEU:HB2	2:B:863:THR:HA	2.02	0.40
1:A:189:ILE:HD11	1:A:205:HIS:HD2	1.86	0.40
2:B:806:GLY:O	2:B:809:PRO:HD2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/382 (82%)	301 (96%)	12 (4%)	0	100	100
2	B	408/437 (93%)	402 (98%)	6 (2%)	0	100	100
3	P	2/11 (18%)	2 (100%)	0	0	100	100
All	All	723/830 (87%)	705 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/344 (86%)	294 (100%)	0	100	100
2	B	346/370 (94%)	345 (100%)	1 (0%)	94	86
3	P	4/9 (44%)	4 (100%)	0	100	100
All	All	644/723 (89%)	643 (100%)	1 (0%)	95	87

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	851	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	818	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	ACY	A	3001	-	1,3,3	2.43	1 (100%)	0,3,3	0.00	-
7	ACY	A	3002	-	1,3,3	2.39	1 (100%)	0,3,3	0.00	-
7	ACY	A	3004	-	1,3,3	2.53	1 (100%)	0,3,3	0.00	-
7	ACY	B	3003	-	1,3,3	2.41	1 (100%)	0,3,3	0.00	-
7	ACY	B	3005	-	1,3,3	2.45	1 (100%)	0,3,3	0.00	-
4	SUC	B	3010	-	24,24,24	1.44	2 (8%)	36,36,36	0.88	1 (2%)
6	FAR	P	2010	3	14,14,14	0.71	0	16,16,16	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ACY	A	3001	-	-	0/0/0/0	0/0/0/0
7	ACY	A	3002	-	-	0/0/0/0	0/0/0/0
7	ACY	A	3004	-	-	0/0/0/0	0/0/0/0
7	ACY	B	3003	-	-	0/0/0/0	0/0/0/0
7	ACY	B	3005	-	-	0/0/0/0	0/0/0/0
4	SUC	B	3010	-	-	0/12/51/51	0/2/2/2
6	FAR	P	2010	3	-	0/14/14/14	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	3010	SUC	C4-C5	2.36	1.58	1.53
7	A	3002	ACY	CH3-C	2.39	1.52	1.48
7	B	3003	ACY	CH3-C	2.41	1.52	1.48
7	A	3001	ACY	CH3-C	2.43	1.52	1.48
7	B	3005	ACY	CH3-C	2.45	1.52	1.48
7	A	3004	ACY	CH3-C	2.53	1.52	1.48
4	B	3010	SUC	C3-C2	5.10	1.65	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	3010	SUC	O3-C3-C2	-2.07	105.68	110.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	3001	ACY	4	0
7	A	3002	ACY	11	0
7	B	3003	ACY	2	0
6	P	2010	FAR	10	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/382 (82%)	0.76	27 (8%) 13 12	11, 19, 33, 48	0
2	B	410/437 (93%)	0.58	29 (7%) 19 19	9, 14, 24, 34	0
3	P	4/11 (36%)	3.16	3 (75%) 0 0	13, 16, 22, 25	0
All	All	729/830 (87%)	0.67	59 (8%) 15 14	9, 16, 28, 48	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	55	PHE	14.9
1	A	369	THR	6.6
1	A	304	PRO	6.2
2	B	515	SER	5.9
1	A	302	LEU	5.8
3	P	2006	CYS	5.4
2	B	880	ALA	5.2
1	A	306	HIS	5.2
1	A	307	SER	5.2
2	B	879	GLY	5.1
1	A	301	ASP	4.8
1	A	328	ASP	4.6
1	A	305	SER	4.5
1	A	368	SER	4.0
2	B	881	MET	3.9
1	A	303	GLN	3.8
2	B	923	PHE	3.6
2	B	924	GLU	3.4
1	A	109	ARG	3.4
2	B	878	SER	3.3
1	A	145	GLN	3.2
1	A	300	LEU	3.2
2	B	825	MET	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	P	2008	LEU	3.1
1	A	326	GLN	3.1
3	P	2007	VAL	3.0
2	B	620	GLN	3.0
1	A	329	ASN	3.0
2	B	882	LEU	2.9
1	A	56	VAL	2.9
1	A	91	ILE	2.8
1	A	327	CYS	2.7
2	B	766	ARG	2.7
2	B	588	GLN	2.7
2	B	813	ARG	2.6
2	B	846	PRO	2.6
2	B	917	GLN	2.6
1	A	85	ASN	2.5
2	B	532	ARG	2.5
2	B	777	VAL	2.5
1	A	59	ASP	2.5
2	B	618	ILE	2.4
2	B	803	TRP	2.4
2	B	621	ILE	2.4
2	B	584	ARG	2.3
2	B	617	PRO	2.3
1	A	81	ASN	2.3
1	A	84	PRO	2.3
2	B	719	PRO	2.3
2	B	516	SER	2.2
1	A	150	GLU	2.2
2	B	753	PHE	2.2
2	B	580	HIS	2.1
2	B	819	GLY	2.1
1	A	268	LYS	2.1
2	B	747	ALA	2.1
2	B	628	GLN	2.1
1	A	297	ASN	2.1
1	A	58	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	ACY	A	3002	4/4	0.82	0.24	4.33	40,41,41,41	0
7	ACY	B	3005	4/4	0.65	0.25	2.36	37,37,38,38	0
6	FAR	P	2010	15/15	0.88	0.22	2.11	16,19,24,24	0
4	SUC	B	3010	23/23	0.91	0.15	1.69	17,20,21,21	0
7	ACY	A	3004	4/4	0.97	0.09	-0.89	15,15,15,15	0
5	ZN	B	1001	1/1	1.00	0.04	-1.57	13,13,13,13	0
7	ACY	B	3003	4/4	0.71	0.23	-	40,41,41,41	0
7	ACY	A	3001	4/4	0.71	0.24	-	40,41,41,41	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.