



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 02:29 AM GMT

PDB ID : 2HFW
Title : Structural and kinetic analysis of proton shuttle residues in the active site of human carbonic anhydrase III
Authors : Elder, I.; Fisher, S.Z.; Laipis, P.J.; Tu, C.K.; McKenna, R.; Silverman, D.N.
Deposited on : 2006-06-26
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 2101 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	257	2065	1317	361	381	6	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	64	HIS	LYS	ENGINEERED	UNP P07451
A	67	ASN	ARG	ENGINEERED	UNP P07451
A	182	SER	CYS	ENGINEERED	UNP P07451
A	188	SER	CYS	ENGINEERED	UNP P07451

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	35	Total O 35 35	0	0

3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Carbonic anhydrase 3



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.37 Å 51.67 Å 117.66 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 19.61 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.50) 86.8 (19.61-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ ¹	1.58 (at 2.50 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R , R_{free}	0.204 , 0.239 0.212 , 0.238	Depositor DCC
R_{free} test set	448 reflections (5.49%)	DCC
Wilson B-factor (Å ²)	39.9	Xtriage
Anisotropy	0.419	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 40.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 8508 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2101	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	2.18	63/2134 (3.0%)	1.18	14/2905 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (63) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	191	TYR	CD1-CE1	-9.80	1.24	1.39
1	A	191	TYR	CE2-CZ	-9.57	1.26	1.38
1	A	218	VAL	CB-CG2	-7.86	1.36	1.52
1	A	117	GLU	CD-OE1	-7.72	1.17	1.25
1	A	206	CYS	CB-SG	-7.66	1.69	1.82
1	A	191	TYR	CE1-CZ	-7.57	1.28	1.38
1	A	114	TYR	CE1-CZ	-7.56	1.28	1.38
1	A	47	TRP	CE3-CZ3	-6.77	1.26	1.38
1	A	205	GLU	CD-OE2	-6.76	1.18	1.25
1	A	196	GLY	C-O	-6.76	1.12	1.23
1	A	192	TRP	CE3-CZ3	-6.75	1.26	1.38
1	A	187	ALA	CA-CB	-6.73	1.38	1.52
1	A	194	TYR	CE2-CZ	-6.71	1.29	1.38
1	A	106	GLU	CD-OE2	-6.64	1.18	1.25
1	A	69	VAL	CB-CG1	-6.63	1.39	1.52
1	A	205	GLU	CD-OE1	-6.52	1.18	1.25
1	A	117	GLU	CD-OE2	-6.24	1.18	1.25
1	A	121	VAL	CB-CG2	-6.23	1.39	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	97	TRP	CG-CD1	-6.23	1.28	1.36
1	A	191	TYR	CZ-OH	-6.21	1.27	1.37
1	A	121	VAL	CB-CG1	-6.09	1.40	1.52
1	A	47	TRP	CG-CD1	-6.08	1.28	1.36
1	A	93	PHE	CD1-CE1	-6.08	1.27	1.39
1	A	143	VAL	CB-CG1	-6.08	1.40	1.52
1	A	256	VAL	CB-CG2	-6.08	1.40	1.52
1	A	191	TYR	CD2-CE2	-6.01	1.30	1.39
1	A	140	GLY	C-O	-5.99	1.14	1.23
1	A	194	TYR	CD1-CE1	-5.97	1.30	1.39
1	A	160	PHE	CD1-CE1	-5.94	1.27	1.39
1	A	93	PHE	CD2-CE2	-5.82	1.27	1.39
1	A	147	PHE	CD2-CE2	-5.80	1.27	1.39
1	A	260	PHE	CE2-CZ	-5.79	1.26	1.37
1	A	204	GLU	CD-OE1	-5.78	1.19	1.25
1	A	208	VAL	CB-CG2	-5.76	1.40	1.52
1	A	185	PHE	CD1-CE1	-5.75	1.27	1.39
1	A	68	VAL	CB-CG1	-5.74	1.40	1.52
1	A	74	TYR	CD2-CE2	-5.72	1.30	1.39
1	A	114	TYR	CD2-CE2	-5.71	1.30	1.39
1	A	197	SER	CB-OG	-5.66	1.34	1.42
1	A	123	TRP	CG-CD1	-5.66	1.28	1.36
1	A	163	ALA	CA-CB	-5.65	1.40	1.52
1	A	7	TYR	CD1-CE1	-5.62	1.30	1.39
1	A	211	LEU	CG-CD2	-5.54	1.31	1.51
1	A	185	PHE	CD2-CE2	-5.53	1.28	1.39
1	A	176	PHE	CE1-CZ	-5.53	1.26	1.37
1	A	180	ASP	CB-CG	-5.49	1.40	1.51
1	A	25	GLY	C-O	-5.47	1.14	1.23
1	A	109	VAL	CB-CG1	-5.44	1.41	1.52
1	A	198	PHE	CE1-CZ	-5.40	1.27	1.37
1	A	106	GLU	CD-OE1	-5.38	1.19	1.25
1	A	185	PHE	CE2-CZ	-5.34	1.27	1.37
1	A	122	HIS	C-O	-5.30	1.13	1.23
1	A	176	PHE	CD2-CE2	-5.27	1.28	1.39
1	A	49	VAL	CB-CG1	-5.18	1.42	1.52
1	A	198	PHE	CD1-CE1	-5.14	1.28	1.39
1	A	191	TYR	CG-CD1	-5.11	1.32	1.39
1	A	192	TRP	CG-CD1	-5.10	1.29	1.36
1	A	88	TYR	CE2-CZ	-5.10	1.31	1.38
1	A	114	TYR	CG-CD2	-5.06	1.32	1.39
1	A	201	PRO	CB-CG	-5.05	1.24	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	160	PHE	CD2-CE2	-5.04	1.29	1.39
1	A	209	TRP	CG-CD1	-5.03	1.29	1.36
1	A	123	TRP	CE3-CZ3	-5.02	1.29	1.38

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	52	ASP	CB-CG-OD2	10.54	127.79	118.30
1	A	101	ASP	CB-CG-OD2	7.68	125.21	118.30
1	A	221	ASP	CB-CG-OD2	7.31	124.88	118.30
1	A	130	THR	N-CA-CB	7.25	124.08	110.30
1	A	84	LEU	C-N-CD	-6.63	106.01	120.60
1	A	190	ASP	CB-CG-OD2	6.57	124.21	118.30
1	A	206	CYS	CA-CB-SG	5.55	124.00	114.00
1	A	210	LEU	CB-CG-CD2	-5.49	101.67	111.00
1	A	164	LEU	CB-CG-CD1	-5.41	101.81	111.00
1	A	37	ASP	CB-CG-OD2	5.39	123.16	118.30
1	A	165	ASP	CB-CG-OD2	5.29	123.06	118.30
1	A	102	ASP	CB-CG-OD2	5.23	123.01	118.30
1	A	211	LEU	CA-CB-CG	5.17	127.19	115.30
1	A	155	GLY	N-CA-C	5.09	125.82	113.10

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	13	PRO	Peptide
1	A	49	VAL	Peptide
1	A	85	PRO	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2065	0	1974	108	1
2	A	1	0	0	0	0
3	A	35	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2101	0	1974	108	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:TRP:CH2	1:A:64:HIS:HD2	1.55	1.24
1:A:238:PRO:O	1:A:239:VAL:HG23	1.41	1.17
1:A:5:TRP:CH2	1:A:64:HIS:CD2	2.34	1.16
1:A:6:GLY:HA3	1:A:11:ASN:ND2	1.69	1.07
1:A:166:LYS:O	1:A:172:LYS:HD3	1.61	1.01
1:A:36:LYS:H	1:A:36:LYS:HD3	1.25	1.01
1:A:36:LYS:CD	1:A:36:LYS:H	1.76	0.97
1:A:17:HIS:HA	1:A:20:PHE:O	1.69	0.92
1:A:35:THR:OG1	1:A:110:ASP:OD1	1.86	0.92
1:A:64:HIS:O	1:A:241:LEU:HD21	1.74	0.88
1:A:5:TRP:CZ2	1:A:64:HIS:CD2	2.65	0.84
1:A:238:PRO:O	1:A:239:VAL:CG2	2.24	0.83
1:A:228:SER:O	1:A:230:LEU:HD23	1.79	0.82
1:A:228:SER:O	1:A:229:LEU:O	2.01	0.78
1:A:36:LYS:CD	1:A:36:LYS:N	2.49	0.75
1:A:6:GLY:HA3	1:A:11:ASN:HD22	1.51	0.74
1:A:169:THR:CG2	1:A:170:LYS:O	2.38	0.72
1:A:166:LYS:NZ	3:A:328:HOH:O	2.23	0.71
1:A:150:ILE:CG2	1:A:151:GLY:N	2.54	0.68
1:A:92:GLN:OE1	1:A:94:HIS:HD2	1.75	0.68
1:A:228:SER:O	1:A:229:LEU:C	2.31	0.67
1:A:61:ASN:C	1:A:61:ASN:OD1	2.30	0.67
1:A:24:LYS:HG3	1:A:24:LYS:O	1.94	0.66
1:A:169:THR:HG22	1:A:170:LYS:O	1.97	0.65
1:A:22:ASN:O	1:A:24:LYS:N	2.30	0.65
1:A:35:THR:O	3:A:403:HOH:O	2.16	0.64
1:A:64:HIS:ND1	1:A:64:HIS:N	2.45	0.63
1:A:129:ASN:O	1:A:130:THR:HG22	1.98	0.63
1:A:129:ASN:O	1:A:130:THR:CG2	2.46	0.63
1:A:167:ILE:O	1:A:169:THR:N	2.31	0.62
1:A:219:SER:OG	1:A:222:GLN:HG3	2.00	0.62
1:A:228:SER:C	1:A:229:LEU:O	2.34	0.62
1:A:167:ILE:O	1:A:168:LYS:C	2.37	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:ASN:C	1:A:24:LYS:H	2.04	0.61
1:A:15:HIS:ND1	1:A:18:GLU:OE2	2.35	0.60
1:A:156:GLU:N	1:A:156:GLU:OE2	2.35	0.59
1:A:5:TRP:CZ3	1:A:64:HIS:CD2	2.89	0.58
1:A:92:GLN:OE1	1:A:94:HIS:CD2	2.57	0.57
1:A:51:TYR:HA	1:A:77:SER:HB3	1.87	0.57
1:A:169:THR:HG23	1:A:170:LYS:O	2.04	0.56
1:A:161:LEU:O	1:A:225:LYS:HE2	2.05	0.56
1:A:150:ILE:HG23	1:A:151:GLY:N	2.20	0.56
1:A:63:GLY:HA3	1:A:170:LYS:HE2	1.87	0.55
1:A:35:THR:HG22	3:A:332:HOH:O	2.07	0.54
1:A:249:GLN:HB3	1:A:250:PRO:HD2	1.88	0.54
1:A:69:VAL:HG12	1:A:70:PHE:N	2.22	0.54
1:A:225:LYS:O	1:A:228:SER:OG	2.25	0.53
1:A:30:PRO:HG3	1:A:106:GLU:HB3	1.90	0.52
1:A:234:GLU:C	1:A:236:GLU:H	2.14	0.52
1:A:138:ARG:NH2	1:A:195:GLN:OE1	2.38	0.52
1:A:89:ARG:O	1:A:122:HIS:HA	2.10	0.52
1:A:180:ASP:O	1:A:183:SER:OG	2.26	0.51
1:A:22:ASN:C	1:A:24:LYS:N	2.64	0.51
1:A:232:SER:O	1:A:233:ALA:O	2.30	0.50
1:A:237:PRO:O	1:A:238:PRO:O	2.30	0.50
1:A:169:THR:HG23	1:A:233:ALA:HA	1.94	0.49
1:A:65:THR:HG23	1:A:67:ASN:OD1	2.12	0.49
1:A:218:VAL:HG22	1:A:219:SER:N	2.28	0.49
1:A:167:ILE:HA	1:A:172:LYS:HB3	1.95	0.48
1:A:69:VAL:CG1	1:A:70:PHE:N	2.76	0.48
1:A:156:GLU:HG2	1:A:183:SER:HB2	1.94	0.48
1:A:5:TRP:HH2	1:A:64:HIS:HD2	1.45	0.47
1:A:98:GLY:O	1:A:227:ARG:HD2	2.14	0.47
1:A:234:GLU:O	1:A:236:GLU:N	2.48	0.47
1:A:228:SER:O	1:A:230:LEU:CD2	2.58	0.47
1:A:166:LYS:O	1:A:172:LYS:CD	2.49	0.47
1:A:53:GLY:O	1:A:178:LYS:HA	2.15	0.47
1:A:5:TRP:CZ2	1:A:64:HIS:NE2	2.83	0.47
1:A:129:ASN:C	1:A:130:THR:CG2	2.84	0.46
1:A:94:HIS:O	1:A:95:LEU:HG	2.16	0.45
1:A:236:GLU:HB3	1:A:237:PRO:HD2	1.98	0.45
1:A:29:SER:HB3	1:A:30:PRO:HA	1.98	0.45
1:A:237:PRO:O	1:A:238:PRO:C	2.54	0.45
1:A:61:ASN:OD1	1:A:61:ASN:O	2.34	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:LEU:HA	1:A:85:PRO:HD3	1.54	0.45
1:A:129:ASN:C	1:A:130:THR:HG23	2.36	0.45
1:A:46:PRO:HB3	3:A:348:HOH:O	2.17	0.45
1:A:150:ILE:HD12	1:A:150:ILE:HA	1.58	0.45
1:A:24:LYS:HE3	1:A:24:LYS:O	2.16	0.44
1:A:95:LEU:HD11	1:A:118:LEU:HD13	2.00	0.44
1:A:246:ARG:HA	1:A:247:PRO:HD3	1.58	0.44
1:A:180:ASP:C	1:A:180:ASP:OD1	2.55	0.44
1:A:149:LYS:HG3	1:A:149:LYS:HZ3	1.53	0.44
1:A:150:ILE:N	1:A:150:ILE:HD13	2.33	0.44
1:A:180:ASP:HA	1:A:181:PRO:HD3	1.77	0.43
1:A:72:ASP:OD2	1:A:72:ASP:N	2.51	0.43
1:A:156:GLU:HA	1:A:159:ILE:HD12	2.00	0.43
1:A:20:PHE:HA	1:A:21:PRO:HD3	1.64	0.43
1:A:46:PRO:HA	3:A:348:HOH:O	2.18	0.43
1:A:237:PRO:HA	1:A:238:PRO:HD2	1.47	0.42
1:A:11:ASN:O	1:A:12:GLY:O	2.36	0.42
1:A:11:ASN:N	1:A:11:ASN:OD1	2.52	0.42
1:A:59:ILE:HG22	1:A:176:PHE:HB2	2.01	0.42
1:A:198:PHE:CE1	1:A:206:CYS:SG	3.12	0.42
1:A:41:ASP:OD2	1:A:41:ASP:C	2.55	0.42
1:A:14:ASP:N	1:A:14:ASP:OD1	2.33	0.42
1:A:234:GLU:C	1:A:236:GLU:N	2.73	0.42
1:A:94:HIS:O	1:A:118:LEU:HD12	2.18	0.42
1:A:151:GLY:N	1:A:218:VAL:O	2.53	0.42
1:A:95:LEU:HD11	1:A:118:LEU:CD1	2.50	0.41
1:A:36:LYS:HD2	1:A:36:LYS:N	2.33	0.41
1:A:120:LEU:HB3	1:A:122:HIS:HE1	1.85	0.41
1:A:12:GLY:HA2	1:A:13:PRO:HD3	1.78	0.41
1:A:38:ILE:HG21	1:A:38:ILE:HD13	1.79	0.41
1:A:247:PRO:HA	1:A:248:PRO:HD3	1.78	0.41
1:A:64:HIS:C	1:A:241:LEU:HD21	2.38	0.41
1:A:84:LEU:HB3	1:A:85:PRO:HD2	2.01	0.41
1:A:65:THR:CG2	1:A:67:ASN:OD1	2.69	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:GLU:OE1	1:A:113:LYS:NZ[4_456]	1.80	0.40

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	255/260 (98%)	218 (86%)	27 (11%)	10 (4%)	4 5

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	14	ASP
1	A	168	LYS
1	A	229	LEU
1	A	233	ALA
1	A	238	PRO
1	A	12	GLY
1	A	23	ALA
1	A	165	ASP
1	A	9	SER
1	A	21	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	226/228 (99%)	204 (90%)	22 (10%)	10 19

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	33	LEU
1	A	35	THR
1	A	36	LYS
1	A	58	THR
1	A	60	LEU
1	A	64	HIS
1	A	77	SER
1	A	80	ARG
1	A	99	SER
1	A	100	SER
1	A	149	LYS
1	A	150	ILE
1	A	152	HIS
1	A	169	THR
1	A	172	LYS
1	A	189	ARG
1	A	228	SER
1	A	232	SER
1	A	235	ASN
1	A	242	VAL
1	A	250	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	HIS
1	A	94	HIS
1	A	253	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/260 (98%)	-0.10	6 (2%) 64 67	16, 35, 63, 84	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	168	LYS	4.1
1	A	235	ASN	3.8
1	A	9	SER	3.3
1	A	237	PRO	2.9
1	A	167	ILE	2.3
1	A	238	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no carbohydrates in this entry.

6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ZN	A	262	1/1	0.99	0.09	-	26,26,26,26	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.