



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:32 AM GMT

PDB ID : 2HII
Title : heterotrimeric PCNA sliding clamp
Authors : Pascal, J.M.; Tsodikov, O.V.; Ellenberger, T.
Deposited on : 2006-06-29
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	<div> <div>67%</div> <div>25%</div> <div>5%</div> <div>.</div> </div>
1	X	257	<div> <div>63%</div> <div>27%</div> <div>7%</div> <div>.</div> </div>
2	B	245	<div> <div>61%</div> <div>31%</div> <div>7%</div> <div>.</div> </div>
2	Y	245	<div> <div>62%</div> <div>29%</div> <div>7%</div> <div>.</div> </div>
3	C	252	<div> <div>66%</div> <div>24%</div> <div>6%</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Z	252	<div><div><div>%</div><div><div></div></div><div>64%</div><div>27%</div><div>5%</div><div>•</div></div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PCNA1 (SSO0397).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	Se	0	0	0
			1927	1228	309	381	9			
1	X	249	Total	C	N	O	Se	0	0	0
			1927	1228	309	381	9			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	47	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	50	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	157	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	182	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	204	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	220	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	229	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	241	MSE	MET	MODIFIED RESIDUE	UNP P57766
A	250	LEU	-	CLONING ARTIFACT	UNP P57766
A	251	GLU	-	CLONING ARTIFACT	UNP P57766
A	252	HIS	-	EXPRESSION TAG	UNP P57766
A	253	HIS	-	EXPRESSION TAG	UNP P57766
A	254	HIS	-	EXPRESSION TAG	UNP P57766
A	255	HIS	-	EXPRESSION TAG	UNP P57766
A	256	HIS	-	EXPRESSION TAG	UNP P57766
A	257	HIS	-	EXPRESSION TAG	UNP P57766
X	1	MSE	MET	MODIFIED RESIDUE	UNP P57766
X	47	MSE	MET	MODIFIED RESIDUE	UNP P57766
X	50	MSE	MET	MODIFIED RESIDUE	UNP P57766
X	157	MSE	MET	MODIFIED RESIDUE	UNP P57766
X	182	MSE	MET	MODIFIED RESIDUE	UNP P57766
X	204	MSE	MET	MODIFIED RESIDUE	UNP P57766
X	220	MSE	MET	MODIFIED RESIDUE	UNP P57766
X	229	MSE	MET	MODIFIED RESIDUE	UNP P57766

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
X	241	MSE	MET	MODIFIED RESIDUE	UNP P57766
X	250	LEU	-	CLONING ARTIFACT	UNP P57766
X	251	GLU	-	CLONING ARTIFACT	UNP P57766
X	252	HIS	-	EXPRESSION TAG	UNP P57766
X	253	HIS	-	EXPRESSION TAG	UNP P57766
X	254	HIS	-	EXPRESSION TAG	UNP P57766
X	255	HIS	-	EXPRESSION TAG	UNP P57766
X	256	HIS	-	EXPRESSION TAG	UNP P57766
X	257	HIS	-	EXPRESSION TAG	UNP P57766

- Molecule 2 is a protein called PCNA2 (SSO1047).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	243	Total	C	N	O	Se	0	0	0
			1922	1237	301	380	4			
2	Y	243	Total	C	N	O	Se	0	0	0
			1922	1237	301	380	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	MSE	MET	MODIFIED RESIDUE	UNP Q97Z84
B	199	MSE	MET	MODIFIED RESIDUE	UNP Q97Z84
B	208	MSE	MET	MODIFIED RESIDUE	UNP Q97Z84
B	215	MSE	MET	MODIFIED RESIDUE	UNP Q97Z84
Y	2	MSE	MET	MODIFIED RESIDUE	UNP Q97Z84
Y	199	MSE	MET	MODIFIED RESIDUE	UNP Q97Z84
Y	208	MSE	MET	MODIFIED RESIDUE	UNP Q97Z84
Y	215	MSE	MET	MODIFIED RESIDUE	UNP Q97Z84

- Molecule 3 is a protein called PCNA3 (SSO0405).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	243	Total	C	N	O	Se	0	0	0
			1926	1224	311	387	4			
3	Z	243	Total	C	N	O	Se	0	0	0
			1926	1224	311	387	4			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	1	MSE	MET	MODIFIED RESIDUE	UNP P57765

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	55	MSE	MET	MODIFIED RESIDUE	UNP P57765
C	75	MSE	MET	MODIFIED RESIDUE	UNP P57765
C	230	MSE	MET	MODIFIED RESIDUE	UNP P57765
C	245	LEU	-	CLONING ARTIFACT	UNP P57765
C	246	GLU	-	CLONING ARTIFACT	UNP P57765
C	247	HIS	-	EXPRESSION TAG	UNP P57765
C	248	HIS	-	EXPRESSION TAG	UNP P57765
C	249	HIS	-	EXPRESSION TAG	UNP P57765
C	250	HIS	-	EXPRESSION TAG	UNP P57765
C	251	HIS	-	EXPRESSION TAG	UNP P57765
C	252	HIS	-	EXPRESSION TAG	UNP P57765
Z	1	MSE	MET	MODIFIED RESIDUE	UNP P57765
Z	55	MSE	MET	MODIFIED RESIDUE	UNP P57765
Z	75	MSE	MET	MODIFIED RESIDUE	UNP P57765
Z	230	MSE	MET	MODIFIED RESIDUE	UNP P57765
Z	245	LEU	-	CLONING ARTIFACT	UNP P57765
Z	246	GLU	-	CLONING ARTIFACT	UNP P57765
Z	247	HIS	-	EXPRESSION TAG	UNP P57765
Z	248	HIS	-	EXPRESSION TAG	UNP P57765
Z	249	HIS	-	EXPRESSION TAG	UNP P57765
Z	250	HIS	-	EXPRESSION TAG	UNP P57765
Z	251	HIS	-	EXPRESSION TAG	UNP P57765
Z	252	HIS	-	EXPRESSION TAG	UNP P57765


- Molecule 2: PCNA2 (SSO1047)



- Molecule 3: PCNA3 (SSO0405)



- Molecule 3: PCNA3 (SSO0405)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	36.52Å 87.18Å 135.87Å 89.80° 86.36° 78.33°	Depositor
Resolution (Å)	20.00 – 2.79 45.20 – 2.79	Depositor EDS
% Data completeness (in resolution range)	92.8 (20.00-2.79) 92.7 (45.20-2.79)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.56 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.215 , 0.282 0.206 , 0.271	Depositor DCC
R_{free} test set	1892 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	39.9	Xtriage
Anisotropy	0.906	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 28.4	EDS
Estimated twinning fraction	0.000 for -h,-h+k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 37655 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11550	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.04 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.8350e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/1947	0.68	1/2610 (0.0%)
1	X	0.51	0/1947	0.68	0/2610
2	B	0.47	0/1954	0.68	2/2636 (0.1%)
2	Y	0.55	0/1954	0.73	2/2636 (0.1%)
3	C	0.53	0/1949	0.68	1/2622 (0.0%)
3	Z	0.51	0/1949	0.65	0/2622
All	All	0.52	0/11700	0.68	6/15736 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	217	LEU	CA-CB-CG	5.94	128.95	115.30
1	A	125	LEU	CA-CB-CG	5.73	128.47	115.30
2	Y	217	LEU	CA-CB-CG	5.37	127.64	115.30
2	Y	91	LEU	CA-CB-CG	5.31	127.50	115.30
2	B	135	LEU	CA-CB-CG	5.13	127.11	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1927	0	1976	50	0
1	X	1927	0	1976	57	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1922	0	1920	58	0
2	Y	1922	0	1920	60	0
3	C	1926	0	1936	53	0
3	Z	1926	0	1936	57	0
All	All	11550	0	11664	311	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 311 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:149:ASP:HB2	2:Y:110:PHE:HZ	1.13	1.13
2:Y:215:MSE:HE2	2:Y:229:PHE:CE2	1.89	1.06
1:X:149:ASP:HB3	2:Y:82:ARG:HE	1.16	1.05
2:Y:137:THR:HB	2:Y:215:MSE:HE3	1.33	1.03
2:Y:80:LEU:O	2:Y:83:VAL:HG22	1.61	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/257 (96%)	237 (96%)	9 (4%)	1 (0%)	39	74
1	X	247/257 (96%)	238 (96%)	8 (3%)	1 (0%)	39	74
2	B	241/245 (98%)	228 (95%)	11 (5%)	2 (1%)	24	58
2	Y	241/245 (98%)	231 (96%)	8 (3%)	2 (1%)	24	58
3	C	241/252 (96%)	235 (98%)	4 (2%)	2 (1%)	24	58
3	Z	241/252 (96%)	233 (97%)	6 (2%)	2 (1%)	24	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1458/1508 (97%)	1402 (96%)	46 (3%)	10 (1%)	26	62

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	124	ILE
2	B	148	SER
3	C	190	LYS
2	Y	182	ASN
3	Z	190	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/219 (100%)	190 (87%)	29 (13%)	5	14
1	X	219/219 (100%)	182 (83%)	37 (17%)	2	7
2	B	218/215 (101%)	178 (82%)	40 (18%)	2	6
2	Y	218/215 (101%)	177 (81%)	41 (19%)	2	6
3	C	219/224 (98%)	187 (85%)	32 (15%)	4	11
3	Z	219/224 (98%)	189 (86%)	30 (14%)	4	13
All	All	1312/1316 (100%)	1103 (84%)	209 (16%)	3	9

5 of 209 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	183	LEU
1	X	85	LYS
3	Z	105	THR
3	C	194	ASN
1	X	49	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
2	Y	27	ASN
2	Y	121	GLN
3	Z	106	ASN
1	X	240	HIS
3	Z	119	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/257 (93%)	-0.04	1 (0%) 93 90	28, 32, 37, 44	0
1	X	240/257 (93%)	0.04	2 (0%) 87 81	27, 32, 36, 40	0
2	B	239/245 (97%)	0.07	2 (0%) 87 81	28, 32, 36, 42	0
2	Y	239/245 (97%)	-0.16	0 100 100	28, 32, 35, 40	0
3	C	239/252 (94%)	-0.09	4 (1%) 73 63	28, 32, 37, 49	0
3	Z	239/252 (94%)	-0.00	3 (1%) 79 71	28, 32, 36, 46	0
All	All	1436/1508 (95%)	-0.03	12 (0%) 87 81	27, 32, 36, 49	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	83	SER	3.8
3	C	122	PRO	3.6
1	A	83	SER	3.3
2	B	120	THR	2.8
3	C	124	ILE	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.