wwPDB X-ray Structure Validation Summary Report (i)

Feb 1, 2016 – 09:12 AM GMT

PDB ID	:	3HL6
Title	:	Staphylococcus aureus pathogenicity island 3 ORF9 protein
Authors	:	Kruse, A.C.; Huseby, M.; Shi, K.; Digre, J.; Schlievert, P.M.; Ohlendorf, D.H.; Earhart, C.A.
Deposited on	:	2009-05-26
Resolution	:	2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	15% 58% 21% 5% 16%
1	B	225	10% 56% 26% • 15%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3143 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pathogenicity island protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	189	1536	980	244	302	10	0	0	0
1	B	192	1547	989	246	302	10	0	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ASP	ASN	CONFLICT	UNP Q5HHI6
A	214	LEU	-	EXPRESSION TAG	UNP Q5HHI6
A	215	ALA	-	EXPRESSION TAG	UNP Q5HHI6
A	216	ALA	-	EXPRESSION TAG	UNP Q5HHI6
A	217	ALA	-	EXPRESSION TAG	UNP Q5HHI6
A	218	LEU	-	EXPRESSION TAG	UNP Q5HHI6
A	219	GLU	-	EXPRESSION TAG	UNP Q5HHI6
A	220	HIS	-	EXPRESSION TAG	UNP Q5HHI6
A	221	HIS	-	EXPRESSION TAG	UNP Q5HHI6
A	222	HIS	-	EXPRESSION TAG	UNP Q5HHI6
A	223	HIS	-	EXPRESSION TAG	UNP Q5HHI6
A	224	HIS	-	EXPRESSION TAG	UNP Q5HHI6
A	225	HIS	-	EXPRESSION TAG	UNP Q5HHI6
B	2	ASP	ASN	CONFLICT	UNP Q5HHI6
B	214	LEU	-	EXPRESSION TAG	UNP Q5HHI6
B	215	ALA	-	EXPRESSION TAG	UNP Q5HHI6
B	216	ALA	-	EXPRESSION TAG	UNP Q5HHI6
B	217	ALA	-	EXPRESSION TAG	UNP Q5HHI6
B	218	LEU	-	EXPRESSION TAG	UNP Q5HHI6
B	219	GLU	-	EXPRESSION TAG	UNP Q5HHI6
B	220	HIS	-	EXPRESSION TAG	UNP Q5HHI6
B	221	HIS	-	EXPRESSION TAG	UNP Q5HHI6
B	222	HIS	-	EXPRESSION TAG	UNP Q5HHI6
B	223	HIS	-	EXPRESSION TAG	UNP Q5HHI6
B	224	HIS	-	EXPRESSION TAG	UNP Q5HHI6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	225	HIS	-	EXPRESSION TAG	UNP Q5HHI6

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
2	B	1	1	1	0	0
			Total	Cl		
2	A	1	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	22	22	22	0	0
			Total	O		
3	B	36	36	36	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pathogenicity island protein

Chain A:



- Molecule 1: Pathogenicity island protein

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	82.74Å 78.58Å 83.54Å 90.00° 107.41° 90.00°	Depositor
Resolution (Å)	30.00 – 2.50 28.21 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.6 (30.00-2.50) 96.7 (28.21-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$< I/\sigma(I) >$ [1]	2.70 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.228 , 0.276 0.226 , 0.276	Depositor DCC
R_{free} test set	889 reflections (5.46%)	DCC
Wilson B-factor (Å^2)	55.5	Xtriage
Anisotropy	0.189	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 48.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 17182 reflections	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	3143	wwPDB-VP
Average B, all atoms (Å^2)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.17 % of the origin peak, indicating pseudo translational symmetry.The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.2106e-03.The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.67	1/1565 (0.1%)	0.67	0/2111
1	B	0.71	1/1576 (0.1%)	0.70	0/2127
All	All	0.69	2/3141 (0.1%)	0.68	0/4238

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	73	PRO	N-CD	8.71	1.60	1.47
1	A	148	SER	CB-OG	7.74	1.52	1.42

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	187	GLU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1536	0	1499	69	0
1	B	1547	0	1518	69	0
2	A	1	0	0	1	0
2	B	1	0	0	0	0
3	A	22	0	0	2	0
3	B	36	0	0	2	0
All	All	3143	0	3017	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 137 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:ILE:CD1	1:B:3:ILE:CG1	1.80	1.57
1:B:122:ALA:HB1	1:B:189:ALA:O	1.15	1.22
1:A:58:TYR:CE1	1:A:60:LYS:HG2	1.75	1.21
1:B:60:LYS:HD3	1:B:60:LYS:H	0.98	1.15
1:A:191:GLN:HA	1:A:191:GLN:NE2	1.63	1.11

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	185/225 (82%)	170 (92%)	15 (8%)	0	100 100
1	B	188/225 (84%)	180 (96%)	7 (4%)	1 (0%)	34 55
All	All	373/450 (83%)	350 (94%)	22 (6%)	1 (0%)	46 68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	27	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	172/202 (85%)	154 (90%)	18 (10%)	8 16
1	B	171/202 (85%)	155 (91%)	16 (9%)	11 20
All	All	343/404 (85%)	309 (90%)	34 (10%)	10 18

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	185	GLN
1	B	58	TYR
1	B	188	ASN
1	A	195	LYS
1	A	62	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	191	GLN
1	B	202	ASN
1	B	119	GLN
1	A	119	GLN
1	B	188	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	189/225 (84%)	0.94	33 (17%)	2	2	40, 60, 91, 100	0
1	B	192/225 (85%)	0.83	23 (11%)	6	6	37, 56, 79, 89	0
All	All	381/450 (84%)	0.89	56 (14%)	3	3	37, 58, 85, 100	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	93	HIS	7.9
1	A	186	ASP	6.5
1	B	92	LEU	6.3
1	B	94	ASP	6.1
1	A	30	VAL	6.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
2	CL	A	226	1/1	0.89	0.12	-0.90	69,69,69,69	0
2	CL	B	226	1/1	0.92	0.08	-1.53	64,64,64,64	0

6.5 Other polymers (i)

There are no such residues in this entry.