



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:17 PM GMT

PDB ID : 4HU4
Title : Crystal structure of EAL domain of the E. coli DosP - dimeric form
Authors : Tarnawski, M.; Barends, T.R.M.; Hartmann, E.; Schlichting, I.
Deposited on : 2012-11-02
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	<div> <div>4%</div> <div>63%</div> <div>21%</div> <div>15%</div> </div>
1	B	292	<div> <div>4%</div> <div>57%</div> <div>25%</div> <div>15%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxygen sensor protein DosP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	247	Total	C	N	O	S	0	0	0
			1962	1253	341	358	10			
1	B	247	Total	C	N	O	S	0	0	0
			1962	1253	341	358	10			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	508	MET	-	EXPRESSION TAG	UNP P76129
A	509	GLY	-	EXPRESSION TAG	UNP P76129
A	510	SER	-	EXPRESSION TAG	UNP P76129
A	511	SER	-	EXPRESSION TAG	UNP P76129
A	512	HIS	-	EXPRESSION TAG	UNP P76129
A	513	HIS	-	EXPRESSION TAG	UNP P76129
A	514	HIS	-	EXPRESSION TAG	UNP P76129
A	515	HIS	-	EXPRESSION TAG	UNP P76129
A	516	HIS	-	EXPRESSION TAG	UNP P76129
A	517	HIS	-	EXPRESSION TAG	UNP P76129
A	518	SER	-	EXPRESSION TAG	UNP P76129
A	519	SER	-	EXPRESSION TAG	UNP P76129
A	520	GLY	-	EXPRESSION TAG	UNP P76129
A	521	LEU	-	EXPRESSION TAG	UNP P76129
A	522	VAL	-	EXPRESSION TAG	UNP P76129
A	523	PRO	-	EXPRESSION TAG	UNP P76129
A	524	ARG	-	EXPRESSION TAG	UNP P76129
A	525	GLY	-	EXPRESSION TAG	UNP P76129
A	526	SER	-	EXPRESSION TAG	UNP P76129
A	527	HIS	-	EXPRESSION TAG	UNP P76129
A	528	MET	-	EXPRESSION TAG	UNP P76129
B	508	MET	-	EXPRESSION TAG	UNP P76129
B	509	GLY	-	EXPRESSION TAG	UNP P76129
B	510	SER	-	EXPRESSION TAG	UNP P76129
B	511	SER	-	EXPRESSION TAG	UNP P76129

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	512	HIS	-	EXPRESSION TAG	UNP P76129
B	513	HIS	-	EXPRESSION TAG	UNP P76129
B	514	HIS	-	EXPRESSION TAG	UNP P76129
B	515	HIS	-	EXPRESSION TAG	UNP P76129
B	516	HIS	-	EXPRESSION TAG	UNP P76129
B	517	HIS	-	EXPRESSION TAG	UNP P76129
B	518	SER	-	EXPRESSION TAG	UNP P76129
B	519	SER	-	EXPRESSION TAG	UNP P76129
B	520	GLY	-	EXPRESSION TAG	UNP P76129
B	521	LEU	-	EXPRESSION TAG	UNP P76129
B	522	VAL	-	EXPRESSION TAG	UNP P76129
B	523	PRO	-	EXPRESSION TAG	UNP P76129
B	524	ARG	-	EXPRESSION TAG	UNP P76129
B	525	GLY	-	EXPRESSION TAG	UNP P76129
B	526	SER	-	EXPRESSION TAG	UNP P76129
B	527	HIS	-	EXPRESSION TAG	UNP P76129
B	528	MET	-	EXPRESSION TAG	UNP P76129

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	57	Total O 57 57	0	0
2	B	59	Total O 59 59	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.60Å 171.17Å 48.83Å 90.00° 90.03° 90.00°	Depositor
Resolution (Å)	19.99 – 2.40 19.99 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.99-2.40) 99.6 (19.99-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 2.41Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.233 , 0.280 0.224 , 0.273	Depositor DCC
R_{free} test set	1220 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	48.4	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 32.5	EDS
Estimated twinning fraction	0.480 for h,-k,-l 0.430 for h,-k,-l	Xtriage
Reported twinning fraction	0.480 for h,-k,-l	Depositor
L-test for twinning ²	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.20$	Xtriage
Outliers	0 of 22645 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4040	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/2003	0.41	0/2714
1	B	0.22	0/2003	0.42	0/2714
All	All	0.22	0/4006	0.41	0/5428

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1962	0	1976	37	0
1	B	1962	0	1976	50	0
2	A	57	0	0	3	0
2	B	59	0	0	8	0
All	All	4040	0	3952	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:752:VAL:HG12	1:A:772:VAL:HB	1.68	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:541:ARG:HD3	1:B:597:LEU:HD21	1.72	0.71
1:A:641:PHE:O	1:A:643:SER:N	2.23	0.71
1:B:612:ILE:HD13	1:B:650:VAL:HG22	1.74	0.68
1:B:685:GLN:NE2	2:B:835:HOH:O	2.28	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/292 (83%)	218 (90%)	22 (9%)	3 (1%)	16	23
1	B	243/292 (83%)	220 (90%)	20 (8%)	3 (1%)	16	23
All	All	486/584 (83%)	438 (90%)	42 (9%)	6 (1%)	16	23

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	628	HIS
1	A	642	ARG
1	B	583	ASP
1	A	659	ILE
1	B	659	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/253 (85%)	210 (98%)	4 (2%)	65	83
1	B	214/253 (85%)	204 (95%)	10 (5%)	32	50
All	All	428/506 (85%)	414 (97%)	14 (3%)	45	66

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	597	LEU
1	B	606	ASN
1	B	674	MET
1	B	559	LEU
1	B	646	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/292 (84%)	0.27	12 (4%) 33 34	14, 32, 62, 75	0
1	B	247/292 (84%)	0.32	13 (5%) 30 30	13, 32, 61, 74	0
All	All	494/584 (84%)	0.30	25 (5%) 32 32	13, 32, 62, 75	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	597	LEU	6.6
1	A	678	THR	3.8
1	A	637	SER	3.8
1	B	680	ILE	3.3
1	A	698	ASP	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.