



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:18 PM GMT

PDB ID : 4HXA
Title : Crystal structure of 13D9 FAB
Authors : Johal, A.R.; Jarrell, H.C.; Khieu, N.H.; Letts, J.A.; Landry, R.C.; Jachymek, W.; Yang, Q.; Jennings, H.J.; Brisson, J.R.; Evans, S.V.
Deposited on : 2012-11-09
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1799 (2.08-2.04)
Clashscore	102246	1910 (2.08-2.04)
Ramachandran outliers	100387	1893 (2.08-2.04)
Sidechain outliers	100360	1893 (2.08-2.04)
RSRZ outliers	91569	1802 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	213	
2	H	221	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 13D9 FAB LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	213	Total	C	N	O	S	0	4	0
			1662	1035	270	347	10			

- Molecule 2 is a protein called 13D9 FAB HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	221	Total	C	N	O	S	0	2	0
			1694	1062	291	334	7			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	187	Total	O	0	0
			187	187		
3	H	174	Total	O	0	0
			174	174		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 13D9 FAB LIGHT CHAIN



• Molecule 2: 13D9 FAB HEAVY CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.75Å 100.28Å 132.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.06 19.83 – 2.06	Depositor EDS
% Data completeness (in resolution range)	88.4 (20.00-2.06) 88.4 (19.83-2.06)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.45 (at 2.06Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.213 , 0.286 0.211 , 0.283	Depositor DCC
R_{free} test set	1556 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.446	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	0 of 31108 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3717	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	1.03	1/1710 (0.1%)	0.90	3/2321 (0.1%)
2	H	1.03	1/1738 (0.1%)	1.16	11/2369 (0.5%)
All	All	1.03	2/3448 (0.1%)	1.04	14/4690 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1
2	H	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	27	SER	C-N	10.39	1.57	1.34
2	H	111	VAL	CB-CG2	5.75	1.65	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	66	ARG	NE-CZ-NH2	-18.96	110.82	120.30
2	H	66	ARG	NE-CZ-NH1	16.55	128.58	120.30
1	L	73	LEU	CA-CB-CG	8.71	135.32	115.30
2	H	44	ARG	NE-CZ-NH1	-7.29	116.66	120.30
1	L	73	LEU	CB-CG-CD1	6.95	122.82	111.00
2	H	44	ARG	NE-CZ-NH2	6.72	123.66	120.30
2	H	136	VAL	CB-CA-C	-6.61	98.83	111.40
2	H	181	VAL	CB-CA-C	-6.30	99.43	111.40
2	H	66	ARG	CD-NE-CZ	6.19	132.26	123.60
1	L	108	ARG	NE-CZ-NH2	-6.11	117.25	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	80	LEU	CB-CG-CD1	5.73	120.74	111.00
2	H	18	LEU	CA-CB-CG	5.37	127.64	115.30
2	H	174	LEU	CA-CB-CG	5.35	127.60	115.30
2	H	82	MET	CG-SD-CE	-5.34	91.66	100.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	133	GLY	Peptide
1	L	27	SER	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1662	0	1566	38	0
2	H	1694	0	1669	42	1
3	H	174	0	0	8	1
3	L	187	0	0	7	0
All	All	3717	0	3235	74	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:25:SER:HA	3:H:356:HOH:O	1.34	1.25
2:H:133:GLY:HA3	2:H:134:SER:CB	1.96	0.96
2:H:133:GLY:HA3	2:H:134:SER:HB3	1.48	0.94
1:L:38:GLN:HE22	2:H:39:GLN:HE22	1.25	0.82
1:L:142:LYS:O	1:L:163[B]:TRP:CZ3	2.36	0.79
1:L:190:ASN:HD21	1:L:212:ASN:H	1.32	0.78
1:L:144:ILE:HG22	1:L:163[B]:TRP:CH2	2.20	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:142:LYS:O	1:L:163[B]:TRP:HZ3	1.71	0.73
2:H:3:GLN:HB2	2:H:25:SER:HB3	1.70	0.73
1:L:114:THR:HG21	3:H:418:HOH:O	1.90	0.71
1:L:160:LEU:HD21	2:H:171:GLN:NE2	2.05	0.71
1:L:31:ASN:HD22	1:L:32:TYR:H	1.40	0.70
2:H:25:SER:CA	3:H:356:HOH:O	2.07	0.68
2:H:67:PHE:CZ	2:H:82:MET:HE2	2.29	0.66
1:L:163[B]:TRP:HZ2	3:L:319:HOH:O	1.79	0.64
2:H:159:LEU:HD13	2:H:181:VAL:HG11	1.79	0.64
2:H:83:ARG:HG3	2:H:85[B]:GLU:HG2	1.80	0.63
1:L:156:GLN:HA	3:L:360:HOH:O	1.99	0.62
2:H:3:GLN:HB2	2:H:25:SER:CB	2.30	0.60
1:L:4:LEU:HD12	1:L:25:VAL:HG13	1.83	0.60
1:L:160:LEU:HD21	2:H:171:GLN:HE21	1.65	0.60
1:L:160:LEU:CD2	2:H:171:GLN:NE2	2.65	0.60
1:L:105:GLU:OE1	1:L:142:LYS:NZ	2.36	0.59
2:H:143:LYS:HA	2:H:176:THR:HB	1.84	0.59
1:L:119:PRO:HD2	2:H:213:ARG:NH1	2.18	0.59
1:L:214:CYS:HB3	3:H:376:HOH:O	2.04	0.58
1:L:190:ASN:ND2	1:L:212:ASN:H	2.00	0.57
2:H:188:TRP:CD1	2:H:193:ILE:HD12	2.40	0.57
1:L:160:LEU:CD2	2:H:171:GLN:HE21	2.18	0.57
2:H:66:ARG:HD3	3:H:302:HOH:O	2.05	0.56
1:L:142:LYS:O	1:L:163[B]:TRP:CH2	2.60	0.54
1:L:31:ASN:HD22	1:L:32:TYR:N	2.04	0.54
1:L:108:ARG:HD3	1:L:109:ALA:O	2.07	0.54
1:L:214:CYS:OXT	3:L:374:HOH:O	2.18	0.53
1:L:141:PRO:O	1:L:198:HIS:HE1	1.92	0.53
2:H:154:TRP:CE2	2:H:181:VAL:HG22	2.44	0.52
2:H:75:LYS:CE	2:H:79:TYR:OH	2.58	0.52
2:H:133:GLY:HA3	2:H:134:SER:OG	2.09	0.52
1:L:90:GLN:HE22	1:L:93:GLY:H	1.57	0.51
1:L:2:GLU:HG2	1:L:30:VAL:CG1	2.41	0.51
1:L:114:THR:HG22	3:L:483:HOH:O	2.12	0.50
3:L:443:HOH:O	2:H:164:HIS:HD2	1.94	0.50
1:L:12:SER:HA	1:L:105:GLU:O	2.12	0.49
1:L:213:GLU:O	1:L:214:CYS:HB2	2.13	0.49
2:H:163:VAL:HG22	2:H:181:VAL:HG13	1.95	0.48
2:H:143:LYS:HG2	2:H:176:THR:HB	1.95	0.47
2:H:66:ARG:CD	3:H:302:HOH:O	2.62	0.47
1:L:198:HIS:HD2	1:L:200:THR:OG1	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:147:PRO:O	2:H:199:HIS:HE1	1.98	0.46
1:L:144:ILE:HG22	1:L:163[B]:TRP:HH2	1.79	0.46
2:H:52(B):LYS:HA	2:H:52(B):LYS:CE	2.46	0.46
2:H:75:LYS:HE2	2:H:79:TYR:OH	2.16	0.45
1:L:150:ILE:HD11	1:L:179:LEU:HD21	1.98	0.45
2:H:176:THR:HG22	3:H:354:HOH:O	2.15	0.45
1:L:163[B]:TRP:HE3	3:L:413:HOH:O	1.99	0.44
2:H:72:ASP:OD1	2:H:75:LYS:HD3	2.17	0.44
1:L:90:GLN:NE2	1:L:92:SER:H	2.16	0.44
1:L:163[A]:TRP:CD1	1:L:175:MET:HG3	2.52	0.44
2:H:133:GLY:CA	2:H:134:SER:CB	2.81	0.43
2:H:83:ARG:HE	2:H:85[B]:GLU:CD	2.22	0.43
2:H:202:SER:O	2:H:204:THR:HG23	2.19	0.43
1:L:173:TYR:HB3	3:L:413:HOH:O	2.19	0.43
1:L:182:THR:OG1	1:L:185[B]:GLU:HB2	2.19	0.43
2:H:85[A]:GLU:H	2:H:85[A]:GLU:CD	2.22	0.42
2:H:66:ARG:NH2	2:H:86:ASP:OD2	2.34	0.42
2:H:176:THR:CG2	3:H:354:HOH:O	2.68	0.42
2:H:53:ASP:HB3	2:H:55:TYR:H	1.83	0.42
2:H:11:LEU:HD11	2:H:112:SER:HB3	2.02	0.42
2:H:35:HIS:CE1	2:H:50:VAL:HG23	2.54	0.41
2:H:103:TRP:N	2:H:103:TRP:CD1	2.88	0.41
2:H:52(B):LYS:HA	2:H:52(B):LYS:HE3	2.01	0.41
1:L:203:SER:HA	1:L:204:PRO:HD2	1.82	0.41
2:H:1:GLN:O	2:H:26:GLY:HA3	2.21	0.41
1:L:145:ASN:HB3	1:L:197:THR:HB	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:215:PRO:O	3:H:309:HOH:O[2_554]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	215/213 (101%)	209 (97%)	6 (3%)	0	100	100
2	H	221/221 (100%)	213 (96%)	6 (3%)	2 (1%)	21	10
All	All	436/434 (100%)	422 (97%)	12 (3%)	2 (0%)	34	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	134	SER
2	H	54	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	192/188 (102%)	179 (93%)	13 (7%)	20	11
2	H	195/193 (101%)	171 (88%)	24 (12%)	6	2
All	All	387/381 (102%)	350 (90%)	37 (10%)	11	4

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	1	GLU
1	L	2	GLU
1	L	31	ASN
1	L	73	LEU
1	L	90	GLN
1	L	108	ARG
1	L	114	THR
1	L	143	ASP
1	L	156	GLN
1	L	159	VAL
1	L	160	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	169	LYS
1	L	212	ASN
2	H	3	GLN
2	H	18	LEU
2	H	25	SER
2	H	43	LYS
2	H	44	ARG
2	H	52(B)	LYS
2	H	58	ASN
2	H	80	LEU
2	H	93[A]	SER
2	H	93[B]	SER
2	H	109	VAL
2	H	115	LYS
2	H	124	LEU
2	H	135	SER
2	H	136	VAL
2	H	138	LEU
2	H	142	VAL
2	H	159	LEU
2	H	170	LEU
2	H	171	GLN
2	H	174	LEU
2	H	176	THR
2	H	181	VAL
2	H	187	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	31	ASN
1	L	38	GLN
1	L	90	GLN
1	L	190	ASN
1	L	198	HIS
1	L	212	ASN
2	H	31	ASN
2	H	58	ASN
2	H	164	HIS
2	H	171	GLN
2	H	199	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	213/213 (100%)	-0.28	3 (1%) 78 80	21, 30, 42, 53	0
2	H	221/221 (100%)	0.01	12 (5%) 29 32	22, 31, 48, 67	0
All	All	434/434 (100%)	-0.13	15 (3%) 48 53	21, 30, 45, 67	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	55	TYR	7.9
2	H	54	ILE	7.3
2	H	133	GLY	5.9
2	H	131	THR	4.2
2	H	132	THR	3.7
2	H	52(B)	LYS	3.5
2	H	53	ASP	3.4
2	H	52(C)	SER	3.1
1	L	163[A]	TRP	2.7
2	H	93[A]	SER	2.6
1	L	56	SER	2.5
2	H	1	GLN	2.2
2	H	37	LEU	2.2
2	H	128	CYS	2.2
1	L	126	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.