



Full wwPDB X-ray Structure Validation Report ⓘ

May 31, 2016 – 05:05 PM EDT

PDB ID : 5HX5
Title : APOBEC3F Catalytic Domain Crystal Structure
Authors : Shaban, N.M.; Shi, K.; Aihara, H.; Harris, R.S.
Deposited on : 2016-01-29
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674

i

X-RAY DIFFRACTION

A.

 R_{free}

electron density. The numeric value is given above the bar.

1	A	199
---	---	-----

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA dC->dU-editing enzyme APOBEC-3F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	0	0	0
			1464	942	241	272	9			
1	B	168	Total	C	N	O	S	8	0	0
			1417	915	229	264	9			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	175	GLY	-	expression tag	UNP Q8IUX4
A	176	PRO	-	expression tag	UNP Q8IUX4
A	177	LEU	-	expression tag	UNP Q8IUX4
A	178	GLY	-	expression tag	UNP Q8IUX4
A	179	SER	-	expression tag	UNP Q8IUX4
A	180	PRO	-	expression tag	UNP Q8IUX4
A	181	GLY	-	expression tag	UNP Q8IUX4
A	182	ILE	-	expression tag	UNP Q8IUX4
A	183	PRO	-	expression tag	UNP Q8IUX4
A	184	GLY	-	expression tag	UNP Q8IUX4
A	196	ASP	TYR	engineered mutation	UNP Q8IUX4
A	247	GLY	HIS	engineered mutation	UNP Q8IUX4
A	248	ARG	CYS	engineered mutation	UNP Q8IUX4
A	259	ALA	CYS	engineered mutation	UNP Q8IUX4
A	302	LYS	PHE	engineered mutation	UNP Q8IUX4
A	310	ASP	TRP	engineered mutation	UNP Q8IUX4
A	355	ASP	LYS	engineered mutation	UNP Q8IUX4
A	358	ASP	LYS	engineered mutation	UNP Q8IUX4
A	363	ASP	PHE	engineered mutation	UNP Q8IUX4
B	175	GLY	-	expression tag	UNP Q8IUX4
B	176	PRO	-	expression tag	UNP Q8IUX4
B	177	LEU	-	expression tag	UNP Q8IUX4
B	178	GLY	-	expression tag	UNP Q8IUX4
B	179	SER	-	expression tag	UNP Q8IUX4
B	180	PRO	-	expression tag	UNP Q8IUX4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	181	GLY	-	expression tag	UNP Q8IUX4
B	182	ILE	-	expression tag	UNP Q8IUX4
B	183	PRO	-	expression tag	UNP Q8IUX4
B	184	GLY	-	expression tag	UNP Q8IUX4
B	196	ASP	TYR	engineered mutation	UNP Q8IUX4
B	247	GLY	HIS	engineered mutation	UNP Q8IUX4
B	248	ARG	CYS	engineered mutation	UNP Q8IUX4
B	259	ALA	CYS	engineered mutation	UNP Q8IUX4
B	302	LYS	PHE	engineered mutation	UNP Q8IUX4
B	310	ASP	TRP	engineered mutation	UNP Q8IUX4
B	355	ASP	LYS	engineered mutation	UNP Q8IUX4
B	358	ASP	LYS	engineered mutation	UNP Q8IUX4
B	363	ASP	PHE	engineered mutation	UNP Q8IUX4

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	31	Total O 31 31	0	0
3	B	30	Total O 30 30	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3F



- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3F



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.94Å 50.90Å 100.86Å 90.00° 90.75° 90.00°	Depositor
Resolution (Å)	35.94 – 2.33 35.94 – 2.33	Depositor EDS
% Data completeness (in resolution range)	93.2 (35.94-2.33) 93.7 (35.94-2.33)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 2.34Å)	Xtriage
Refinement program	PHENIX (dev_2313: ???)	Depositor
R, R_{free}	0.227 , 0.264 0.230 , 0.238	Depositor DCC
R_{free} test set	772 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	44.6	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.126 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2944	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/1510	0.43	0/2046
1	B	0.26	0/1465	0.45	0/1989
All	All	0.27	0/2975	0.44	0/4035

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1464	0	1345	17	0
1	B	1417	0	1292	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	0	0	1	0
3	B	30	0	0	3	0
All	All	2944	0	2637	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:223:GLU:HB3	1:B:270:GLU:HB2	1.64	0.80
1:B:205:LYS:O	1:B:305:ARG:NH1	2.22	0.72
1:B:309:PHE:O	3:B:501:HOH:O	2.16	0.63
1:A:225:VAL:HA	1:A:231:VAL:HG22	1.80	0.62
1:B:302:LYS:NZ	3:B:504:HOH:O	2.36	0.59
1:A:252:ARG:HE	1:A:286:GLU:HG3	1.70	0.56
1:B:203:HIS:O	1:B:276:SER:HB2	2.10	0.52
1:B:319:ARG:O	1:B:323:GLN:HG2	2.11	0.51
1:B:278:SER:HB3	1:B:307:TYR:H	1.76	0.51
1:B:226:LYS:HB2	1:B:229:SER:HB3	1.93	0.50
1:A:305:ARG:NH2	3:A:503:HOH:O	2.41	0.49
1:A:319:ARG:O	1:A:323:GLN:HG2	2.12	0.48
1:A:222:MET:HE1	1:A:234:LYS:HD2	1.96	0.48
1:B:289:GLU:OE2	1:B:293:ARG:NH1	2.47	0.47
1:B:313:ASP:OD1	1:B:313:ASP:N	2.45	0.47
1:B:239:ARG:NH2	3:B:506:HOH:O	2.48	0.46
1:B:257:TRP:O	1:B:261:ASP:HB2	2.16	0.46
1:A:300:THR:HG23	1:A:327:SER:HB3	1.98	0.45
1:B:217:TRP:CZ2	1:B:239:ARG:HD2	2.51	0.45
1:A:224:VAL:O	1:A:231:VAL:HA	2.17	0.45
1:A:227:HIS:O	1:A:227:HIS:ND1	2.41	0.45
1:A:213:ARG:HA	1:A:213:ARG:CZ	2.47	0.44
1:A:220:PHE:HA	1:A:272:THR:O	2.17	0.44
1:B:203:HIS:O	1:B:305:ARG:NH2	2.49	0.43
1:B:221:THR:HG23	1:B:343:PHE:HZ	1.84	0.43
1:A:334:LYS:HD2	1:A:334:LYS:HA	1.84	0.42
1:A:340:TRP:CD1	1:A:351:PHE:HB2	2.54	0.42
1:A:219:CYS:HB2	1:A:274:TYR:HB2	2.01	0.42
1:B:340:TRP:CD1	1:B:351:PHE:HB2	2.54	0.42
1:B:339:CYS:HA	1:B:343:PHE:HD2	1.85	0.41
1:A:207:LEU:HD23	1:A:360:ASN:OD1	2.19	0.41
1:A:255:LEU:HD11	1:A:287:VAL:HG22	2.03	0.41
1:A:223:GLU:HB3	1:A:270:GLU:HB2	2.02	0.41
1:A:278:SER:OG	1:A:318:LEU:HD21	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/199 (84%)	164 (98%)	4 (2%)	0	100	100
1	B	162/199 (81%)	157 (97%)	5 (3%)	0	100	100
All	All	330/398 (83%)	321 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	157/178 (88%)	153 (98%)	4 (2%)	55	68
1	B	154/178 (86%)	150 (97%)	4 (3%)	54	66
All	All	311/356 (87%)	303 (97%)	8 (3%)	54	66

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	227	HIS
1	A	261	ASP
1	A	309	PHE
1	A	310	ASP
1	B	248	ARG
1	B	261	ASP
1	B	308	TYR
1	B	309	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	203	HIS
1	B	315	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	174/199 (87%)	0.08	3 (1%) 73 82	31, 51, 87, 113	0
1	B	168/199 (84%)	0.41	9 (5%) 29 42	38, 68, 113, 147	1 (0%)
All	All	342/398 (85%)	0.24	12 (3%) 48 59	31, 59, 106, 147	1 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	202	PHE	3.9
1	B	231	VAL	3.6
1	A	226	LYS	3.5
1	B	198	HIS	3.2
1	A	263	LEU	3.1
1	B	346	ASN	2.9
1	B	201	TYR	2.7
1	B	359	TYR	2.7
1	B	308	TYR	2.5
1	B	265	PRO	2.2
1	B	270	GLU	2.2
1	A	373	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ZN	A	401	1/1	0.99	0.07	-	54,54,54,54	0
2	ZN	B	401	1/1	0.99	0.06	-	67,67,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.