



wwPDB X-ray Structure Validation Summary Report ⓘ

May 4, 2016 – 02:28 PM EDT

PDB ID : 5HYN
Title : Structure of Human Polycomb Repressive Complex 2 (PRC2) with oncogenic histone H3K27M peptide
Authors : Zhang, Y.; Justin, N.; Wilson, J.R.; Gamblin, S.J.
Deposited on : 2016-02-01
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027457
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2184 (3.00-2.92)
Clashscore	102246	2552 (3.00-2.92)
Ramachandran outliers	100387	2468 (3.00-2.92)
Sidechain outliers	100360	2471 (3.00-2.92)
RSRZ outliers	91569	2201 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	746	<div> <div>7%</div> <div> <div>47%</div> <div>28%</div> <div>•</div> <div>22%</div> </div> </div>
1	F	746	<div> <div>8%</div> <div> <div>48%</div> <div>25%</div> <div>•</div> <div>24%</div> </div> </div>
1	K	746	<div> <div>10%</div> <div> <div>50%</div> <div>23%</div> <div>•</div> <div>25%</div> </div> </div>
1	Q	746	<div> <div>8%</div> <div> <div>45%</div> <div>27%</div> <div>•</div> <div>24%</div> </div> </div>
2	B	367	<div> <div>3%</div> <div> <div>62%</div> <div>37%</div> <div>••</div> </div> </div>
2	G	367	<div> <div>4%</div> <div> <div>67%</div> <div>31%</div> <div>••</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	L	367	
2	R	367	
3	C	129	
3	H	129	
3	M	129	
3	S	129	
4	D	13	
4	I	13	
4	O	13	
4	T	13	
5	E	12	
5	J	12	
5	P	12	
5	U	12	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ZN	A	1004	-	-	-	X
6	ZN	F	1003	-	-	-	X
6	ZN	K	1004	-	-	-	X
6	ZN	K	1007	-	-	-	X
6	ZN	Q	1004	-	-	-	X
6	ZN	Q	1007	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 35028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase EZH2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	581	Total	C	N	O	S	0	0	0
			4673	2932	825	874	42			
1	F	568	Total	C	N	O	S	0	0	0
			4567	2869	804	852	42			
1	K	562	Total	C	N	O	S	0	0	0
			4521	2836	799	844	42			
1	Q	565	Total	C	N	O	S	0	0	0
			4542	2850	801	849	42			

- Molecule 2 is a protein called Polycomb protein EED.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	365	Total	C	N	O	S	0	0	0
			2959	1873	521	543	22			
2	G	365	Total	C	N	O	S	0	0	0
			2959	1873	521	543	22			
2	L	365	Total	C	N	O	S	0	0	0
			2959	1873	521	543	22			
2	R	365	Total	C	N	O	S	0	0	0
			2959	1873	521	543	22			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	75	GLY	-	expression tag	UNP O75530
B	76	SER	-	expression tag	UNP O75530
G	75	GLY	-	expression tag	UNP O75530
G	76	SER	-	expression tag	UNP O75530
L	75	GLY	-	expression tag	UNP O75530
L	76	SER	-	expression tag	UNP O75530
R	75	GLY	-	expression tag	UNP O75530
R	76	SER	-	expression tag	UNP O75530

- Molecule 3 is a protein called Polycomb protein SUZ12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	125	Total	C	N	O	S	0	0	0
			1042	657	180	193	12			
3	H	124	Total	C	N	O	S	0	0	0
			1032	651	177	192	12			
3	M	125	Total	C	N	O	S	0	0	0
			1042	657	180	193	12			
3	S	125	Total	C	N	O	S	0	0	0
			1042	657	180	193	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	557	GLY	-	expression tag	UNP Q15022
H	557	GLY	-	expression tag	UNP Q15022
M	557	GLY	-	expression tag	UNP Q15022
S	557	GLY	-	expression tag	UNP Q15022

- Molecule 4 is a protein called H3K27M.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	9	Total	C	N	O	S	0	0	0
			63	38	13	11	1			
4	I	9	Total	C	N	O	S	0	0	0
			63	38	13	11	1			
4	O	9	Total	C	N	O	S	0	0	0
			63	38	13	11	1			
4	T	9	Total	C	N	O	S	0	0	0
			63	38	13	11	1			


- Molecule 5 is a protein called JARID2 K116me3.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	11	Total	C	N	O	0	0	0
			96	60	21	15			
5	J	10	Total	C	N	O	0	0	0
			85	54	17	14			
5	P	9	Total	C	N	O	0	0	0
			77	48	16	13			
5	U	10	Total	C	N	O	0	0	0
			85	54	17	14			

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	Q	8	Total	Zn	0	0
			8	8		
6	A	8	Total	Zn	0	0
			8	8		
6	K	8	Total	Zn	0	0
			8	8		
6	F	8	Total	Zn	0	0
			8	8		


- Molecule 7 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C₁₄H₂₀N₆O₅S).





• Molecule 1: Histone-lysine N-methyltransferase EZH2





Chain B:

3% 62% 37%

GLY
SER
K77
Y60
S81
F82
V85
L88
D91
P95
L96
F101
H104
S105
K106
D109
P110
V112
F113
S118
M119
R120
L123
Y124
E125
S128
Q129
G130
L134
Q136
S137
Y138
V139
D140
A141
E145
T153
Y154
S159
H160
P161
L162

V165
R169
G170
I171
I172
R173
M176
P177
M180
K184
H185
Y186
W187
G188
H189
G190
M191
A192
I193
M194
K197
F198
H199
D202
P203
M204
L205
L206
D207
S208
V209
S210
K211
D212
D213
A214
L215
I220
D223
T224
V226
L225
A227
D228
P229
V232
H235
R236
D237
P238
V239
L240
S241
A242
D245
L246
L247
G248
E249
S253
C254
G255
M256
D257
H258
S259
L260
W263
R269
I274
S277
Y280
H281
P282
T285
L286
R287
P288
Q292
K293
L294
R302
R306
V309
D310
L315
L318
K322
S323
C324
N326
W331
P332
R336
S348
V349
V350
T351
L352
L353
F356
D357
Y358
C361
K366
M370
K376
K385
L386
Y387
V388
K389
D390
V393
E394
D395
P396
R397
K398
K399
K400
C401
L402
T405
T416
S417
F418
S419
S422
L425
L426
A427
D430
I434
W437
D438
P439


[illegible]

Chain L:


68% 29%

GLY SER K77 Y80 S81 F82 N86 S87 L88 D91 Q94 P95 F101 F113 S118 N119 R120 V121 T122 L123 H127 E131 I132 R133 L134 L135 Q136 V139 E145 C150 D155 S159 H160 P161 A164 V165 R169 G170 I171 I172 P177 M180


I183 K184 H185 Y186 V187 G188 H189 G190 N191 A192 I193 M194 E195 L196 K197 F198 H199 P200 R201 E202 L205 L206 L207 S208 V209 S210 K211 D212 L215 R216 D223 A227 T228 F229 G230 G231 V232 E238 A242 L246 L247 C254 G255 K256 D257 H258 T265 N266 R267 T268 Q292 K293 I294 E295 R306 V309 L315 G316 D317 L318 I319 L320 S323 C324 E325 N326 W331 K335 M336 S348 I352 L353 G354 R355 Y358 C361 D362 M376 L386 E392 E393 E394 D395 P396 H397 K398 A399 K400 C401 T402 T403 L404 R420 D421 D422




• Molecule 2: Polycomb protein EED




• Molecule 3: Polycomb protein SUZ12




• Molecule 3: Polycomb protein SUZ12



• Molecule 3: Polycomb protein SUZ12




- Molecule 3: Polycomb protein SUZ12




- Molecule 4: H3K27M




- Molecule 4: H3K27M




- Molecule 4: H3K27M



- Molecule 4: H3K27M



- Molecule 5: JARID2 K116me3




- Molecule 5: JARID2 K116me3



● Molecule 5: JARID2 K116me3

Chain P: 



● Molecule 5: JARID2 K116me3

Chain U: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	131.64Å 171.51Å 274.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	95.01 – 2.95 104.43 – 2.94	Depositor EDS
% Data completeness (in resolution range)	99.8 (95.01-2.95) 99.9 (104.43-2.94)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.96Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, R_{free}	0.219 , 0.273 0.235 , 0.281	Depositor DCC
R_{free} test set	12405 reflections (4.95%)	DCC
Wilson B-factor (Å ²)	63.5	Xtriage
Anisotropy	0.627	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 72.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	35028	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.92 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.4013e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SAH, M3L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.66	6/4777 (0.1%)	0.80	8/6443 (0.1%)
1	F	0.67	6/4667 (0.1%)	0.86	14/6293 (0.2%)
1	K	0.61	1/4621 (0.0%)	0.84	8/6232 (0.1%)
1	Q	0.72	6/4641 (0.1%)	0.93	13/6258 (0.2%)
2	B	0.50	0/3034	0.75	1/4107 (0.0%)
2	G	0.57	1/3034 (0.0%)	0.84	5/4107 (0.1%)
2	L	0.58	2/3034 (0.1%)	0.81	3/4107 (0.1%)
2	R	0.59	0/3034	0.82	1/4107 (0.0%)
3	C	0.54	0/1063	0.85	2/1427 (0.1%)
3	H	0.58	1/1052 (0.1%)	0.78	1/1412 (0.1%)
3	M	0.68	3/1063 (0.3%)	0.83	2/1427 (0.1%)
3	S	0.51	0/1063	0.74	2/1427 (0.1%)
4	D	0.50	0/63	0.86	0/83
4	I	0.50	0/63	0.72	0/83
4	O	0.57	0/63	1.07	1/83 (1.2%)
4	T	0.50	0/63	0.88	0/83
5	E	0.60	0/84	0.82	0/110
5	J	0.80	0/73	1.43	1/96 (1.0%)
5	P	0.56	0/65	0.62	0/85
5	U	0.95	1/73 (1.4%)	1.05	0/96
All	All	0.62	27/35630 (0.1%)	0.84	62/48066 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	F	0	1
1	K	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Q	0	4
2	G	0	2
2	L	0	3
2	R	0	1
3	H	0	1
All	All	0	15

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	33	ARG	NE-CZ	-12.29	1.17	1.33
1	F	181	TYR	CB-CG	-10.42	1.36	1.51
1	A	32	PHE	CE1-CZ	8.05	1.52	1.37
1	A	249	GLU	CD-OE1	7.87	1.34	1.25
2	G	396	PRO	N-CD	7.52	1.58	1.47

The worst 5 of 62 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	Q	443	LEU	CB-CG-CD1	-13.41	88.20	111.00
1	Q	33	ARG	NE-CZ-NH2	-11.52	114.54	120.30
1	Q	70	ILE	CG1-CB-CG2	-10.19	88.99	111.40
5	J	6	ARG	NE-CZ-NH2	-9.21	115.69	120.30
2	G	324	CYS	CA-CB-SG	-8.72	98.31	114.00

There are no chirality outliers.

5 of 15 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	102	ASN	Peptide
1	A	134	MET	Peptide
1	F	564	ALA	Peptide
2	G	392	GLU	Peptide
2	G	395	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4673	0	4524	247	1
1	F	4567	0	4427	192	1
1	K	4521	0	4373	180	1
1	Q	4542	0	4400	233	1
2	B	2959	0	2881	148	0
2	G	2959	0	2881	98	0
2	L	2959	0	2881	94	0
2	R	2959	0	2881	98	0
3	C	1042	0	1021	56	0
3	H	1032	0	1014	37	0
3	M	1042	0	1021	49	0
3	S	1042	0	1021	35	0
4	D	63	0	68	3	0
4	I	63	0	68	4	0
4	O	63	0	68	4	0
4	T	63	0	68	12	0
5	E	96	0	106	3	0
5	J	85	0	90	5	0
5	P	77	0	79	1	0
5	U	85	0	90	8	0
6	A	8	0	0	0	0
6	F	8	0	0	0	0
6	K	8	0	0	0	0
6	Q	8	0	0	0	0
7	A	26	0	19	6	0
7	F	26	0	19	2	0
7	K	26	0	19	1	0
7	Q	26	0	19	1	0
All	All	35028	0	34038	1288	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 1288 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Q:432:TRP:CH2	1:Q:461:LYS:HG3	1.65	1.32
1:Q:318:LYS:NZ	1:Q:324:CYS:SG	2.26	1.08
1:Q:432:TRP:HD1	1:Q:469:PHE:CD1	1.75	1.05
1:Q:432:TRP:HH2	1:Q:461:LYS:CG	1.70	1.04
1:A:430:VAL:HG21	1:A:465:GLN:HE21	1.19	1.03

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:568:THR:OG1	1:Q:561:ARG:NH1[3_644]	2.11	0.09
1:F:561:ARG:NH1	1:K:568:THR:OG1[2_545]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	569/746 (76%)	534 (94%)	34 (6%)	1 (0%)	52	86
1	F	558/746 (75%)	526 (94%)	31 (6%)	1 (0%)	52	86
1	K	552/746 (74%)	527 (96%)	23 (4%)	2 (0%)	39	78
1	Q	555/746 (74%)	527 (95%)	27 (5%)	1 (0%)	52	86
2	B	363/367 (99%)	348 (96%)	15 (4%)	0	100	100
2	G	363/367 (99%)	355 (98%)	8 (2%)	0	100	100
2	L	363/367 (99%)	354 (98%)	9 (2%)	0	100	100
2	R	363/367 (99%)	352 (97%)	11 (3%)	0	100	100
3	C	123/129 (95%)	120 (98%)	3 (2%)	0	100	100
3	H	122/129 (95%)	119 (98%)	2 (2%)	1 (1%)	24	64
3	M	123/129 (95%)	118 (96%)	4 (3%)	1 (1%)	24	64
3	S	123/129 (95%)	121 (98%)	2 (2%)	0	100	100
4	D	7/13 (54%)	7 (100%)	0	0	100	100
4	I	7/13 (54%)	7 (100%)	0	0	100	100
4	O	7/13 (54%)	7 (100%)	0	0	100	100
4	T	7/13 (54%)	7 (100%)	0	0	100	100
5	E	8/12 (67%)	8 (100%)	0	0	100	100
5	J	7/12 (58%)	7 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	P	6/12 (50%)	6 (100%)	0	0	100	100
5	U	7/12 (58%)	6 (86%)	1 (14%)	0	100	100
All	All	4233/5068 (84%)	4056 (96%)	170 (4%)	7 (0%)	52	86

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	68	VAL
1	K	66	GLN
1	F	139	LEU
3	M	582	ASP
1	K	265	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	520/667 (78%)	496 (95%)	24 (5%)	33	71
1	F	507/667 (76%)	486 (96%)	21 (4%)	37	74
1	K	503/667 (75%)	481 (96%)	22 (4%)	35	72
1	Q	505/667 (76%)	485 (96%)	20 (4%)	38	75
2	B	328/329 (100%)	322 (98%)	6 (2%)	66	89
2	G	328/329 (100%)	319 (97%)	9 (3%)	52	84
2	L	328/329 (100%)	326 (99%)	2 (1%)	90	97
2	R	328/329 (100%)	327 (100%)	1 (0%)	94	98
3	C	119/121 (98%)	118 (99%)	1 (1%)	86	95
3	H	118/121 (98%)	115 (98%)	3 (2%)	55	85
3	M	119/121 (98%)	117 (98%)	2 (2%)	68	90
3	S	119/121 (98%)	118 (99%)	1 (1%)	86	95
4	D	6/7 (86%)	6 (100%)	0	100	100
4	I	6/7 (86%)	6 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	O	6/7 (86%)	6 (100%)	0	100	100
4	T	6/7 (86%)	6 (100%)	0	100	100
5	E	8/9 (89%)	6 (75%)	2 (25%)	1	3
5	J	7/9 (78%)	7 (100%)	0	100	100
5	P	6/9 (67%)	6 (100%)	0	100	100
5	U	7/9 (78%)	7 (100%)	0	100	100
All	All	3874/4532 (86%)	3760 (97%)	114 (3%)	50	82

5 of 114 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	663	CYS
3	H	579	MET
1	Q	525	HIS
1	F	668	ASN
2	G	223	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 102 such sidechains are listed below:

Mol	Chain	Res	Type
2	G	286	ASN
1	K	326	GLN
2	R	286	ASN
2	G	349	ASN
5	J	10	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	M3L	E	7	5	9,11,12	0.48	0	12,14,16	1.31	2 (16%)
5	M3L	J	7	5	9,11,12	0.52	0	12,14,16	1.29	1 (8%)
5	M3L	P	7	5	9,11,12	0.45	0	12,14,16	0.84	0
5	M3L	U	7	5	9,11,12	0.49	0	12,14,16	1.50	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	M3L	E	7	5	-	0/8/10/12	0/0/0/0
5	M3L	J	7	5	-	0/8/10/12	0/0/0/0
5	M3L	P	7	5	-	0/8/10/12	0/0/0/0
5	M3L	U	7	5	-	0/8/10/12	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	U	7	M3L	CM3-NZ-CM1	-3.36	100.27	108.96
5	E	7	M3L	CM3-NZ-CM1	-3.27	100.49	108.96
5	J	7	M3L	CM2-NZ-CM1	-3.25	100.56	108.96
5	E	7	M3L	CM2-NZ-CM1	2.00	114.14	108.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	7	M3L	1	0
5	J	7	M3L	2	0
5	U	7	M3L	1	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 36 ligands modelled in this entry, 32 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	SAH	A	1009	-	22,28,28	1.04	2 (9%)	18,40,40	2.87	5 (27%)
7	SAH	F	1009	-	22,28,28	1.12	2 (9%)	18,40,40	2.81	5 (27%)
7	SAH	K	1009	-	22,28,28	1.07	2 (9%)	18,40,40	3.03	4 (22%)
7	SAH	Q	1009	-	22,28,28	1.06	2 (9%)	18,40,40	2.79	4 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	SAH	A	1009	-	-	0/7/31/31	0/3/3/3
7	SAH	F	1009	-	-	0/7/31/31	0/3/3/3
7	SAH	K	1009	-	-	0/7/31/31	0/3/3/3
7	SAH	Q	1009	-	-	0/7/31/31	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1009	SAH	C2-N1	2.10	1.38	1.33
7	Q	1009	SAH	C2-N1	2.15	1.38	1.33
7	K	1009	SAH	C2-N1	2.22	1.38	1.33
7	F	1009	SAH	C2-N1	2.67	1.39	1.33
7	A	1009	SAH	C2-N3	3.22	1.37	1.32

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1009	SAH	N3-C2-N1	-10.58	120.56	128.87
7	K	1009	SAH	N3-C2-N1	-10.57	120.57	128.87
7	Q	1009	SAH	N3-C2-N1	-9.53	121.38	128.87
7	F	1009	SAH	N3-C2-N1	-9.16	121.67	128.87
7	K	1009	SAH	C5'-SD-CG	-5.60	85.42	102.42

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1009	SAH	6	0
7	F	1009	SAH	2	0
7	K	1009	SAH	1	0
7	Q	1009	SAH	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	581/746 (77%)	0.60	52 (8%) 12 6	40, 84, 159, 196	0
1	F	568/746 (76%)	0.67	58 (10%) 9 5	40, 92, 162, 224	0
1	K	562/746 (75%)	0.79	75 (13%) 4 2	43, 92, 165, 189	0
1	Q	565/746 (75%)	0.70	63 (11%) 7 4	45, 86, 165, 202	0
2	B	365/367 (99%)	0.40	11 (3%) 54 33	51, 79, 121, 171	0
2	G	365/367 (99%)	0.37	14 (3%) 44 26	49, 76, 115, 177	0
2	L	365/367 (99%)	0.29	4 (1%) 82 65	41, 65, 103, 140	0
2	R	365/367 (99%)	0.28	5 (1%) 78 59	35, 61, 92, 147	0
3	C	125/129 (96%)	0.47	5 (4%) 42 25	48, 72, 148, 172	0
3	H	124/129 (96%)	0.51	8 (6%) 22 12	58, 89, 143, 187	0
3	M	125/129 (96%)	0.74	11 (8%) 12 6	56, 92, 135, 158	0
3	S	125/129 (96%)	0.66	12 (9%) 10 5	54, 91, 139, 159	0
4	D	9/13 (69%)	0.16	0 100 100	67, 86, 111, 118	0
4	I	9/13 (69%)	0.16	0 100 100	62, 81, 103, 113	0
4	O	9/13 (69%)	1.70	2 (22%) 1 1	68, 76, 88, 93	9 (100%)
4	T	9/13 (69%)	1.72	2 (22%) 1 1	70, 73, 81, 94	9 (100%)
5	E	10/12 (83%)	0.34	1 (10%) 9 5	67, 89, 139, 140	0
5	J	9/12 (75%)	0.03	0 100 100	63, 82, 127, 133	0
5	P	8/12 (66%)	0.52	1 (12%) 5 2	69, 84, 117, 140	0
5	U	9/12 (75%)	0.87	0 100 100	71, 95, 133, 134	0
All	All	4307/5068 (84%)	0.56	324 (7%) 17 9	35, 79, 155, 224	18 (0%)

The worst 5 of 324 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	311	THR	13.6
1	Q	314	ALA	12.5
1	Q	235	GLY	11.8
1	F	74	VAL	10.5
1	F	75	SER	10.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	M3L	P	7	12/13	0.95	0.26	-	39,57,65,71	0
5	M3L	E	7	12/13	0.97	0.26	-	40,58,78,90	0
5	M3L	U	7	12/13	0.97	0.24	-	34,59,68,68	0
5	M3L	J	7	12/13	0.98	0.22	-	41,52,63,65	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	ZN	K	1004	1/1	0.98	0.19	2.68	70,70,70,70	0
6	ZN	F	1003	1/1	0.98	0.21	2.57	65,65,65,65	0
6	ZN	A	1004	1/1	0.98	0.24	2.32	63,63,63,63	0
6	ZN	Q	1007	1/1	0.99	0.22	2.29	54,54,54,54	0
6	ZN	K	1007	1/1	0.98	0.21	2.17	54,54,54,54	0
6	ZN	Q	1004	1/1	0.99	0.18	2.10	77,77,77,77	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	ZN	F	1004	1/1	0.99	0.21	1.68	65,65,65,65	0
6	ZN	Q	1003	1/1	0.98	0.18	1.42	69,69,69,69	0
6	ZN	Q	1008	1/1	0.99	0.21	1.34	50,50,50,50	0
6	ZN	F	1008	1/1	0.98	0.21	1.18	54,54,54,54	0
6	ZN	A	1008	1/1	0.99	0.21	1.16	58,58,58,58	0
6	ZN	F	1001	1/1	0.99	0.20	0.89	63,63,63,63	0
6	ZN	F	1007	1/1	0.99	0.21	0.81	54,54,54,54	0
6	ZN	A	1003	1/1	0.99	0.22	0.74	51,51,51,51	0
6	ZN	K	1006	1/1	0.98	0.21	0.74	63,63,63,63	0
6	ZN	K	1001	1/1	0.97	0.22	0.73	69,69,69,69	0
6	ZN	K	1003	1/1	0.98	0.17	0.57	66,66,66,66	0
6	ZN	A	1007	1/1	0.98	0.21	0.57	52,52,52,52	0
6	ZN	K	1008	1/1	0.98	0.21	0.44	59,59,59,59	0
6	ZN	F	1005	1/1	0.99	0.20	0.41	67,67,67,67	0
6	ZN	A	1005	1/1	0.99	0.21	0.38	56,56,56,56	0
6	ZN	Q	1006	1/1	0.98	0.22	0.38	66,66,66,66	0
6	ZN	Q	1005	1/1	0.99	0.19	0.20	72,72,72,72	0
6	ZN	F	1006	1/1	0.99	0.20	0.08	60,60,60,60	0
6	ZN	A	1006	1/1	0.99	0.20	0.07	58,58,58,58	0
6	ZN	Q	1001	1/1	0.99	0.21	0.04	60,60,60,60	0
6	ZN	K	1005	1/1	0.99	0.17	-0.00	74,74,74,74	0
6	ZN	A	1001	1/1	0.99	0.20	-0.08	54,54,54,54	0
7	SAH	Q	1009	26/26	0.88	0.21	-0.34	55,77,177,321	0
7	SAH	F	1009	26/26	0.92	0.20	-0.38	40,75,86,146	0
7	SAH	A	1009	26/26	0.94	0.21	-0.39	58,77,90,93	0
7	SAH	K	1009	26/26	0.89	0.21	-0.40	60,74,184,324	0
6	ZN	Q	1002	1/1	0.99	0.15	-0.55	128,128,128,128	0
6	ZN	F	1002	1/1	0.98	0.16	-0.94	110,110,110,110	0
6	ZN	A	1002	1/1	0.98	0.16	-1.13	64,64,64,64	0
6	ZN	K	1002	1/1	0.92	0.12	-1.33	122,122,122,122	0

6.5 Other polymers ⓘ

There are no such residues in this entry.