



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:39 AM GMT

PDB ID : 2HZS
Title : Structure of the Mediator head submodule Med8C/18/20
Authors : Lariviere, L.; Geiger, S.; Hoepfner, S.; Rother, S.; Straesser, K.; Cramer, P.
Deposited on : 2006-08-09
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	<div><div>17%</div><div><div></div><div>62%</div><div>30%</div><div>5%</div><div>.</div></div></div>
1	C	209	<div><div>15%</div><div><div></div><div>61%</div><div>32%</div><div>.</div><div>.</div></div></div>
1	E	209	<div><div>11%</div><div><div></div><div>62%</div><div>32%</div><div>.</div><div>.</div></div></div>
1	G	209	<div><div>19%</div><div><div></div><div>63%</div><div>30%</div><div>.</div><div>.</div></div></div>
2	B	306	<div><div>2%</div><div><div></div><div>50%</div><div>24%</div><div>.</div><div>.</div><div>23%</div></div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	306	<div><div><div></div><div></div><div></div><div></div></div><div>5%49%19%30%</div></div>
2	F	306	<div><div><div></div><div></div><div></div><div></div></div><div>4%46%28%23%</div></div>
2	H	306	<div><div><div></div><div></div><div></div><div></div></div><div>4%50%20%5%25%</div></div>
3	I	27	<div><div><div></div><div></div><div></div><div></div></div><div>4%63%19%19%</div></div>
3	J	27	<div><div><div></div><div></div><div></div><div></div></div><div>56%15%30%</div></div>
3	K	27	<div><div><div></div><div></div><div></div><div></div></div><div>52%7%41%</div></div>
3	L	27	<div><div><div></div><div></div><div></div><div></div></div><div>56%7%37%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14086 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA polymerase II mediator complex subunit 20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	0	0
			1560	987	263	305	5			
1	C	202	Total	C	N	O	S	0	0	0
			1554	984	262	303	5			
1	E	203	Total	C	N	O	S	0	0	0
			1560	987	263	305	5			
1	G	203	Total	C	N	O	S	0	0	0
			1560	987	263	305	5			

- Molecule 2 is a protein called RNA polymerase II mediator complex subunit 18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	235	Total	C	N	O	S	0	0	0
			1844	1175	302	358	9			
2	D	214	Total	C	N	O	S	0	0	0
			1671	1065	273	326	7			
2	F	235	Total	C	N	O	S	0	0	0
			1829	1156	298	367	8			
2	H	230	Total	C	N	O	S	0	0	0
			1803	1146	297	351	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	274	VAL	ALA	ENGINEERED	UNP P32585
D	274	VAL	ALA	ENGINEERED	UNP P32585
F	274	VAL	ALA	ENGINEERED	UNP P32585
H	274	VAL	ALA	ENGINEERED	UNP P32585

- Molecule 3 is a protein called RNA polymerase II mediator complex subunit 8.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	I	22	Total	C	N	O	0	0	0
			192	123	37	32			
3	J	19	Total	C	N	O	0	0	0
			156	102	25	29			
3	K	16	Total	C	N	O	0	0	0
			131	87	19	25			
3	L	17	Total	C	N	O	0	0	0
			141	93	22	26			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	211	HIS	-	EXPRESSION TAG	UNP P38304
I	212	HIS	-	EXPRESSION TAG	UNP P38304
I	213	HIS	-	EXPRESSION TAG	UNP P38304
I	214	HIS	-	EXPRESSION TAG	UNP P38304
I	215	HIS	-	EXPRESSION TAG	UNP P38304
I	216	HIS	-	EXPRESSION TAG	UNP P38304
J	211	HIS	-	EXPRESSION TAG	UNP P38304
J	212	HIS	-	EXPRESSION TAG	UNP P38304
J	213	HIS	-	EXPRESSION TAG	UNP P38304
J	214	HIS	-	EXPRESSION TAG	UNP P38304
J	215	HIS	-	EXPRESSION TAG	UNP P38304
J	216	HIS	-	EXPRESSION TAG	UNP P38304
K	211	HIS	-	EXPRESSION TAG	UNP P38304
K	212	HIS	-	EXPRESSION TAG	UNP P38304
K	213	HIS	-	EXPRESSION TAG	UNP P38304
K	214	HIS	-	EXPRESSION TAG	UNP P38304
K	215	HIS	-	EXPRESSION TAG	UNP P38304
K	216	HIS	-	EXPRESSION TAG	UNP P38304
L	211	HIS	-	EXPRESSION TAG	UNP P38304
L	212	HIS	-	EXPRESSION TAG	UNP P38304
L	213	HIS	-	EXPRESSION TAG	UNP P38304
L	214	HIS	-	EXPRESSION TAG	UNP P38304
L	215	HIS	-	EXPRESSION TAG	UNP P38304
L	216	HIS	-	EXPRESSION TAG	UNP P38304

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	O	0	0
			3	3		
4	B	34	Total	O	0	0
			34	34		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total 2	O 2	0	0
4	D	12	Total 12	O 12	0	0
4	E	2	Total 2	O 2	0	0
4	F	13	Total 13	O 13	0	0
4	H	15	Total 15	O 15	0	0
4	I	4	Total 4	O 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA polymerase II mediator complex subunit 20




- Molecule 1: RNA polymerase II mediator complex subunit 20




- Molecule 1: RNA polymerase II mediator complex subunit 20







• Molecule 1: RNA polymerase II mediator complex subunit 20



• Molecule 2: RNA polymerase II mediator complex subunit 18



• Molecule 2: RNA polymerase II mediator complex subunit 18



- Chain F:
-
- | Amino Acid | Frequency (%) |
|------------|---------------|
| V2 | 46% |
| Q3 | 46% |
| Q4 | 46% |
| L6 | 46% |
| S6 | 46% |
| L7 | 46% |
| S7 | 46% |
| F8 | 46% |
| I11 | 46% |
| G12 | 46% |
| D13 | 46% |
| L19 | 46% |
| T25 | 46% |
| I26 | 46% |
| S27 | 46% |
| P31 | 46% |
| V40 | 46% |
| S46 | 46% |
| Y47 | 46% |
| D48 | 46% |
| V49 | 46% |
| E50 | 46% |
| N51 | 46% |
| V52 | 46% |
| N53 | 46% |
| S54 | 46% |
| R55 | 46% |
| N56 | 46% |
| Q57 | 46% |
| L58 | 46% |
| V59 | 46% |
| E60 | 46% |
| R63 | 46% |
| I64 | 46% |
| K65 | 46% |
| L66 | 46% |
| S67 | 46% |
| K68 | 46% |
| E69 | 46% |
| V70 | 46% |
| P71 | 46% |
| Y74 | 46% |
| L75 | 46% |
| I76 | 46% |
| D77 | 46% |
| E78 | 46% |
| THR | 28% |
| MET | 28% |
| ASN | 28% |
| ASP | 28% |
| LYS | 28% |
| P70 | 28% |
| I71 | 28% |
| D143 | 4% |
| D144 | 4% |
| L145 | 4% |

- Chain H:
-
- 4% 50% 20% 5% 25%
- T271 E272 L275 K279 K280 Y285 I286 F287 L288 S295 S298 A301 H11 Q3 Q4 L5 S6 L7 I11 Y16 I20 S21 T22 I26 S27 P31 W41 M44 D48 V49 E50 N51 V52 N53 S54 R55 N56 Q57 L58 V59 R63 L66 T79 N80 M81 D82 R83 I89 L90 LYS SER PHE THR ASN ASP LYS PRO LEU ASN S160 V270

- Chain I:
-
- | Amino Acid Type | Percentage |
|-----------------|------------|
| 4% | 4% |
| 63% | 63% |
| 19% | 19% |
| 19% | 19% |
- Legend:
- | Amino Acid Type |
|-----------------|
| SER |
| LYS |
| PRO |
| SER |
| LYS |
| P195 |
| K203 |
| T207 |
| G208 |
| E209 |
| K210 |
| H211 |
| H215 |
| H216 |

- Chain J:  56% 15% 30%
- | Label | Color |
|-------|--------|
| SER | Grey |
| LYS | Grey |
| PRO | Grey |
| S193 | Green |
| K194 | Yellow |
| P195 | Yellow |
| V201 | Yellow |
| F206 | Yellow |
| H211 | Green |
| HIS | Grey |
| HIS | Grey |
| HIS | Grey |
| HIS | Grey |
| HIS | Grey |

- 



- Molecule 3: RNA polymerase II mediator complex subunit 8



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.36Å 115.76Å 129.16Å 90.00° 98.49° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 19.79 – 2.67	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.70) 99.1 (19.79-2.67)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.67Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.236 , 0.272 0.232 , 0.270	Depositor DCC
R_{free} test set	3047 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	57.6	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 61527 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14086	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/1584	0.87	8/2145 (0.4%)
1	C	0.38	0/1577	0.63	1/2134 (0.0%)
1	E	0.35	0/1584	0.64	1/2145 (0.0%)
1	G	0.36	0/1584	0.59	0/2145
2	B	0.46	0/1876	0.73	2/2541 (0.1%)
2	D	0.44	0/1700	0.68	1/2304 (0.0%)
2	F	0.45	0/1861	0.70	0/2526
2	H	0.45	0/1835	0.70	0/2486
3	I	0.45	0/201	0.60	0/269
3	J	0.49	0/160	0.59	0/214
3	K	0.55	0/134	0.65	0/179
3	L	0.52	0/145	0.59	0/194
All	All	0.42	0/14241	0.69	13/19282 (0.1%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	138	ARG	NE-CZ-NH1	-14.18	113.21	120.30
1	A	40	ARG	NE-CZ-NH2	-13.12	113.74	120.30
1	A	138	ARG	NE-CZ-NH2	12.83	126.72	120.30
1	A	40	ARG	NE-CZ-NH1	9.33	124.97	120.30
2	B	303	GLY	N-CA-C	-7.56	94.19	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1560	0	1576	70	0
1	C	1554	0	1573	66	0
1	E	1560	0	1576	61	0
1	G	1560	0	1576	52	0
2	B	1844	0	1851	79	0
2	D	1671	0	1676	53	1
2	F	1829	0	1811	87	1
2	H	1803	0	1809	67	0
3	I	192	0	168	5	0
3	J	156	0	150	5	0
3	K	131	0	126	2	0
3	L	141	0	133	2	0
4	A	3	0	0	0	0
4	B	34	0	0	2	0
4	C	2	0	0	0	0
4	D	12	0	0	0	0
4	E	2	0	0	0	0
4	F	13	0	0	1	0
4	H	15	0	0	0	0
4	I	4	0	0	0	0
All	All	14086	0	14025	522	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 522 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:LEU:HD11	1:A:152:LEU:HD12	1.35	1.06
1:G:142:LEU:HD11	1:G:152:LEU:HD12	1.36	1.05
2:F:265:ILE:HD12	2:F:265:ILE:H	1.25	1.02
1:C:142:LEU:HD11	1:C:152:LEU:HD12	1.37	1.01
1:E:142:LEU:HD11	1:E:152:LEU:HD12	1.37	1.00

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:25:THR:OG1	2:F:25:THR:OG1[2_646]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/209 (95%)	180 (90%)	17 (8%)	2 (1%)	19	45
1	C	196/209 (94%)	174 (89%)	18 (9%)	4 (2%)	9	24
1	E	199/209 (95%)	180 (90%)	18 (9%)	1 (0%)	34	63
1	G	199/209 (95%)	176 (88%)	21 (11%)	2 (1%)	19	45
2	B	229/306 (75%)	212 (93%)	12 (5%)	5 (2%)	8	22
2	D	208/306 (68%)	196 (94%)	10 (5%)	2 (1%)	19	45
2	F	231/306 (76%)	209 (90%)	19 (8%)	3 (1%)	15	37
2	H	226/306 (74%)	208 (92%)	10 (4%)	8 (4%)	4	10
3	I	20/27 (74%)	20 (100%)	0	0	100	100
3	J	17/27 (63%)	14 (82%)	3 (18%)	0	100	100
3	K	14/27 (52%)	13 (93%)	1 (7%)	0	100	100
3	L	15/27 (56%)	14 (93%)	1 (7%)	0	100	100
All	All	1753/2168 (81%)	1596 (91%)	130 (7%)	27 (2%)	13	32

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	83	LYS
2	B	84	PRO
2	F	56	ASN
2	F	261	ARG
2	H	56	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/177 (97%)	161 (94%)	11 (6%)	22	47
1	C	172/177 (97%)	163 (95%)	9 (5%)	29	58
1	E	172/177 (97%)	162 (94%)	10 (6%)	25	52
1	G	172/177 (97%)	160 (93%)	12 (7%)	19	42
2	B	213/280 (76%)	199 (93%)	14 (7%)	21	45
2	D	193/280 (69%)	179 (93%)	14 (7%)	17	39
2	F	212/280 (76%)	197 (93%)	15 (7%)	18	41
2	H	208/280 (74%)	193 (93%)	15 (7%)	18	41
3	I	21/26 (81%)	21 (100%)	0	100	100
3	J	18/26 (69%)	18 (100%)	0	100	100
3	K	15/26 (58%)	15 (100%)	0	100	100
3	L	16/26 (62%)	16 (100%)	0	100	100
All	All	1584/1932 (82%)	1484 (94%)	100 (6%)	22	48

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	246	ILE
1	E	181	GLU
2	H	233	GLN
2	D	275	LEU
1	E	52	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	277	ASN
1	E	173	HIS
2	H	243	ASN
1	E	68	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	195	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/209 (97%)	0.90	35 (17%) 2 1	35, 77, 120, 127	0
1	C	202/209 (96%)	0.83	31 (15%) 3 2	34, 81, 121, 127	0
1	E	203/209 (97%)	0.64	24 (11%) 6 5	35, 74, 117, 122	0
1	G	203/209 (97%)	0.85	39 (19%) 2 1	34, 79, 118, 122	0
2	B	235/306 (76%)	-0.07	7 (2%) 54 54	28, 45, 72, 111	0
2	D	214/306 (69%)	0.05	15 (7%) 19 17	29, 49, 100, 109	0
2	F	235/306 (76%)	0.01	13 (5%) 29 27	29, 48, 91, 108	0
2	H	230/306 (75%)	0.03	11 (4%) 34 33	30, 47, 92, 104	0
3	I	22/27 (81%)	0.13	1 (4%) 37 36	35, 44, 61, 82	0
3	J	19/27 (70%)	0.12	0 100 100	40, 50, 83, 93	0
3	K	16/27 (59%)	-0.28	0 100 100	37, 46, 64, 69	0
3	L	17/27 (62%)	-0.19	0 100 100	36, 46, 70, 85	0
All	All	1799/2168 (82%)	0.36	176 (9%) 10 7	28, 57, 115, 127	0

The worst 5 of 176 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	65	HIS	9.9
1	C	65	HIS	8.0
1	E	65	HIS	7.4
2	D	50	GLU	6.1
1	A	181	GLU	6.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.