



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:19 AM GMT

PDB ID : 3I24
Title : Crystal Structure of a HIT family hydrolase protein from *Vibrio fischeri*.
Northeast Structural Genomics Consortium target id VfR176
Authors : Seetharaman, J.; Abashidze, M.; Forouhar, F; Janjua, H.; Xiao, R.; Ciccocanti, C.; Foote, E.L.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2009-06-29
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	<div> <div>2%</div> <div>79%</div> <div>17%</div> <div>.</div> </div>
1	B	149	<div> <div>6%</div> <div>77%</div> <div>16%</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NA	A	153	-	-	-	X
2	NA	A	158	-	-	-	X
2	NA	A	159	-	-	-	X
2	NA	B	156	-	-	-	X
2	NA	B	157	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2679 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIT family hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	143	Total	C	N	O	S	0	0	0
			1135	727	201	204	3			
1	B	139	Total	C	N	O	S	0	0	0
			1105	709	195	198	3			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	142	LEU	-	EXPRESSION TAG	UNP Q5E3V1
A	143	GLU	-	EXPRESSION TAG	UNP Q5E3V1
A	144	HIS	-	EXPRESSION TAG	UNP Q5E3V1
A	145	HIS	-	EXPRESSION TAG	UNP Q5E3V1
A	146	HIS	-	EXPRESSION TAG	UNP Q5E3V1
A	147	HIS	-	EXPRESSION TAG	UNP Q5E3V1
A	148	HIS	-	EXPRESSION TAG	UNP Q5E3V1
A	149	HIS	-	EXPRESSION TAG	UNP Q5E3V1
B	142	LEU	-	EXPRESSION TAG	UNP Q5E3V1
B	143	GLU	-	EXPRESSION TAG	UNP Q5E3V1
B	144	HIS	-	EXPRESSION TAG	UNP Q5E3V1
B	145	HIS	-	EXPRESSION TAG	UNP Q5E3V1
B	146	HIS	-	EXPRESSION TAG	UNP Q5E3V1
B	147	HIS	-	EXPRESSION TAG	UNP Q5E3V1
B	148	HIS	-	EXPRESSION TAG	UNP Q5E3V1
B	149	HIS	-	EXPRESSION TAG	UNP Q5E3V1

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	8	Total	Na	0	0
			8	8		
2	A	11	Total	Na	0	0
			11	11		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	207	Total 207	O 207	0	0
3	B	213	Total 213	O 213	0	0

- Molecule 1: HIT family hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.43Å 69.31Å 49.08Å 90.00° 109.44° 90.00°	Depositor
Resolution (Å)	27.74 – 1.50 27.74 – 1.50	Depositor EDS
% Data completeness (in resolution range)	90.3 (27.74-1.50) 91.4 (27.74-1.50)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.60 (at 1.50Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.197 , 0.215 0.197 , 0.216	Depositor DCC
R_{free} test set	1922 reflections (4.89%)	DCC
Wilson B-factor (Å ²)	9.3	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 79669 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2679	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/1159	0.62	1/1575 (0.1%)
1	B	0.28	0/1129	0.61	0/1535
All	All	0.28	0/2288	0.61	1/3110 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	7	PRO	N-CA-CB	5.28	109.64	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1135	0	1150	23	0
1	B	1105	0	1127	19	0
2	A	11	0	0	0	0
2	B	8	0	0	0	0
3	A	207	0	0	7	0
3	B	213	0	0	3	0
All	All	2679	0	2277	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34:TRP:HE1	1:A:93:HIS:HD2	1.23	0.84
1:B:42:GLU:HG2	1:B:43:GLU:HG3	1.61	0.83
1:B:86:LEU:H	1:B:86:LEU:HD23	1.54	0.72
1:A:9:LEU:HB2	3:A:215:HOH:O	1.92	0.70
1:A:30:ASP:HB2	1:A:117:ARG:HA	1.74	0.70
1:A:34:TRP:HE1	1:A:93:HIS:CD2	2.07	0.68
1:A:9:LEU:HD12	3:A:215:HOH:O	1.93	0.66
1:A:6:HIS:N	3:A:215:HOH:O	2.34	0.61
1:A:117:ARG:HD2	1:A:122:GLN:OE1	2.01	0.61
1:B:48:HIS:HD2	3:B:230:HOH:O	1.84	0.60
1:A:25:LEU:HG	3:A:228:HOH:O	2.05	0.57
1:B:14:ILE:HD11	1:B:122:GLN:NE2	2.20	0.56
1:B:6:HIS:HE1	1:B:8:ARG:HH11	1.54	0.55
1:A:41:ILE:HD13	1:A:54:GLN:NE2	2.24	0.52
1:B:32:GLY:HA3	1:B:33:PRO:C	2.31	0.51
1:B:79:ASN:ND2	1:B:97:ARG:HH11	2.09	0.50
1:A:79:ASN:ND2	1:A:97:ARG:HH11	2.11	0.49
1:B:60:LYS:HE3	3:B:369:HOH:O	2.13	0.49
1:A:41:ILE:HB	1:A:44:LEU:HD11	1.95	0.48
1:B:141:ARG:HG2	3:B:248:HOH:O	2.13	0.48
1:A:32:GLY:HA3	1:A:33:PRO:C	2.33	0.48
1:B:44:LEU:HD21	1:B:50:MET:CG	2.44	0.47
1:A:30:ASP:HB2	1:A:117:ARG:CA	2.44	0.46
1:A:68:LEU:HD23	1:A:68:LEU:C	2.36	0.46
1:A:92:ILE:N	1:A:92:ILE:HD12	2.31	0.46
1:A:23:LYS:HB3	3:A:228:HOH:O	2.15	0.46
1:B:18:ASN:HD21	1:B:23:LYS:NZ	2.15	0.44
1:A:116:ILE:HD11	3:A:248:HOH:O	2.16	0.44
1:B:8:ARG:HG3	1:B:8:ARG:HH11	1.83	0.43
1:B:92:ILE:HD12	1:B:92:ILE:N	2.33	0.43
1:A:75:PRO:HA	1:A:98:PHE:CE1	2.53	0.42
1:A:51:THR:OG1	1:A:54:GLN:HG3	2.20	0.42
1:B:75:PRO:HA	1:B:98:PHE:CE1	2.54	0.42
1:B:18:ASN:ND2	1:B:23:LYS:HG3	2.35	0.42
1:A:6:HIS:HB3	3:A:215:HOH:O	2.19	0.42
1:A:70:GLU:OE1	1:B:48:HIS:HE1	2.04	0.41
1:B:75:PRO:HB3	1:B:96:ALA:HB1	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:68:LEU:HD23	1:B:68:LEU:C	2.41	0.41
1:A:70:GLU:HA	1:A:75:PRO:HD2	2.03	0.40
1:B:86:LEU:HG	1:B:87:VAL:N	2.37	0.40
1:A:75:PRO:HB3	1:A:96:ALA:HB1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/149 (95%)	140 (99%)	1 (1%)	0	100	100
1	B	137/149 (92%)	135 (98%)	2 (2%)	0	100	100
All	All	278/298 (93%)	275 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/134 (94%)	125 (99%)	1 (1%)	86	70
1	B	124/134 (92%)	124 (100%)	0	100	100
All	All	250/268 (93%)	249 (100%)	1 (0%)	93	84

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	93	HIS
1	B	18	ASN
1	B	48	HIS
1	B	49	HIS
1	B	79	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 19 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	143/149 (95%)	0.22	3 (2%) 67 70	6, 9, 18, 26	0
1	B	139/149 (93%)	0.33	9 (6%) 22 23	6, 11, 21, 29	0
All	All	282/298 (94%)	0.27	12 (4%) 39 41	6, 10, 20, 29	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	86	LEU	5.6
1	A	118	ALA	5.1
1	B	141	ARG	3.4
1	A	119	GLN	3.1
1	B	43	GLU	2.9
1	B	113	THR	2.8
1	B	115	VAL	2.5
1	B	44	LEU	2.5
1	B	119	GLN	2.4
1	B	42	GLU	2.3
1	A	8	ARG	2.2
1	B	3	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NA	B	156	1/1	0.99	0.16	9.83	10,10,10,10	0
2	NA	A	158	1/1	0.96	0.19	5.85	9,9,9,9	0
2	NA	A	159	1/1	0.99	0.12	3.31	8,8,8,8	0
2	NA	B	157	1/1	0.98	0.20	2.96	9,9,9,9	0
2	NA	A	153	1/1	0.98	0.17	2.57	7,7,7,7	0
2	NA	B	155	1/1	0.99	0.17	1.89	8,8,8,8	0
2	NA	B	153	1/1	0.99	0.13	1.64	8,8,8,8	0
2	NA	B	151	1/1	0.99	0.11	1.08	7,7,7,7	0
2	NA	A	160	1/1	0.99	0.11	0.97	7,7,7,7	0
2	NA	A	156	1/1	0.99	0.11	0.71	8,8,8,8	0
2	NA	A	154	1/1	0.99	0.12	0.71	8,8,8,8	0
2	NA	A	155	1/1	0.99	0.10	0.30	8,8,8,8	0
2	NA	A	151	1/1	0.99	0.10	0.02	5,5,5,5	0
2	NA	B	150	1/1	0.99	0.09	-0.11	8,8,8,8	0
2	NA	B	152	1/1	0.98	0.10	-0.21	7,7,7,7	0
2	NA	A	157	1/1	1.00	0.08	-0.88	9,9,9,9	0
2	NA	B	154	1/1	0.99	0.07	-1.16	10,10,10,10	0
2	NA	A	152	1/1	0.99	0.07	-1.58	7,7,7,7	0
2	NA	A	150	1/1	0.99	0.07	-	5,5,5,5	0

6.5 Other polymers [i](#)

There are no such residues in this entry.