



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:21 AM GMT

PDB ID : 3I4U  
Title : Crystal Structure Analysis of a helicase associated domain  
Authors : Kudlinzki, D.; Ficner, R.  
Deposited on : 2009-07-02  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	<div> <div>9%</div> <div>81%</div> <div>7%</div> <div>11%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	11	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	14	-	-	-	X
3	GOL	A	16	-	-	-	X
3	GOL	A	17	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2223 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent RNA helicase DHX8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	1988	1270	350	351	17	0	5	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	935	MET	-	EXPRESSION TAG	UNP Q14562
A	936	GLY	-	EXPRESSION TAG	UNP Q14562
A	937	ASP	-	EXPRESSION TAG	UNP Q14562
A	938	ARG	-	EXPRESSION TAG	UNP Q14562
A	939	GLY	-	EXPRESSION TAG	UNP Q14562
A	940	PRO	-	EXPRESSION TAG	UNP Q14562
A	941	GLU	-	EXPRESSION TAG	UNP Q14562
A	942	PHE	-	EXPRESSION TAG	UNP Q14562
A	943	GLU	-	EXPRESSION TAG	UNP Q14562
A	944	LEU	-	EXPRESSION TAG	UNP Q14562
A	945	GLY	-	EXPRESSION TAG	UNP Q14562
A	946	THR	-	EXPRESSION TAG	UNP Q14562
A	947	ARG	-	EXPRESSION TAG	UNP Q14562
A	948	GLY	-	EXPRESSION TAG	UNP Q14562
A	949	SER	-	EXPRESSION TAG	UNP Q14562
A	1184	LEU	-	EXPRESSION TAG	UNP Q14562
A	1185	GLU	-	EXPRESSION TAG	UNP Q14562
A	1186	VAL	-	EXPRESSION TAG	UNP Q14562
A	1187	ASP	-	EXPRESSION TAG	UNP Q14562
A	1188	LEU	-	EXPRESSION TAG	UNP Q14562
A	1189	GLN	-	EXPRESSION TAG	UNP Q14562
A	1190	GLY	-	EXPRESSION TAG	UNP Q14562
A	1191	ASP	-	EXPRESSION TAG	UNP Q14562
A	1192	HIS	-	EXPRESSION TAG	UNP Q14562
A	1193	GLY	-	EXPRESSION TAG	UNP Q14562
A	1194	LEU	-	EXPRESSION TAG	UNP Q14562
A	1195	SER	-	EXPRESSION TAG	UNP Q14562

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1196	ALA	-	EXPRESSION TAG	UNP Q14562
A	1197	TRP	-	EXPRESSION TAG	UNP Q14562
A	1198	SER	-	EXPRESSION TAG	UNP Q14562
A	1199	HIS	-	EXPRESSION TAG	UNP Q14562
A	1200	PRO	-	EXPRESSION TAG	UNP Q14562
A	1201	GLN	-	EXPRESSION TAG	UNP Q14562
A	1202	PHE	-	EXPRESSION TAG	UNP Q14562
A	1203	GLU	-	EXPRESSION TAG	UNP Q14562
A	1204	LYS	-	EXPRESSION TAG	UNP Q14562

- Molecule 2 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Br 3 3	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	190	Total	O	0	0
			190	190		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP-dependent RNA helicase DHX8



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.85Å 78.85Å 88.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.48 – 2.10 29.48 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.5 (29.48-2.10) 98.6 (29.48-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.64 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.193 , 0.245 0.194 , 0.248	Depositor DCC
$R_{free}$ test set	843 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.7	Xtriage
Anisotropy	0.164	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 54.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 16688 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2223	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/2042	0.47	0/2750

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1988	0	2021	17	0
2	A	3	0	0	0	0
3	A	42	0	56	9	0
4	A	190	0	0	0	0
All	All	2223	0	2077	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1172:TRP:HE1	3:A:16:GOL:H2	1.52	0.74
1:A:1040:GLY:HA2	1:A:1141:LEU:HD13	1.79	0.65
1:A:1088:ARG:HD2	3:A:12:GOL:H2	1.83	0.60
1:A:1183:VAL:O	1:A:1183:VAL:HG12	2.02	0.60
1:A:1055:ASN:HD21	3:A:14:GOL:C2	2.20	0.54
1:A:1055:ASN:HD21	3:A:14:GOL:H2	1.73	0.53
1:A:1080:LYS:HB2	3:A:11:GOL:H2	1.90	0.53
1:A:1018:PHE:CZ	1:A:1048:VAL:HG21	2.47	0.50
1:A:1084:GLY:HA3	3:A:13:GOL:C1	2.43	0.49
1:A:1122:GLU:HG2	1:A:1134:TYR:HE1	1.78	0.49
1:A:1039:GLU:HB2	1:A:1043:LEU:HD12	1.96	0.48
1:A:1172:TRP:NE1	3:A:16:GOL:H2	2.26	0.45
1:A:1084:GLY:HA3	3:A:13:GOL:H12	1.98	0.45
1:A:1067:ILE:HG21	1:A:1072:LEU:HD11	2.01	0.43
1:A:991:LYS:HE2	1:A:995[B]:MET:HG3	2.01	0.43
1:A:960:GLN:HG3	1:A:1113[B]:ARG:CZ	2.49	0.42
1:A:1084:GLY:HA3	3:A:13:GOL:H11	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/270 (90%)	233 (96%)	8 (3%)	1 (0%)	39	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1183	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	219/239 (92%)	216 (99%)	3 (1%)	74 80

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	978[A]	ARG
1	A	978[B]	ARG
1	A	1141	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1025	GLN
1	A	1055	ASN
1	A	1145	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	11	-	5,5,5	0.44	0	5,5,5	0.17	0
3	GOL	A	12	-	5,5,5	0.31	0	5,5,5	0.32	0
3	GOL	A	13	-	5,5,5	0.31	0	5,5,5	0.29	0
3	GOL	A	14	-	5,5,5	0.33	0	5,5,5	0.23	0
3	GOL	A	15	-	5,5,5	0.35	0	5,5,5	0.28	0
3	GOL	A	16	-	5,5,5	0.34	0	5,5,5	0.39	0
3	GOL	A	17	-	5,5,5	0.35	0	5,5,5	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	11	-	-	0/4/4/4	0/0/0/0
3	GOL	A	12	-	-	0/4/4/4	0/0/0/0
3	GOL	A	13	-	-	0/4/4/4	0/0/0/0
3	GOL	A	14	-	-	0/4/4/4	0/0/0/0
3	GOL	A	15	-	-	0/4/4/4	0/0/0/0
3	GOL	A	16	-	-	0/4/4/4	0/0/0/0
3	GOL	A	17	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	11	GOL	1	0
3	A	12	GOL	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	13	GOL	3	0
3	A	14	GOL	2	0
3	A	16	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	241/270 (89%)	0.57	24 (9%)	9 13	21, 33, 67, 91	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1194	LEU	9.3
1	A	1184	LEU	8.0
1	A	1183	VAL	6.2
1	A	1098	LYS	4.6
1	A	1193	GLY	4.4
1	A	948	GLY	4.3
1	A	1191	ASP	4.3
1	A	1016	ASN	3.2
1	A	1107	ILE	3.1
1	A	1022	LYS	2.9
1	A	1019	TYR	2.4
1	A	1121	GLN	2.4
1	A	1182	LYS	2.3
1	A	1015	GLN	2.3
1	A	1023	ASP	2.3
1	A	1127	LEU	2.2
1	A	1008	ILE	2.1
1	A	970	GLU	2.1
1	A	1129	ASP	2.1
1	A	1021	PRO	2.1
1	A	1070	ARG	2.1
1	A	1122	GLU	2.1
1	A	1012	LEU	2.0
1	A	969	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	17	6/6	0.74	0.22	8.88	60,61,61,61	0
3	GOL	A	11	6/6	0.81	0.28	6.00	39,42,43,44	0
3	GOL	A	16	6/6	0.62	0.24	4.80	59,59,60,61	0
3	GOL	A	14	6/6	0.78	0.18	2.14	49,52,53,53	0
3	GOL	A	15	6/6	0.90	0.17	0.98	53,54,54,54	0
3	GOL	A	12	6/6	0.88	0.31	-	47,47,47,48	0
2	BR	A	2	1/1	0.98	0.10	-	43,43,43,43	1
2	BR	A	1	1/1	1.00	0.07	-	38,38,38,38	0
2	BR	A	3	1/1	0.99	0.04	-	55,55,55,55	1
3	GOL	A	13	6/6	0.79	0.41	-	53,55,56,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.