



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:42 AM GMT

PDB ID : 2IBU
Title : Crystallographic and kinetic studies of human mitochondrial acetoacetyl-CoA thiolase (T2): the importance of potassium and chloride for its structure and function
Authors : Haapalainen, A.M.; Wierenga, R.K.
Deposited on : 2006-09-12
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	<div> <div>93%</div> <div>5% •</div> </div>
1	B	395	<div> <div>94%</div> <div>5% •</div> </div>
1	C	395	<div> <div>93%</div> <div>5% ••</div> </div>
1	D	395	<div> <div>94%</div> <div>6% •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	COA	A	6001	-	-	-	X
3	COA	B	6002	-	-	-	X
3	COA	C	6003	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 12936 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl-CoA acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	391	Total	C	N	O	S	0	6	0
			2891	1825	490	554	22			
1	B	391	Total	C	N	O	S	0	5	0
			2889	1825	490	553	21			
1	C	393	Total	C	N	O	S	0	2	0
			2895	1827	493	554	21			
1	D	393	Total	C	N	O	S	0	0	0
			2888	1820	493	554	21			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	MET	-	INITIATING METHIONINE	UNP P24752
A	34	ALA	VAL	ENGINEERED	UNP P24752
A	126	CSO	CYS	MODIFIED RESIDUE	UNP P24752
B	33	MET	-	INITIATING METHIONINE	UNP P24752
B	34	ALA	VAL	ENGINEERED	UNP P24752
B	126	CSO	CYS	MODIFIED RESIDUE	UNP P24752
C	33	MET	-	INITIATING METHIONINE	UNP P24752
C	34	ALA	VAL	ENGINEERED	UNP P24752
C	126	CSO	CYS	MODIFIED RESIDUE	UNP P24752
D	33	MET	-	INITIATING METHIONINE	UNP P24752
D	34	ALA	VAL	ENGINEERED	UNP P24752
D	126	CSO	CYS	MODIFIED RESIDUE	UNP P24752

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cl	0	0
			2	2		
2	D	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Cl	0	0
			1	1		

- Molecule 3 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	B	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	C	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	D	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	421	Total	O	0	0
			421	421		
5	B	381	Total	O	0	0
			381	381		
5	C	180	Total	O	0	0
			180	180		
5	D	189	Total	O	0	0
			189	189		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acetyl-CoA acetyltransferase




- Molecule 1: Acetyl-CoA acetyltransferase



- Molecule 1: Acetyl-CoA acetyltransferase



- Molecule 1: Acetyl-CoA acetyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.25Å 106.57Å 101.50Å 90.00° 102.90° 90.00°	Depositor
Resolution (Å)	19.46 – 1.90 19.45 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.46-1.90) 100.0 (19.45-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.166 , 0.198 0.166 , 0.199	Depositor DCC
R_{free} test set	6131 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	18.3	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 122616 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12936	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, GOL, CSO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/2949	0.62	0/3990
1	B	0.54	0/2943	0.62	0/3984
1	C	0.45	0/2937	0.60	0/3973
1	D	0.45	0/2922	0.58	0/3952
All	All	0.51	0/11751	0.61	0/15899

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2891	0	2981	13	0
1	B	2889	0	2982	11	0
1	C	2895	0	2988	13	0
1	D	2888	0	2976	11	0
2	A	2	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	48	0	32	0	0
3	B	48	0	32	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	48	0	32	0	0
3	D	48	0	32	0	0
4	A	6	0	8	0	0
5	A	421	0	0	1	0
5	B	381	0	0	2	0
5	C	180	0	0	0	0
5	D	189	0	0	0	0
All	All	12936	0	12063	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48[A]:THR:HG22	5:A:6121:HOH:O	1.82	0.78
1:C:308:PRO:HB2	1:C:427:LEU:HD23	1.80	0.63
1:C:402:LEU:HB2	1:C:426:LYS:HG3	1.81	0.63
1:D:163:MET:HE3	1:D:175:LEU:HD21	1.83	0.61
1:B:354:GLU:HB3	1:B:381:VAL:HG23	1.84	0.58
1:D:76:ILE:CD1	1:D:113:LEU:HD11	2.34	0.58
1:D:135:MET:HA	1:D:138:GLN:HE21	1.70	0.56
1:B:395:VAL:O	1:B:399[B]:THR:HG23	2.07	0.54
1:A:76:ILE:CD1	1:A:113:LEU:HD11	2.37	0.54
1:C:115:ILE:H	1:C:115:ILE:HD13	1.72	0.53
1:B:176:GLU:HG3	5:B:6151:HOH:O	2.10	0.51
1:A:336[A]:VAL:HG11	1:A:423:LEU:HB2	1.92	0.51
1:A:354:GLU:HB3	1:A:381:VAL:HG23	1.93	0.51
1:C:354:GLU:HB3	1:C:381:VAL:HG23	1.93	0.51
1:D:163:MET:CE	1:D:175:LEU:HD21	2.42	0.50
1:C:269:THR:HG21	1:C:276:GLY:C	2.32	0.50
1:A:137:SER:O	1:A:141[A]:MET:HG3	2.12	0.49
1:D:373:LYS:C	1:D:376:ILE:HD11	2.34	0.48
1:D:196:CYS:HB2	1:D:325:PHE:HE2	1.78	0.48
1:C:399:THR:O	1:C:426:LYS:HE3	2.14	0.48
1:C:354:GLU:HG2	1:C:360:VAL:HG21	1.94	0.48
1:B:309:LEU:O	1:B:426:LYS:HD2	2.15	0.47
1:D:51:PRO:HD3	1:D:236:ILE:O	2.16	0.46
1:B:349:MET:HG3	1:B:398:LEU:CD2	2.46	0.46
1:B:76:ILE:CD1	1:B:113[A]:LEU:HD11	2.46	0.46
1:B:354:GLU:HG2	1:B:360:VAL:HG21	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:354:GLU:HG2	1:A:360:VAL:HG21	1.98	0.45
1:A:306:VAL:O	1:A:308:PRO:HD3	2.17	0.45
1:A:135:MET:HA	1:A:138:GLN:HE21	1.81	0.45
1:C:402:LEU:CB	1:C:426:LYS:HG3	2.46	0.45
1:A:395:VAL:O	1:A:399[B]:THR:HG23	2.16	0.44
3:B:6002:COA:H52A	5:B:6340:HOH:O	2.18	0.44
1:B:51:PRO:HD3	1:B:236:ILE:O	2.18	0.44
1:D:207:ALA:O	1:D:211:GLN:HG3	2.17	0.44
1:C:370:ASP:HA	1:C:371:PRO:HD2	1.86	0.44
1:A:39:LEU:CD2	1:B:39:LEU:HD23	2.47	0.43
1:A:39:LEU:HD21	1:B:39:LEU:HD23	2.00	0.43
1:D:169:PRO:HD2	1:D:173:VAL:HG22	2.01	0.43
1:C:351:GLU:HG2	1:C:398:LEU:HD13	2.01	0.42
1:D:373:LYS:HA	1:D:376:ILE:HD11	2.02	0.42
1:C:307:THR:HA	1:C:308:PRO:HD3	1.83	0.42
1:C:116:SER:O	1:C:118:PRO:HD3	2.19	0.42
1:C:340:VAL:O	1:C:340:VAL:HG12	2.20	0.42
1:D:84:GLU:H	1:D:84:GLU:CD	2.24	0.41
1:A:314:ALA:HB3	1:A:336[A]:VAL:HG13	2.02	0.41
1:B:328:ALA:N	1:B:329:PRO:CD	2.84	0.41
1:A:72:ILE:HG12	1:A:149[A]:VAL:HG11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	394/395 (100%)	387 (98%)	6 (2%)	1 (0%)	46	35
1	B	393/395 (100%)	386 (98%)	6 (2%)	1 (0%)	46	35
1	C	392/395 (99%)	384 (98%)	7 (2%)	1 (0%)	46	35
1	D	390/395 (99%)	383 (98%)	6 (2%)	1 (0%)	46	35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1569/1580 (99%)	1540 (98%)	25 (2%)	4 (0%)	46 35

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125	VAL
1	B	125	VAL
1	C	125	VAL
1	D	125	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	311/308 (101%)	309 (99%)	2 (1%)	90 90
1	B	310/308 (101%)	309 (100%)	1 (0%)	94 95
1	C	309/308 (100%)	298 (96%)	11 (4%)	42 30
1	D	307/308 (100%)	302 (98%)	5 (2%)	70 66
All	All	1237/1232 (100%)	1218 (98%)	19 (2%)	70 69

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	VAL
1	A	175	LEU
1	B	389	MET
1	C	115	ILE
1	C	125	VAL
1	C	238	VAL
1	C	267	LEU
1	C	286	LEU
1	C	295	LEU
1	C	305	ASN
1	C	309	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	398	LEU
1	C	426	LYS
1	C	427	LEU
1	D	174	LYS
1	D	304	LEU
1	D	309	LEU
1	D	389	MET
1	D	427	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	GLN
1	A	138	GLN
1	B	93	ASN
1	B	138	GLN
1	B	199	ASN
1	C	93	ASN
1	D	93	ASN
1	D	138	GLN
1	D	164	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	126	1	3,6,7	0.36	0	1,6,8	2.71	1 (100%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSO	B	126	1	3,6,7	0.44	0	1,6,8	2.41	1 (100%)
1	CSO	C	126	1	3,6,7	0.65	0	1,6,8	2.14	1 (100%)
1	CSO	D	126	1	3,6,7	0.51	0	1,6,8	2.07	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	126	1	-	0/1/5/7	0/0/0/0
1	CSO	B	126	1	-	0/1/5/7	0/0/0/0
1	CSO	C	126	1	-	0/1/5/7	0/0/0/0
1	CSO	D	126	1	-	0/1/5/7	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	CSO	O-C-CA	-2.71	118.42	125.49
1	B	126	CSO	O-C-CA	-2.41	119.20	125.49
1	C	126	CSO	O-C-CA	-2.14	119.91	125.49
1	D	126	CSO	O-C-CA	-2.07	120.09	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	4001	-	5,5,5	0.26	0	5,5,5	0.28	0
3	COA	A	6001	-	40,50,50	1.65	6 (15%)	50,75,75	1.39	3 (6%)
3	COA	B	6002	-	40,50,50	1.62	5 (12%)	50,75,75	1.41	4 (8%)
3	COA	C	6003	-	40,50,50	1.65	6 (15%)	50,75,75	1.32	2 (4%)
3	COA	D	6004	-	40,50,50	1.65	6 (15%)	50,75,75	1.34	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	4001	-	-	0/4/4/4	0/0/0/0
3	COA	A	6001	-	-	0/44/64/64	0/3/3/3
3	COA	B	6002	-	-	0/44/64/64	0/3/3/3
3	COA	C	6003	-	-	0/44/64/64	0/3/3/3
3	COA	D	6004	-	-	0/44/64/64	0/3/3/3

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	6001	COA	C2A-N3A	2.02	1.35	1.32
3	D	6004	COA	O4B-C1B	2.16	1.43	1.41
3	C	6003	COA	O4B-C1B	2.18	1.44	1.41
3	D	6004	COA	C5A-C4A	2.68	1.46	1.40
3	A	6001	COA	C5A-C4A	2.77	1.46	1.40
3	C	6003	COA	C5A-C4A	2.78	1.46	1.40
3	B	6002	COA	C5A-C4A	2.79	1.46	1.40
3	B	6002	COA	P2A-O5A	3.51	1.69	1.54
3	C	6003	COA	P2A-O5A	3.59	1.70	1.54
3	B	6002	COA	P1A-O2A	3.61	1.70	1.54
3	A	6001	COA	P2A-O5A	3.61	1.70	1.54
3	A	6001	COA	P1A-O2A	3.62	1.70	1.54
3	D	6004	COA	P2A-O5A	3.68	1.70	1.54
3	D	6004	COA	P1A-O2A	3.69	1.70	1.54
3	C	6003	COA	P1A-O2A	3.69	1.70	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	6002	COA	P3B-O8A	4.42	1.70	1.54
3	C	6003	COA	P3B-O8A	4.47	1.70	1.54
3	D	6004	COA	P3B-O8A	4.48	1.70	1.54
3	A	6001	COA	P3B-O8A	4.49	1.70	1.54
3	C	6003	COA	P3B-O7A	5.96	1.70	1.51
3	D	6004	COA	P3B-O7A	5.97	1.70	1.51
3	B	6002	COA	P3B-O7A	6.03	1.71	1.51
3	A	6001	COA	P3B-O7A	6.05	1.71	1.51

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	6002	COA	N3A-C2A-N1A	-7.01	123.53	128.89
3	D	6004	COA	N3A-C2A-N1A	-6.62	123.83	128.89
3	C	6003	COA	N3A-C2A-N1A	-6.32	124.06	128.89
3	A	6001	COA	N3A-C2A-N1A	-6.11	124.22	128.89
3	C	6003	COA	P2A-O3A-P1A	-4.47	120.19	132.73
3	D	6004	COA	P2A-O3A-P1A	-4.01	121.46	132.73
3	A	6001	COA	C7P-C6P-C5P	-3.97	105.77	112.31
3	B	6002	COA	P2A-O3A-P1A	-3.80	122.05	132.73
3	A	6001	COA	P2A-O3A-P1A	-3.80	122.06	132.73
3	B	6002	COA	C1B-N9A-C4A	-2.06	123.84	126.94
3	B	6002	COA	C2A-N1A-C6A	2.14	122.59	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	6002	COA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/395 (98%)	-0.47	5 (1%) 79 82	9, 16, 31, 58	13 (3%)
1	B	390/395 (98%)	-0.38	7 (1%) 71 74	10, 18, 33, 50	10 (2%)
1	C	392/395 (99%)	0.03	13 (3%) 50 53	17, 29, 47, 68	6 (1%)
1	D	392/395 (99%)	0.07	17 (4%) 39 42	17, 29, 46, 68	8 (2%)
All	All	1564/1580 (98%)	-0.19	42 (2%) 58 61	9, 23, 43, 68	37 (2%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	244	GLY	6.1
1	C	35	SER	5.2
1	D	274	GLU	5.1
1	C	243	LYS	5.0
1	D	273	LYS	5.0
1	C	275	ASN	3.9
1	C	242	VAL	3.9
1	D	275	ASN	3.6
1	A	243	LYS	3.5
1	D	244	GLY	3.1
1	D	241	THR	3.0
1	D	246	PRO	3.0
1	C	36	LYS	2.9
1	C	245	GLN	2.8
1	D	261	PHE	2.7
1	D	233	ASN	2.6
1	A	245	GLN	2.6
1	B	244	GLY	2.6
1	A	273	LYS	2.5
1	A	244	GLY	2.5
1	D	243	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	242	VAL	2.5
1	B	243	LYS	2.4
1	D	268	LYS	2.4
1	B	246	PRO	2.3
1	D	403	LYS	2.3
1	C	246	PRO	2.3
1	C	241	THR	2.3
1	A	246	PRO	2.2
1	D	404	GLN	2.2
1	D	270	VAL	2.2
1	D	258	ARG	2.2
1	C	202	LYS	2.2
1	C	205	ASN	2.1
1	B	273	LYS	2.1
1	B	38[A]	THR	2.1
1	D	205	ASN	2.1
1	C	305	ASN	2.1
1	D	262	SER	2.1
1	D	245	GLN	2.1
1	C	274	GLU	2.0
1	B	245	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CSO	B	126	7/8	0.95	0.07	-	13,14,21,24	0
1	CSO	D	126	7/8	0.93	0.09	-	21,23,27,33	0
1	CSO	A	126	7/8	0.97	0.07	-	10,11,14,23	0
1	CSO	C	126	7/8	0.96	0.07	-	18,21,24,29	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	COA	B	6002	48/48	0.84	0.20	3.46	30,39,49,50	0
3	COA	A	6001	48/48	0.79	0.20	2.95	32,40,53,54	0
3	COA	C	6003	48/48	0.80	0.21	2.67	36,60,64,65	0
3	COA	D	6004	48/48	0.79	0.20	1.60	45,63,69,69	0
4	GOL	A	4001	6/6	0.90	0.12	1.09	29,30,32,32	0
2	CL	D	2003	1/1	1.00	0.05	-1.79	23,23,23,23	0
2	CL	C	2004	1/1	0.99	0.05	-1.93	25,25,25,25	0
2	CL	A	2001	1/1	1.00	0.06	-2.27	14,14,14,14	0
2	CL	A	2002	1/1	1.00	0.06	-3.73	14,14,14,14	0

6.5 Other polymers

There are no such residues in this entry.