



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:31 PM GMT

PDB ID : 4ILW
Title : Complex of matrix metalloproteinase-10 catalytic domain (MMP-10cd) with
tissue inhibitor of metalloproteinases-2 (TIMP-2)
Authors : Batra, J.; Soares, A.S.; Radisky, E.S.
Deposited on : 2013-01-01
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	<div> <div>5%</div> <div>70%</div> <div>16%</div> <div>6%</div> <div>6%</div> </div>
1	B	194	<div> <div>22%</div> <div>66%</div> <div>24%</div> <div>6%</div> </div>
2	D	165	<div> <div>5%</div> <div>79%</div> <div>13%</div> <div>6%</div> </div>
2	F	165	<div> <div>7%</div> <div>79%</div> <div>13%</div> <div>6%</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5564 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metalloproteinase inhibitor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1419	896	242	264	17			
1	B	182	Total	C	N	O	S	0	0	0
			1419	896	242	264	17			

- Molecule 2 is a protein called Stromelysin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	159	Total	C	N	O	S	0	2	0
			1278	821	207	247	3			
2	F	159	Total	C	N	O	S	0	2	0
			1278	821	207	247	3			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	2	Total	Zn	0	0
			2	2		
3	F	2	Total	Zn	0	0
			2	2		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	3	Total	Ca	0	0
			3	3		
4	F	3	Total	Ca	0	0
			3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	44	Total 44	O 44	0	0
5	B	37	Total 37	O 37	0	0
5	D	48	Total 48	O 48	0	0
5	F	31	Total 31	O 31	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Metalloproteinase inhibitor 2




• Molecule 1: Metalloproteinase inhibitor 2



• Molecule 2: Stromelysin-2



• Molecule 2: Stromelysin-2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	37.69Å 56.93Å 82.59Å 76.14° 79.84° 71.25°	Depositor
Resolution (Å)	40.58 – 2.10 40.55 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.7 (40.58-2.10) 91.4 (40.55-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.216 , 0.263 0.220 , 0.268	Depositor DCC
R_{free} test set	1782 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.628	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 35606 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5564	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.74	0/1452	0.88	3/1959 (0.2%)
1	B	0.69	0/1452	0.85	1/1959 (0.1%)
2	D	0.80	0/1324	0.84	1/1803 (0.1%)
2	F	0.63	0/1324	0.82	0/1803
All	All	0.72	0/5552	0.85	5/7524 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
2	D	0	2
2	F	0	1
All	All	0	7

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	34	ASP	CB-CG-OD2	-6.05	112.85	118.30
1	A	74	VAL	CB-CA-C	-5.68	100.61	111.40
2	D	199	ASP	CB-CG-OD2	5.67	123.40	118.30
1	B	16	ASP	CB-CA-C	-5.34	99.72	110.40
1	A	121	ARG	NE-CZ-NH1	5.17	122.88	120.30

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	171	SER	Peptide
1	A	32	GLY	Peptide
1	B	31	SER	Peptide
1	B	33	ASN	Peptide
2	D	239	TYR	Peptide
2	D	241	SER	Peptide
2	F	239	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1419	0	1388	39	0
1	B	1419	0	1388	35	0
2	D	1278	0	1196	18	0
2	F	1278	0	1196	23	0
3	D	2	0	0	0	0
3	F	2	0	0	0	0
4	D	3	0	0	0	0
4	F	3	0	0	0	0
5	A	44	0	0	3	0
5	B	37	0	0	6	0
5	D	48	0	0	4	0
5	F	31	0	0	2	0
All	All	5564	0	5168	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:VAL:HB	5:B:219:HOH:O	1.74	0.85
1:B:52:MET:SD	1:B:55:GLY:O	2.36	0.84
2:D:116:ARG:HB3	2:D:159:MET:CE	2.08	0.84
1:A:32:GLY:HA3	1:A:33:ASN:HB2	1.60	0.83
1:A:32:GLY:CA	1:A:33:ASN:HB2	2.10	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:106:PRO:HD2	5:D:404:HOH:O	1.80	0.81
1:B:121:ARG:HD2	5:B:201:HOH:O	1.79	0.81
2:F:116:ARG:HB3	2:F:159:MET:CE	2.13	0.78
1:B:121:ARG:NH2	1:B:178:TYR:O	2.17	0.77
2:D:243:THR:O	2:D:245:LEU:HD22	1.86	0.75
2:F:112:HIS:HB3	5:F:429:HOH:O	1.88	0.74
2:F:234:LEU:O	2:F:240:ASN:ND2	2.20	0.73
2:D:116:ARG:HB3	2:D:159:MET:HE1	1.71	0.73
1:B:119:ASN:O	5:B:233:HOH:O	2.05	0.73
2:D:120:TYR:HB3	5:D:426:HOH:O	1.87	0.72
1:B:32:GLY:HA2	1:B:33:ASN:HB2	1.72	0.72
1:B:74:VAL:HG13	1:B:100:LEU:HA	1.71	0.71
1:A:66:ALA:HB2	2:D:171:TYR:CE1	2.24	0.71
1:B:171:SER:O	1:B:172:ASP:OD1	2.09	0.70
1:A:171:SER:O	1:A:172:ASP:HB3	1.91	0.69
2:F:116:ARG:HB3	2:F:159:MET:HE1	1.75	0.69
1:B:113:THR:HG21	5:B:226:HOH:O	1.93	0.67
1:A:74:VAL:HG13	1:A:100:LEU:HA	1.78	0.65
2:F:243:THR:HG22	2:F:244:GLU:N	2.10	0.65
1:A:9:GLN:HE22	1:A:167:CYS:H	1.43	0.64
2:D:116:ARG:HB3	2:D:159:MET:HE2	1.78	0.64
1:A:16:ASP:OD2	1:A:89:LYS:CD	2.46	0.64
1:A:32:GLY:CA	1:A:33:ASN:CB	2.76	0.64
1:A:124:MET:HG2	5:A:238:HOH:O	1.99	0.62
1:B:134:PRO:O	1:B:135:MET:SD	2.58	0.61
1:B:140:ILE:O	1:B:141:SER:OG	2.14	0.61
2:F:240:ASN:OD1	2:F:240:ASN:N	2.32	0.61
2:F:238:LEU:O	2:F:240:ASN:HA	1.99	0.61
1:A:36:TYR:O	2:F:166:GLU:HG3	2.01	0.60
1:B:113:THR:HG22	5:B:211:HOH:O	2.00	0.60
1:A:16:ASP:OD2	1:A:89:LYS:HD3	2.02	0.59
2:F:244:GLU:OE1	2:F:249:ARG:N	2.35	0.59
2:F:116:ARG:HB3	2:F:159:MET:HE2	1.84	0.58
2:D:239:TYR:O	2:D:241:SER:N	2.35	0.58
1:A:34:ASP:HB3	1:A:38:ASN:O	2.03	0.58
1:A:121:ARG:NH1	1:A:180:GLY:HA2	2.19	0.57
1:A:78:VAL:O	1:A:80:GLY:N	2.39	0.55
1:A:172:ASP:CG	1:A:173:GLY:N	2.59	0.55
1:A:22:LYS:HE3	1:A:83:GLU:OE2	2.07	0.55
1:A:172:ASP:CG	1:A:173:GLY:H	2.11	0.54
1:B:34:ASP:HB3	1:B:38:ASN:O	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:213:LEU:HG	2:D:240:ASN:HD21	1.74	0.53
1:A:169:LYS:HB3	5:A:235:HOH:O	2.07	0.52
1:A:172:ASP:OD2	1:A:173:GLY:N	2.35	0.52
2:F:244:GLU:HB2	2:F:248:PHE:HA	1.91	0.52
1:A:16:ASP:OD2	1:A:89:LYS:HD2	2.10	0.52
1:B:85:LEU:HD23	1:B:103:PHE:HE2	1.75	0.52
2:D:234:LEU:HA	2:D:240:ASN:HB2	1.92	0.52
2:D:191:TYR:CE1	5:D:446:HOH:O	2.62	0.51
1:A:32:GLY:HA3	2:D:170:PHE:CE2	2.46	0.51
1:B:140:ILE:O	1:B:141:SER:CB	2.59	0.51
2:D:230:ASN:HD22	2:D:231:THR:N	2.09	0.50
1:A:80:GLY:O	1:A:82:LYS:N	2.40	0.50
2:F:210:ASN:HD22	2:F:213:LEU:H	1.59	0.50
1:B:33:ASN:CB	1:B:40:ILE:H	2.26	0.48
1:B:33:ASN:HB2	1:B:40:ILE:HG22	1.94	0.48
2:F:231:THR:OG1	5:F:426:HOH:O	2.20	0.48
1:B:66:ALA:HB2	2:F:171:TYR:CE1	2.49	0.48
2:D:156:ALA:HB3	2:D:159:MET:HE3	1.96	0.48
1:A:85:LEU:HD23	1:A:103:PHE:HE2	1.77	0.48
1:A:79:GLY:O	1:A:80:GLY:C	2.52	0.47
2:D:156:ALA:H	2:D:159:MET:HE1	1.80	0.47
2:D:156:ALA:HB3	2:D:159:MET:CE	2.45	0.47
1:A:9:GLN:NE2	1:A:167:CYS:H	2.11	0.47
1:B:31:SER:H	1:B:32:GLY:CA	2.27	0.47
1:A:22:LYS:CG	1:A:83:GLU:HG2	2.45	0.47
1:A:22:LYS:HG3	1:A:83:GLU:HG2	1.97	0.46
2:F:156:ALA:H	2:F:159:MET:HE1	1.80	0.46
2:D:210:ASN:HD22	2:D:213:LEU:H	1.63	0.46
1:A:33:ASN:O	1:A:34:ASP:HB3	2.16	0.46
1:A:169:LYS:HB2	1:A:169:LYS:HE2	1.84	0.46
1:B:180:GLY:O	1:B:181:ALA:C	2.55	0.45
1:B:132:ARG:HA	1:B:148:TRP:HB3	1.97	0.45
1:A:170:ARG:O	1:A:171:SER:C	2.54	0.45
2:F:243:THR:O	2:F:245:LEU:HD22	2.17	0.44
1:B:89:LYS:HG2	1:B:97:HIS:CE1	2.52	0.44
1:B:33:ASN:CG	1:B:40:ILE:HB	2.38	0.44
2:F:243:THR:O	2:F:244:GLU:C	2.55	0.44
2:F:243:THR:CG2	2:F:244:GLU:N	2.78	0.44
1:A:32:GLY:HA2	1:A:33:ASN:CB	2.48	0.44
1:B:22:LYS:HE2	1:B:80:GLY:HA2	2.00	0.43
1:A:34:ASP:O	1:A:36:TYR:O	2.36	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:230:ASN:HD22	2:F:231:THR:N	2.16	0.43
1:B:150:ASP:O	1:B:154:GLU:HB2	2.19	0.43
1:A:72:CYS:HB2	5:A:208:HOH:O	2.19	0.43
1:B:48:LYS:HD2	1:B:48:LYS:HA	1.72	0.43
1:B:33:ASN:CG	1:B:34:ASP:N	2.73	0.42
1:B:72:CYS:HB2	5:B:202:HOH:O	2.19	0.42
1:B:6:VAL:HG11	1:B:11:ALA:HB2	2.01	0.42
2:F:156:ALA:HB3	2:F:159:MET:CE	2.50	0.42
2:F:210:ASN:HD21	2:F:212:PHE:HB3	1.84	0.42
2:F:248:PHE:C	2:F:248:PHE:CD2	2.92	0.42
1:A:6:VAL:HG11	1:A:11:ALA:HB2	2.02	0.42
1:B:39:PRO:HB2	1:B:41:LYS:HE3	2.02	0.41
1:A:150:ASP:O	1:A:154:GLU:HB2	2.20	0.41
1:B:33:ASN:OD1	1:B:40:ILE:HB	2.21	0.41
1:B:171:SER:O	1:B:172:ASP:CB	2.69	0.41
1:B:33:ASN:HB3	1:B:40:ILE:H	1.85	0.41
1:B:31:SER:H	1:B:32:GLY:HA3	1.85	0.41
1:A:77:ASP:OD1	1:A:80:GLY:HA3	2.21	0.41
1:A:154:GLU:OE2	1:A:160:HIS:HB2	2.21	0.40
1:A:135:MET:HA	1:A:152:VAL:CG2	2.51	0.40
2:D:123:ASP:HB3	5:D:444:HOH:O	2.20	0.40
1:B:22:LYS:HE2	1:B:80:GLY:CA	2.52	0.40
1:A:72:CYS:HB3	1:A:99:THR:HB	2.03	0.40
2:F:156:ALA:HB3	2:F:159:MET:HE3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	180/194 (93%)	167 (93%)	7 (4%)	6 (3%)	5 1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	180/194 (93%)	165 (92%)	10 (6%)	5 (3%)	6	2
2	D	159/165 (96%)	145 (91%)	12 (8%)	2 (1%)	15	9
2	F	159/165 (96%)	148 (93%)	11 (7%)	0	100	100
All	All	678/718 (94%)	625 (92%)	40 (6%)	13 (2%)	10	4

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	81	LYS
1	B	81	LYS
2	D	240	ASN
1	A	79	GLY
1	A	80	GLY
1	A	172	ASP
1	B	56	PRO
1	A	120	HIS
1	B	120	HIS
1	B	172	ASP
1	B	34	ASP
2	D	244	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/168 (93%)	143 (92%)	13 (8%)	14	9
1	B	156/168 (93%)	140 (90%)	16 (10%)	9	5
2	D	136/139 (98%)	126 (93%)	10 (7%)	17	13
2	F	136/139 (98%)	120 (88%)	16 (12%)	6	3
All	All	584/614 (95%)	529 (91%)	55 (9%)	11	7

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	VAL
1	A	61	GLU
1	A	71	VAL
1	A	74	VAL
1	A	81	LYS
1	A	82	LYS
1	A	100	LEU
1	A	108	ASP
1	A	141	SER
1	A	147	LEU
1	A	152	VAL
1	A	169	LYS
1	A	172	ASP
1	B	30	ASP
1	B	31	SER
1	B	35	ILE
1	B	61	GLU
1	B	74	VAL
1	B	81	LYS
1	B	82	LYS
1	B	100	LEU
1	B	108	ASP
1	B	129	LYS
1	B	135	MET
1	B	136	ILE
1	B	138	CYS
1	B	147	LEU
1	B	152	VAL
1	B	172	ASP
2	D	110	LYS
2	D	123	ASP
2	D	146	LEU
2	D	155	GLU
2	D	190	LEU
2	D	230	ASN
2	D	242	PHE
2	D	245	LEU
2	D	248	PHE
2	D	249	ARG
2	F	110	LYS
2	F	111	THR
2	F	123	ASP
2	F	146	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	170	PHE
2	F	190	LEU
2	F	230	ASN
2	F	239	TYR
2	F	240	ASN
2	F	241	SER
2	F	242	PHE
2	F	244	GLU
2	F	245	LEU
2	F	247	GLN
2	F	248	PHE
2	F	249	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	158	ASN
2	D	210	ASN
2	D	230	ASN
2	D	240	ASN
2	D	252	GLN
2	F	210	ASN
2	F	230	ASN
2	F	252	GLN
2	F	259	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/194 (93%)	0.27	10 (5%) 29 37	17, 34, 66, 106	0
1	B	182/194 (93%)	1.26	42 (23%) 1 1	23, 49, 95, 136	0
2	D	159/165 (96%)	0.30	9 (5%) 27 35	15, 27, 74, 106	0
2	F	159/165 (96%)	0.47	12 (7%) 17 23	25, 40, 70, 111	0
All	All	682/718 (94%)	0.59	73 (10%) 8 11	15, 37, 80, 136	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	246	ALA	8.1
1	B	182	ALA	7.2
2	D	240	ASN	7.1
1	A	182	ALA	7.0
2	F	240	ASN	6.9
1	A	181	ALA	6.6
1	B	134	PRO	5.9
1	B	136	ILE	5.8
2	D	247	GLN	5.7
2	F	242	PHE	5.6
2	D	241	SER	5.4
1	B	137	PRO	5.1
1	B	157	ILE	4.8
2	D	239	TYR	4.8
1	B	152	VAL	4.5
1	B	23	ALA	4.3
1	B	181	ALA	4.2
1	B	79	GLY	4.2
1	B	39	PRO	4.1
2	D	245	LEU	4.1
1	B	151	TRP	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	135	MET	4.0
1	B	107	TRP	3.9
1	B	33	ASN	3.9
1	B	143	PRO	3.8
2	D	243	THR	3.7
1	B	62	PHE	3.7
1	B	139	TYR	3.7
1	B	149	MET	3.4
1	B	80	GLY	3.4
1	A	180	GLY	3.3
2	F	247	GLN	3.3
1	B	141	SER	3.2
2	F	241	SER	3.2
1	B	44	GLN	3.1
1	B	48	LYS	3.1
1	B	32	GLY	3.1
2	D	105	MET	3.0
1	B	155	LYS	3.0
1	B	63	ILE	2.9
1	B	47	ILE	2.8
1	B	156	ASN	2.8
1	A	171	SER	2.7
2	F	170	PHE	2.7
1	B	31	SER	2.7
1	B	40	ILE	2.7
1	B	147	LEU	2.6
1	A	80	GLY	2.6
1	B	81	LYS	2.6
1	B	133	CYS	2.6
1	B	173	GLY	2.6
1	A	173	GLY	2.5
1	A	38	ASN	2.4
2	F	112	HIS	2.4
1	B	43	ILE	2.4
1	A	41	LYS	2.4
2	F	109	ARG	2.4
1	B	60	ILE	2.3
1	B	129	LYS	2.3
1	A	155	LYS	2.3
2	F	239	TYR	2.3
2	F	207	SER	2.2
1	B	36	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	38	ASN	2.2
2	F	231	THR	2.2
1	B	132	ARG	2.2
1	B	153	THR	2.2
1	B	92	GLY	2.2
1	A	31	SER	2.1
2	D	112	HIS	2.1
2	F	111	THR	2.1
2	F	245	LEU	2.0
1	B	82	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ZN	D	301	1/1	1.00	0.13	1.59	17,17,17,17	0
4	CA	F	305	1/1	0.98	0.14	1.08	31,31,31,31	0
3	ZN	F	301	1/1	1.00	0.11	0.18	29,29,29,29	0
4	CA	D	304	1/1	0.97	0.11	-0.07	25,25,25,25	0
4	CA	D	305	1/1	0.97	0.10	-0.64	29,29,29,29	0
3	ZN	F	302	1/1	0.99	0.10	-0.88	33,33,33,33	0
3	ZN	D	302	1/1	0.99	0.11	-1.02	23,23,23,23	0
4	CA	F	303	1/1	0.98	0.08	-1.39	28,28,28,28	0
4	CA	D	303	1/1	1.00	0.09	-2.21	18,18,18,18	0
4	CA	F	304	1/1	0.97	0.07	-2.29	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.