



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 8, 2016 – 05:06 PM EDT

PDB ID : 5IMY
Title : Trapped Toxin
Authors : Lawrence, S.L.; Morton, C.J.; Parker, M.W.
Deposited on : 2016-03-07
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939

i

X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	490	<div> <div>5%</div> <div>75%</div> <div>15%</div> <div>6%</div> <div>••</div> </div>
1	B	490	<div> <div>5%</div> <div>71%</div> <div>19%</div> <div>6%</div> <div>•</div> </div>
2	C	78	<div> <div>3%</div> <div>82%</div> <div>18%</div> </div>
2	D	78	<div> <div>78%</div> <div>18%</div> <div>•</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 17217 atoms, of which 8415 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vaginolysin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	474	Total	C	H	N	O	S	0	2	0
			7356	2338	3637	652	720	9			
1	B	472	Total	C	H	N	O	S	0	0	0
			7308	2322	3614	649	714	9			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	SER	-	expression tag	UNP B2YGA4
A	28	ASN	-	expression tag	UNP B2YGA4
A	30	HIS	MET	conflict	UNP B2YGA4
A	31	MET	ALA	conflict	UNP B2YGA4
A	333	CYS	THR	conflict	UNP B2YGA4
A	348	CYS	ILE	conflict	UNP B2YGA4
B	27	SER	-	expression tag	UNP B2YGA4
B	28	ASN	-	expression tag	UNP B2YGA4
B	30	HIS	MET	conflict	UNP B2YGA4
B	31	MET	ALA	conflict	UNP B2YGA4
B	333	CYS	THR	conflict	UNP B2YGA4
B	348	CYS	ILE	conflict	UNP B2YGA4

- Molecule 2 is a protein called CD59 glycoprotein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	78	Total	C	H	N	O	S	0	0	0
			1208	388	582	106	121	11			
2	D	78	Total	C	H	N	O	S	0	0	0
			1208	388	582	106	121	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	initiating methionine	UNP P13987
D	0	MET	-	initiating methionine	UNP P13987


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	57	Total O 57 57	0	0
3	C	6	Total O 6 6	0	0
3	D	13	Total O 13 13	0	0
3	B	61	Total O 61 61	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Vaginolysin




• Molecule 1: Vaginolysin





• Molecule 2: CD59 glycoprotein





● Molecule 2: CD59 glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	81.89Å 141.71Å 106.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.64 – 2.40 43.20 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.1 (42.64-2.40) 84.3 (43.20-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 2.24Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, R_{free}	0.234 , 0.276 0.239 , 0.288	Depositor DCC
R_{free} test set	1520 reflections (3.37%)	DCC
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	17217	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3813	0.49	0/5183
1	B	0.26	0/3776	0.49	0/5133
2	C	0.29	0/638	0.44	0/863
2	D	0.28	0/638	0.43	0/863
All	All	0.26	0/8865	0.48	0/12042

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	257	ASP	Peptide
1	B	49	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3719	3637	3623	65	2
1	B	3694	3614	3615	67	2
2	C	626	582	580	5	0
2	D	626	582	580	6	0
3	A	57	0	0	5	0
3	B	61	0	0	6	0
3	C	6	0	0	1	0
3	D	13	0	0	1	0
All	All	8802	8415	8398	141	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:LYS:NZ	1:B:383:ASP:OD2	2.04	0.90
1:B:316:GLU:OE1	3:B:601:HOH:O	1.90	0.88
1:A:45:ALA:O	1:A:47:LYS:N	2.08	0.86
2:D:47:PHE:O	2:D:51:THR:OG1	1.96	0.83
1:B:178:LYS:NZ	3:B:602:HOH:O	2.11	0.82

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:ARG:HH22	1:B:353:ASP:OD2[3_646]	1.46	0.14
1:A:162:ARG:NH2	1:B:353:ASP:OD2[3_646]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/490 (97%)	404 (85%)	43 (9%)	27 (6%)	2	1
1	B	470/490 (96%)	396 (84%)	47 (10%)	27 (6%)	2	1
2	C	76/78 (97%)	75 (99%)	1 (1%)	0	100	100
2	D	76/78 (97%)	72 (95%)	4 (5%)	0	100	100
All	All	1096/1136 (96%)	947 (86%)	95 (9%)	54 (5%)	3	1

5 of 54 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	SER
1	A	46	ALA
1	A	76	LYS
1	A	87	GLU
1	A	264	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/416 (98%)	366 (90%)	40 (10%)	10	14
1	B	402/416 (97%)	363 (90%)	39 (10%)	10	15
2	C	72/73 (99%)	65 (90%)	7 (10%)	10	15
2	D	72/73 (99%)	63 (88%)	9 (12%)	6	7
All	All	952/978 (97%)	857 (90%)	95 (10%)	10	14

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	33	LEU
2	D	69	CYS
1	B	464	ARG
2	C	66	LYS
2	D	28	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/490 (96%)	0.17	25 (5%) 30 30	30, 52, 92, 128	0
1	B	472/490 (96%)	0.25	24 (5%) 32 32	33, 56, 91, 139	0
2	C	78/78 (100%)	0.09	2 (2%) 59 58	37, 53, 68, 102	0
2	D	78/78 (100%)	0.11	0 100 100	38, 57, 73, 95	0
All	All	1102/1136 (97%)	0.19	51 (4%) 36 37	30, 54, 89, 139	0

The worst 5 of 51 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	341	ALA	9.2
1	A	43	SER	8.6
1	B	338	GLY	7.8
1	B	340	SER	6.2
1	A	339	GLY	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.