



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:32 AM GMT

PDB ID : 3ISM
Title : Crystal structure of the EndoG/EndoGI complex: Mechanism of EndoG inhibition
Authors : Loll, B.; Gebhardt, M.; Wahle, E.; Meinhart, A.
Deposited on : 2009-08-26
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 3% 87% 8%
1	B	267	 6% 86% 6% 7%
2	C	359	 18% 72% 8% 20%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6757 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CG8862.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	1998	1255	366	369	8	0	3	0
1	B	247	2051	1294	366	383	8	0	13	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	55	MET	-	INITIATING METHIONINE	UNP Q7JXB9
A	311	ALA	-	EXPRESSION TAG	UNP Q7JXB9
A	312	ALA	-	EXPRESSION TAG	UNP Q7JXB9
A	313	ALA	-	EXPRESSION TAG	UNP Q7JXB9
A	314	LEU	-	EXPRESSION TAG	UNP Q7JXB9
A	315	GLU	-	EXPRESSION TAG	UNP Q7JXB9
A	316	HIS	-	EXPRESSION TAG	UNP Q7JXB9
A	317	HIS	-	EXPRESSION TAG	UNP Q7JXB9
A	318	HIS	-	EXPRESSION TAG	UNP Q7JXB9
A	319	HIS	-	EXPRESSION TAG	UNP Q7JXB9
A	320	HIS	-	EXPRESSION TAG	UNP Q7JXB9
A	321	HIS	-	EXPRESSION TAG	UNP Q7JXB9
B	55	MET	-	INITIATING METHIONINE	UNP Q7JXB9
B	311	ALA	-	EXPRESSION TAG	UNP Q7JXB9
B	312	ALA	-	EXPRESSION TAG	UNP Q7JXB9
B	313	ALA	-	EXPRESSION TAG	UNP Q7JXB9
B	314	LEU	-	EXPRESSION TAG	UNP Q7JXB9
B	315	GLU	-	EXPRESSION TAG	UNP Q7JXB9
B	316	HIS	-	EXPRESSION TAG	UNP Q7JXB9
B	317	HIS	-	EXPRESSION TAG	UNP Q7JXB9
B	318	HIS	-	EXPRESSION TAG	UNP Q7JXB9
B	319	HIS	-	EXPRESSION TAG	UNP Q7JXB9
B	320	HIS	-	EXPRESSION TAG	UNP Q7JXB9
B	321	HIS	-	EXPRESSION TAG	UNP Q7JXB9

- Molecule 2 is a protein called CG4930.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	286	2370	1515	384	462	9	0	7	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	2	ALA	SER	ENGINEERED	UNP Q9V3V9

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	B	1	1	1	0	0
3	A	1	1	1	0	0

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	8	4	1	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	148	Total 148	O 148	0	0
5	B	114	Total 114	O 114	0	0
5	C	66	Total 66	O 66	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.79Å 109.15Å 121.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.23 – 2.20 43.22 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.23-2.20) 99.8 (43.22-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.184 , 0.223 0.187 , 0.224	Depositor DCC
R_{free} test set	2412 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	26.8	Xtrriage
Anisotropy	0.204	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.5	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 48241 reflections	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6757	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/2060	0.62	0/2790
1	B	0.49	0/2143	0.56	0/2905
2	C	0.48	0/2446	0.54	0/3314
All	All	0.52	0/6649	0.58	0/9009

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1998	0	1930	10	0
1	B	2051	0	2007	13	0
2	C	2370	0	2287	18	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	8	0	12	2	0
5	A	148	0	0	2	0
5	B	114	0	0	4	0
5	C	66	0	0	0	0
All	All	6757	0	6236	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:206[B]:SER:OG	5:B:365:HOH:O	1.73	1.05
1:B:206[A]:SER:HB2	5:B:365:HOH:O	1.93	0.69
2:C:170:SER:O	2:C:174:ARG:HG2	2.01	0.60
1:B:88[A]:ARG:NH2	5:B:417:HOH:O	2.36	0.59
1:B:280[A]:VAL:HG12	1:B:284:SER:OG	2.04	0.57
2:C:325[B]:CYS:SG	2:C:339:PRO:HD3	2.45	0.57
1:A:100[A]:ARG:HH22	1:B:100:ARG:HH22	1.53	0.56
2:C:219:LEU:HD22	2:C:275:GLU:HG2	1.88	0.55
1:B:69[B]:THR:HG22	1:B:72:ARG:H	1.72	0.54
2:C:67:ARG:HD2	2:C:110:TRP:CG	2.42	0.54
1:A:130:LYS:NZ	5:A:425:HOH:O	2.42	0.52
2:C:314:ASP:O	2:C:318:LYS:HE2	2.11	0.51
1:A:100[A]:ARG:NH2	5:A:406:HOH:O	2.45	0.50
4:A:6989:TRS:H12	2:C:146[A]:LEU:CD2	2.41	0.50
1:A:100[A]:ARG:NH2	1:B:100:ARG:NH2	2.59	0.50
1:B:104:PRO:HG2	1:B:156:MET:HG3	1.96	0.48
2:C:242:GLN:O	2:C:246:LYS:HG2	2.15	0.47
2:C:220:GLU:HA	2:C:223:ARG:HB2	1.96	0.47
2:C:72:ARG:O	2:C:76:ASP:HB2	2.14	0.47
1:A:287[B]:ARG:HH22	1:B:236:ASN:HA	1.79	0.47
2:C:128:ARG:HG2	2:C:163:CYS:SG	2.54	0.46
1:A:167:LYS:NZ	1:A:171:GLU:OE2	2.48	0.46
1:B:206[A]:SER:OG	5:B:399:HOH:O	2.17	0.46
2:C:115:ARG:HG3	2:C:127:TRP:CZ3	2.52	0.45
1:A:184:GLN:OE1	2:C:142[A]:ARG:HD3	2.17	0.44
1:B:108:PHE:CE1	1:B:211:CYS:HB2	2.53	0.43
4:A:6989:TRS:H12	2:C:146[A]:LEU:HD23	1.99	0.43
1:A:131:GLN:HG2	2:C:52[A]:ASP:OD2	2.19	0.43
2:C:67:ARG:HD2	2:C:110:TRP:CD2	2.53	0.43
2:C:341:GLU:H	2:C:341:GLU:CD	2.22	0.42
1:A:88:ARG:NH1	1:A:134:SER:OG	2.52	0.42
2:C:303:VAL:HG13	2:C:308:LEU:HB2	2.03	0.41
1:B:182:VAL:HG21	1:B:267:GLN:HA	2.02	0.41
2:C:327:ARG:CZ	2:C:352:PHE:HB3	2.51	0.40
1:A:108:PHE:CE2	1:A:211:CYS:HB2	2.56	0.40
1:B:66:LEU:HA	1:B:66:LEU:HD12	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/267 (92%)	239 (97%)	6 (2%)	1 (0%)	39	42
1	B	258/267 (97%)	250 (97%)	7 (3%)	1 (0%)	39	42
2	C	287/359 (80%)	279 (97%)	8 (3%)	0	100	100
All	All	791/893 (89%)	768 (97%)	21 (3%)	2 (0%)	46	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	222	ASP
1	B	186	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/235 (93%)	215 (98%)	4 (2%)	66	79
1	B	231/235 (98%)	229 (99%)	2 (1%)	84	92
2	C	257/306 (84%)	257 (100%)	0	100	100
All	All	707/776 (91%)	701 (99%)	6 (1%)	84	93

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	108	PHE
1	A	252	SER
1	A	254	ASP
1	B	108	PHE
1	B	130	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TRS	A	6989	-	7,7,7	0.94	1 (14%)	9,9,9	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TRS	A	6989	-	-	0/9/9/9	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	6989	TRS	C-N	-2.44	1.47	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	6989	TRS	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/267 (91%)	0.25	7 (2%) 55 54	25, 29, 40, 46	0
1	B	247/267 (92%)	0.35	15 (6%) 25 24	24, 30, 41, 48	0
2	C	286/359 (79%)	0.90	64 (22%) 1 1	25, 31, 43, 53	0
All	All	778/893 (87%)	0.52	86 (11%) 7 7	24, 30, 41, 53	0

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	265	PRO	9.7
2	C	21	ALA	7.0
2	C	340	PHE	6.4
2	C	228	ARG	6.0
2	C	238	ASP	5.3
2	C	229	PHE	5.0
2	C	222	TYR	4.9
2	C	270	GLN	4.9
2	C	274	ALA	4.8
2	C	239	SER	4.8
2	C	273	ALA	4.7
2	C	235	ILE	4.5
1	B	65	SER	4.2
1	A	67	THR	4.1
2	C	302	GLU	4.0
1	B	255	HIS	4.0
2	C	220	GLU	3.9
2	C	262	ASP	3.9
1	B	69[A]	THR	3.9
1	A	311	ALA	3.9
2	C	266	SER	3.9
1	A	310	VAL	3.8
2	C	271	ASN	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	23	ASN	3.7
2	C	227	LYS	3.7
2	C	254	GLU	3.7
2	C	245	VAL	3.6
2	C	341	GLU	3.6
2	C	225	LEU	3.6
2	C	267	ALA	3.5
2	C	263	GLY	3.5
1	B	66	LEU	3.4
2	C	256	ILE	3.4
2	C	261	PHE	3.4
2	C	312	TRP	3.3
2	C	22	GLN	3.3
2	C	230	VAL	3.2
2	C	243	ALA	3.2
2	C	264	THR	3.2
1	B	70	PRO	3.1
2	C	258	LYS	3.1
2	C	226	MET	3.0
2	C	310	ASP	2.9
2	C	219	LEU	2.9
2	C	240	VAL	2.8
2	C	344	GLU	2.8
2	C	224	THR	2.8
2	C	247	LYS	2.7
2	C	249	ALA	2.7
2	C	250	ALA	2.7
2	C	352	PHE	2.7
2	C	315	LYS	2.7
2	C	257	TRP	2.7
2	C	251	ALA	2.6
2	C	297	PHE	2.6
2	C	174	ARG	2.6
2	C	308	LEU	2.6
2	C	223	ARG	2.5
1	B	216	TYR	2.5
1	B	68	ALA	2.5
1	B	239	ALA	2.5
1	B	88[A]	ARG	2.5
1	B	240	VAL	2.4
2	C	347	GLY	2.4
1	B	214	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	67	THR	2.4
2	C	272	LYS	2.4
2	C	298	ASN	2.4
2	C	317	VAL	2.3
2	C	242	GLN	2.3
1	B	176	SER	2.3
1	B	103	VAL	2.2
1	A	120[A]	ASP	2.2
1	A	100[A]	ARG	2.1
1	A	103	VAL	2.1
2	C	255	ILE	2.1
2	C	241	HIS	2.1
2	C	269	ASP	2.1
2	C	46	VAL	2.1
2	C	309	LEU	2.1
2	C	275	GLU	2.1
1	A	228	VAL	2.1
2	C	337	PRO	2.1
2	C	334	ASN	2.0
1	B	97	TYR	2.0
2	C	305	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	TRS	A	6989	8/8	0.87	0.14	-0.17	41,44,45,45	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MG	A	1	1/1	0.92	0.04	-3.11	18,18,18,18	0
3	MG	B	1	1/1	0.90	0.05	-4.34	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.