wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 04:33 PM BST

PDB ID	:	1IT1
Title	:	Solution structures of ferrocytochrome c3 from Desulfovibrio vulgaris Miyazaki F
Authors	:	Harada, E.; Fukuoka, Y.; Ohmura, T.; Fukunishi, A.; Kawai, G.; Fujiwara, T.; Akutsu, H.
Deposited on	:	2001-12-29

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 59%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	107	

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 13 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:2-A:36, A:41-A:87, A:92-A:107 (98)	0.15	13

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 5 single-model clusters were found.

Cluster number	Models
1	5, 6, 12, 14, 17
2	3, 13, 18, 20
3	1, 8, 9, 19
4	11, 15
Single-model clusters	2; 4; 7; 10; 16

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1881 atoms, of which 908 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called cytochrome c3.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	107	1581	489	780	150	151	11	0

- Molecule 2 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).



Mol	Chain	Residues	Atoms					
			Total	C	Fe	H	N	O
2	A	1	75	34	1	32	4	4
			Total	C	Fe	H	N	O
2	A	1	75	34	1	32	4	4
			Total	C	Fe	H	N	O
2	A	1	75	34	1	32	4	4
			Total	C	Fe	H	N	O
2	A	1	75	34	1	32	4	4

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: cytochrome c3

Chain A: 13% 62% 16% • 8%



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 13. Colouring as in section 4.1 above.

- Molecule 1: cytochrome c3

Chain A: 11% 58% 21% • 8%



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 300 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.8.5.1
X-PLOR	refinement	3.8.5.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5333
Number of chemical shift lists	5
Total number of shifts	897
Number of shifts mapped to atoms	817
Number of unparsed shifts	0
Number of shifts with mapping errors	80
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	59%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.96±0.00	0±0/763 (0.0±0.0%)	1.30±0.01	1±1/1014 (0.0±0.0%)
All	All	0.96	0/15260 (0.0%)	1.30	10/20280 (0.0%)

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	51	CYS	N-CA-CB	-5.45	100.80	110.60	11	10

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	746	725	725	199±11
2	A	172	128	120	83±8
All	All	18360	17060	16909	4269

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 121.

5 of 686 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:A:201:HEC:HMD1	2:A:201:HEC:CGD	1.02	1.84	7	3
1:A:47:ALA:HB2	1:A:65:TYR:CE1	1.00	1.91	3	17
2:A:201:HEC:HMD1	2:A:201:HEC:O2D	0.98	1.59	2	1
2:A:201:HEC:HMD3	2:A:203:HEC:HBB3	0.97	1.37	13	19
2:A:203:HEC:HBC3	2:A:203:HEC:HMC1	0.96	1.33	8	9

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/107 (91%)	69±3 (71±3%)	25±3 (26±3%)	3±1 (3±1%)	8	40
All	All	1940/2140 (91%)	1375 (71%)	503 (26%)	62 (3%)	8	40

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	35	HIS	20
1	A	48	THR	20
1	A	67	HIS	9
1	A	49	ALA	3
1	A	6	ALA	3

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/85 (95%)	57±3 (70±3%)	24±3 (30±3%)	2	18
All	All	1620/1700 (95%)	1139 (70%)	481 (30%)	2	18

5 of 44 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	100	CYS	20
1	A	9	LEU	20
1	A	84	LEU	20
1	A	105	CYS	20
1	A	24	THR	20

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	HEC	A	201	1	24,50,50	1.86±0.02	2±0 (8±0%)
2	HEC	A	202	1	24,50,50	1.96±0.06	2±0 (6±2%)
2	HEC	A	203	1	24,50,50	1.88±0.04	2±0 (8±0%)
2	HEC	A	204	1	24,50,50	1.79±0.05	2±0 (8±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics

could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	HEC	A	201	1	19,82,82	2.32±0.05	2±0 (10±0%)
2	HEC	A	202	1	19,82,82	2.32±0.10	2±0 (10±1%)
2	HEC	A	203	1	19,82,82	2.47±0.09	2±0 (10±0%)
2	HEC	A	204	1	19,82,82	2.44±0.07	2±0 (10±1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	201	1	-	0±0,6,54,54	0±0,0,8,8
2	HEC	A	202	1	-	0±0,6,54,54	0±0,0,8,8
2	HEC	A	203	1	-	0±0,6,54,54	0±0,0,8,8
2	HEC	A	204	1	-	0±0,6,54,54	0±0,0,8,8

5 of 8 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	202	HEC	C3B-C2B	7.58	1.32	1.40	16	20
2	A	203	HEC	C3B-C2B	7.32	1.33	1.40	16	20
2	A	204	HEC	C3C-C2C	6.21	1.34	1.40	18	19
2	A	201	HEC	C3C-C2C	6.15	1.34	1.40	16	20
2	A	201	HEC	C3B-C2B	5.98	1.34	1.40	14	20

5 of 8 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
2	A	203	HEC	CBC-CAC-C3C	8.21	109.39	127.34	16	20
2	A	204	HEC	CBB-CAB-C3B	7.60	110.72	127.34	3	20
2	A	202	HEC	CBB-CAB-C3B	7.47	111.00	127.34	4	20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
2	A	202	HEC	CBC-CAC-C3C	7.12	111.78	127.34	3	19
2	A	203	HEC	CBB-CAB-C3B	6.96	112.11	127.34	16	20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 59% for the well-defined parts and 59% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 5333

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	817
Number of shifts mapped to atoms	817
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	50

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	0	—	—
$^{13}C_{\beta}$	0	—	—
$^{13}C'$	0	—	—
^{15}N	102	-0.19 ± 0.42	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 59%, i.e. 681 atoms were assigned a chemical shift out of a possible 1155. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	286/482 (59%)	192/192 (100%)	0/196 (0%)	94/94 (100%)
Sidechain	339/568 (60%)	335/341 (98%)	0/204 (0%)	4/23 (17%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	56/105 (53%)	48/58 (83%)	0/38 (0%)	8/9 (89%)
Overall	681/1155 (59%)	575/591 (97%)	0/438 (0%)	106/126 (84%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	20	PHE	HE1	0.80	8.69 – 5.49	-19.7
1	A	5	PRO	HD2	-1.90	5.45 – 1.85	-15.4
1	A	70	HIS	HD2	0.19	9.28 – 4.78	-15.2
1	A	70	HIS	HE1	0.57	10.53 – 5.43	-14.5
1	A	34	HIS	HD2	0.56	9.28 – 4.78	-14.4
1	A	35	HIS	HE1	0.79	10.53 – 5.43	-14.1
1	A	22	HIS	HD2	0.73	9.28 – 4.78	-14.0
1	A	52	HIS	HD2	0.81	9.28 – 4.78	-13.8
1	A	106	HIS	HD2	0.86	9.28 – 4.78	-13.7
1	A	83	HIS	HD2	0.90	9.28 – 4.78	-13.6
1	A	35	HIS	HD2	0.91	9.28 – 4.78	-13.6
1	A	52	HIS	HE1	1.12	10.53 – 5.43	-13.5
1	A	34	HIS	HE1	1.14	10.53 – 5.43	-13.4
1	A	25	HIS	HD2	1.04	9.28 – 4.78	-13.3
1	A	83	HIS	HE1	1.32	10.53 – 5.43	-13.1
1	A	22	HIS	HE1	1.43	10.53 – 5.43	-12.8
1	A	25	HIS	HE1	1.65	10.53 – 5.43	-12.4
1	A	106	HIS	HE1	2.07	10.53 – 5.43	-11.6
1	A	41	GLU	HG3	-0.10	3.31 – 1.21	-11.2
1	A	16	GLN	HE22	2.10	9.27 – 4.77	-10.9
1	A	20	PHE	HE2	3.84	8.69 – 5.49	-10.2
1	A	66	TYR	HE1	4.77	7.86 – 5.56	-8.4
1	A	66	TYR	HE2	4.77	7.86 – 5.56	-8.4
1	A	35	HIS	HB3	-0.16	5.00 – 1.10	-8.2
1	A	5	PRO	HG2	-0.60	3.48 – 0.38	-8.2
1	A	35	HIS	HB2	0.21	4.91 – 1.31	-8.1
1	A	41	GLU	HG2	0.60	3.33 – 1.23	-8.0
1	A	68	ALA	HB1	-0.56	2.61 – 0.11	-7.7
1	A	68	ALA	HB3	-0.56	2.61 – 0.11	-7.7
1	A	68	ALA	HB2	-0.56	2.61 – 0.11	-7.7
1	A	70	HIS	HB2	0.39	4.91 – 1.31	-7.6
1	A	20	PHE	HD2	4.89	8.56 – 5.56	-7.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	22	HIS	HA	1.46	6.81 – 2.41	-7.2
1	A	69	MET	HB2	-0.26	3.73 – 0.33	-6.7
1	A	34	HIS	HB2	0.72	4.91 – 1.31	-6.6
1	A	25	HIS	HB2	0.77	4.91 – 1.31	-6.5
1	A	4	ALA	HB1	-0.17	2.61 – 0.11	-6.1
1	A	4	ALA	HB2	-0.17	2.61 – 0.11	-6.1
1	A	4	ALA	HB3	-0.17	2.61 – 0.11	-6.1
1	A	20	PHE	HZ	4.55	9.11 – 4.91	-5.9
1	A	24	THR	HG22	-0.18	2.29 – -0.01	-5.7
1	A	24	THR	HG21	-0.18	2.29 – -0.01	-5.7
1	A	24	THR	HG23	-0.18	2.29 – -0.01	-5.7
1	A	16	GLN	HB3	0.50	3.37 – 0.67	-5.6
1	A	35	HIS	H	4.45	11.68 – 4.78	-5.5
1	A	34	HIS	HA	2.22	6.81 – 2.41	-5.4
1	A	22	HIS	HB2	1.19	4.91 – 1.31	-5.3
1	A	22	HIS	HB3	1.03	5.00 – 1.10	-5.2
1	A	16	GLN	HE21	4.86	9.53 – 4.93	-5.2
1	A	34	HIS	HB3	1.05	5.00 – 1.10	-5.1

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: BMRB entry 5333

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	20
Number of shifts mapped to atoms	0
Number of unparsed shifts	0
Number of shifts with mapping errors	20
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. First 5 (of 20) occurences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	1	HEM	HBB	2.03	0.02	1
UNMAPPED	1	HEM	HMA	3.3	0.02	1
UNMAPPED	1	HEM	HBAA	3.27	0.02	2
UNMAPPED	1	HEM	HADA	4.54	0.02	2
UNMAPPED	1	HEM	HHB	8.98	0.02	1

7.2.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.2.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 1155. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	0/482 (0%)	0/192 (0%)	0/196 (0%)	0/94 (0%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Sidechain	0/568 (0%)	0/341 (0%)	0/204 (0%)	0/23 (0%)
Aromatic	0/105 (0%)	0/58 (0%)	0/38 (0%)	0/9 (0%)
Overall	0/1155 (0%)	0/591 (0%)	0/438 (0%)	0/126 (0%)

7.2.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots (i)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_2). RCI is only applicable to proteins.

7.3 Chemical shift list 3

File name: BMRB entry 5333

Chemical shift list name: *assigned_chem_shift_list_3*

7.3.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	20
Number of shifts mapped to atoms	0
Number of unparsed shifts	0
Number of shifts with mapping errors	20
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. First 5 (of 20) occurences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	1	HEM	HBB	0.38	0.02	1
UNMAPPED	1	HEM	HMA	3.15	0.02	1
UNMAPPED	1	HEM	HBAA	2.71	0.02	2

Continued on next page...

Continued from previous page...

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	1	HEM	HADA	3.81	0.02	2
UNMAPPED	1	HEM	HHB	9.6	0.02	1

7.3.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.3.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 1155. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	0/482 (0%)	0/192 (0%)	0/196 (0%)	0/94 (0%)
Sidechain	0/568 (0%)	0/341 (0%)	0/204 (0%)	0/23 (0%)
Aromatic	0/105 (0%)	0/58 (0%)	0/38 (0%)	0/9 (0%)
Overall	0/1155 (0%)	0/591 (0%)	0/438 (0%)	0/126 (0%)

7.3.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.3.5 Random Coil Index (RCI) plots (i)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_3). RCI is only applicable to proteins.

7.4 Chemical shift list 4

File name: BMRB entry 5333

Chemical shift list name: *assigned_chem_shift_list_4*

7.4.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	20
Number of shifts mapped to atoms	0
Number of unparsed shifts	0
Number of shifts with mapping errors	20
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. First 5 (of 20) occurences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	1	HEM	HBB	3.0	0.02	1
UNMAPPED	1	HEM	HMA	3.65	0.02	1
UNMAPPED	1	HEM	HBAA	3.15	0.02	2
UNMAPPED	1	HEM	HADA	3.95	0.02	2
UNMAPPED	1	HEM	HHB	10.15	0.02	1

7.4.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.4.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 1155. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	0/482 (0%)	0/192 (0%)	0/196 (0%)	0/94 (0%)
Sidechain	0/568 (0%)	0/341 (0%)	0/204 (0%)	0/23 (0%)
Aromatic	0/105 (0%)	0/58 (0%)	0/38 (0%)	0/9 (0%)
Overall	0/1155 (0%)	0/591 (0%)	0/438 (0%)	0/126 (0%)

7.4.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.4.5 Random Coil Index (RCI) plots (i)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_4). RCI is only applicable to proteins.

7.5 Chemical shift list 5

File name: BMRB entry 5333

Chemical shift list name: *assigned_chem_shift_list_5*

7.5.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	20
Number of shifts mapped to atoms	0
Number of unparsed shifts	0
Number of shifts with mapping errors	20
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. First 5 (of 20) occurences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	1	HEM	HBB	1.92	0.02	1
UNMAPPED	1	HEM	HMA	3.28	0.02	1
UNMAPPED	1	HEM	HBAA	2.91	0.02	2
UNMAPPED	1	HEM	HADA	4.13	0.02	2
UNMAPPED	1	HEM	HHB	9.37	0.02	1

7.5.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.5.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical

shift out of a possible 1155. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	0/482 (0%)	0/192 (0%)	0/196 (0%)	0/94 (0%)
Sidechain	0/568 (0%)	0/341 (0%)	0/204 (0%)	0/23 (0%)
Aromatic	0/105 (0%)	0/58 (0%)	0/38 (0%)	0/9 (0%)
Overall	0/1155 (0%)	0/591 (0%)	0/438 (0%)	0/126 (0%)

7.5.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.5.5 Random Coil Index (RCI) plots (i)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_5). RCI is only applicable to proteins.