



wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:40 PM BST

PDB ID : 3IYQ
EMDB ID: : EMD-5188
Title : tmRNA-SmpB: a journey to the center of the bacterial ribosome
Authors : Weis, F.; Bron, P.; Giudice, E.; Rolland, J.P.; Thomas, D.; Felden, B.; Gillet, R.
Deposited on : 2010-04-16
Resolution : 13.00 Å(reported)
Based on PDB ID : 2OB7, 2CZJ

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/EMValidationReportHelp>

MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk27241


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 13.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
Ramachandran outliers	111179	726
Sidechain outliers	111093	686
RNA backbone	3027	244

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	349	
2	B	149	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called tmRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	349	Total	C	N	O	P	0	0
			7466	3324	1373	2420	349		

- Molecule 2 is a protein called SsrA-binding protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	B	122	Total	C	N	O	0	0
			992	637	181	174		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	145	HIS	-	EXPRESSION TAG	UNP Q8RR57
B	146	HIS	-	EXPRESSION TAG	UNP Q8RR57
B	147	HIS	-	EXPRESSION TAG	UNP Q8RR57
B	148	HIS	-	EXPRESSION TAG	UNP Q8RR57
B	149	HIS	-	EXPRESSION TAG	UNP Q8RR57
B	150	HIS	-	EXPRESSION TAG	UNP Q8RR57

4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of images	49061	Depositor
Resolution determination method	FSC at 0.5 cut-off	Depositor
CTF correction method	Each particle	Depositor
Microscope	JEM2200FS	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	Not provided	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	50000	Depositor
Image detector	Kodak SO163 film	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 2$	RMSZ	$\# Z > 2$
1	A	1.24	1/8346 (0.0%)	1.71	268/13017 (2.1%)
2	B	0.76	0/1010	1.30	7/1354 (0.5%)
All	All	1.20	1/9356 (0.0%)	1.68	275/14371 (1.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	86
2	B	0	4
All	All	0	90

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	219	C	C5'-C4'	5.30	1.57	1.51

The worst 5 of 275 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	310	G	P-O3'-C3'	17.13	140.25	119.70
1	A	239	G	P-O3'-C3'	16.54	139.55	119.70
1	A	309	G	P-O3'-C3'	15.51	138.31	119.70
1	A	157	C	P-O3'-C3'	15.22	137.96	119.70
1	A	193	G	P-O3'-C3'	15.12	137.84	119.70

There are no chirality outliers.

5 of 90 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	2	G	Sidechain
1	A	3	G	Sidechain
1	A	4	G	Sidechain
1	A	7	G	Sidechain
1	A	8	A	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7466	0	3800	4	0
2	B	992	0	1030	2	0
All	All	8458	0	4830	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:82:HIS:CE1	2:B:84:HIS:CD2	2.81	0.67
1:A:18:G:C8	2:B:80:LEU:HB3	2.49	0.48
1:A:97:A:H2'	1:A:101:A:C2	2.51	0.46
1:A:145:C:H4'	1:A:146:U:H3'	2.02	0.41
1:A:208:C:H2'	1:A:209:C:C6	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	120/149 (80%)	96 (80%)	16 (13%)	8 (7%)	1	24

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	49	GLU
2	B	62	PRO
2	B	75	ARG
2	B	10	ALA
2	B	82	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	101/125 (81%)	96 (95%)	5 (5%)	30	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	48	PHE
2	B	62	PRO
2	B	83	LYS
2	B	86	LEU
2	B	121	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	84	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	348/349 (99%)	239 (68%)	47 (13%)

5 of 239 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	4	G
1	A	5	G
1	A	6	U
1	A	7	G
1	A	8	A

5 of 47 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	184	U
1	A	219	C
1	A	331	A
1	A	193	G
1	A	223	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.