



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 09:24 PM BST

PDB ID : 2J11  
Title : p53 tetramerization domain mutant Y327S T329G Q331G  
Authors : Carbajo, R.J.; Mora, P.; Sanchez del Pino, M.M.; Perez-Paya, E.; Pineda-Lucena, A.  
Deposited on : 2006-08-08

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 45%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	31	
1	B	31	
1	C	31	
1	D	31	

## 2 Ensemble composition and analysis

This entry contains 29 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 17 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:327-A:354, B:327-B:354, C:327-C:354, D:327-D:354 (112)	0.48	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 5, 6, 8, 9, 11, 12, 13, 16, 19, 20, 21, 22, 23, 29
2	18, 25, 28
3	7, 17, 26
4	3, 4, 27
5	15, 24
Single-model clusters	10; 14

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1980 atoms, of which 976 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CELLULAR TUMOR ANTIGEN P53.

Mol	Chain	Residues	Atoms						Trace
1	A	31	Total	C	H	N	O	S	0
			495	155	244	46	49	1	
1	B	31	Total	C	H	N	O	S	0
			495	155	244	46	49	1	
1	C	31	Total	C	H	N	O	S	0
			495	155	244	46	49	1	
1	D	31	Total	C	H	N	O	S	0
			495	155	244	46	49	1	

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	327	SER	TYR	ENGINEERED MUTATION	UNP P04637
A	329	GLY	THR	ENGINEERED MUTATION	UNP P04637
A	331	GLY	GLN	ENGINEERED MUTATION	UNP P04637
B	327	SER	TYR	ENGINEERED MUTATION	UNP P04637
B	329	GLY	THR	ENGINEERED MUTATION	UNP P04637
B	331	GLY	GLN	ENGINEERED MUTATION	UNP P04637
C	327	SER	TYR	ENGINEERED MUTATION	UNP P04637
C	329	GLY	THR	ENGINEERED MUTATION	UNP P04637
C	331	GLY	GLN	ENGINEERED MUTATION	UNP P04637
D	327	SER	TYR	ENGINEERED MUTATION	UNP P04637
D	329	GLY	THR	ENGINEERED MUTATION	UNP P04637
D	331	GLY	GLN	ENGINEERED MUTATION	UNP P04637

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: CELLULAR TUMOR ANTIGEN P53




- Molecule 1: CELLULAR TUMOR ANTIGEN P53



- Molecule 1: CELLULAR TUMOR ANTIGEN P53



- Molecule 1: CELLULAR TUMOR ANTIGEN P53



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 20. Colouring as in section 4.1 above.

- Molecule 1: CELLULAR TUMOR ANTIGEN P53






- Molecule 1: CELLULAR TUMOR ANTIGEN P53



- Molecule 1: CELLULAR TUMOR ANTIGEN P53



- Molecule 1: CELLULAR TUMOR ANTIGEN P53



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *CNS*.

Of the 30 calculated structures, 29 were deposited, based on the following criterion: *TOTAL ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
SPARKY	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 7252
Number of chemical shift lists	1
Total number of shifts	924
Number of shifts mapped to atoms	816
Number of unparsed shifts	0
Number of shifts with mapping errors	108
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	45%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	232	228	228	10±3
1	B	232	228	228	10±3
1	C	232	228	228	9±4
1	D	232	228	228	10±4
All	All	26912	26448	26448	740

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

5 of 348 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:344:LEU:HD22	1:C:344:LEU:HD11	1.04	1.30	12	3
1:A:344:LEU:HD21	1:D:344:LEU:HD22	1.00	1.31	20	1
1:B:330:LEU:HD22	1:B:332:ILE:HD11	0.95	1.37	20	5
1:B:344:LEU:HD11	1:C:344:LEU:HD22	0.93	1.35	6	2
1:C:330:LEU:HD23	1:C:332:ILE:HD11	0.92	1.40	29	4



## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	28/31 (90%)	26±1 (93±4%)	2±1 (6±4%)	0±0 (1±1%)	34	78
1	B	28/31 (90%)	26±1 (94±3%)	1±1 (4±3%)	0±0 (1±2%)	18	63
1	C	28/31 (90%)	26±1 (94±3%)	1±1 (5±3%)	0±0 (1±2%)	19	64
1	D	28/31 (90%)	26±1 (94±4%)	1±1 (5±4%)	0±0 (1±2%)	18	63
All	All	3248/3596 (90%)	3044 (94%)	164 (5%)	40 (1%)	21	68

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	D	327	SER	12
1	B	327	SER	12
1	C	327	SER	11
1	A	327	SER	5

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	23/24 (96%)	19±2 (81±8%)	4±2 (19±8%)	5	38
1	B	23/24 (96%)	19±2 (82±8%)	4±2 (18±8%)	6	40
1	C	23/24 (96%)	19±2 (83±9%)	4±2 (17±9%)	6	42
1	D	23/24 (96%)	18±2 (80±8%)	5±2 (20±8%)	5	36
All	All	2668/2784 (96%)	2176 (82%)	492 (18%)	5	39

5 of 75 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	344	LEU	15
1	D	333	ARG	15
1	D	344	LEU	14
1	C	330	LEU	14
1	D	330	LEU	14

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation ⓘ

The completeness of assignment taking into account all chemical shift lists is 45% for the well-defined parts and 45% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 7252

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping ⓘ

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	924
Number of shifts mapped to atoms	816
Number of unparsed shifts	0
Number of shifts with mapping errors	108
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Residue not found in structure. First 5 (of 108) occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
D	6	LYS	HD2	1.27	0.01	2
D	2	TYR	HD2	6.78	0.01	1
B	4	PHE	HE1	7.21	0.01	1
D	4	PHE	HZ	7.13	0.01	1
B	4	PHE	HD1	7.02	0.01	1

#### 7.1.2 Chemical shift referencing ⓘ

No chemical shift referencing corrections were calculated (not enough data).

### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 45%, i.e. 672 atoms were assigned a chemical shift out of a possible 1508. 0 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	200/560 (36%)	200/224 (89%)	0/224 (0%)	0/112 (0%)
Sidechain	412/840 (49%)	412/496 (83%)	0/284 (0%)	0/60 (0%)
Aromatic	60/108 (56%)	60/60 (100%)	0/48 (0%)	0/0 (—%)
Overall	672/1508 (45%)	672/780 (86%)	0/556 (0%)	0/172 (0%)


### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.


### 7.1.5 Random Coil Index (RCI) plots [i](#)

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.


Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:



Random coil index (RCI) for chain C:



Random coil index (RCI) for chain D:

