wwPDB X-ray Structure Validation Summary Report (i)

Sep 12, 2016 – 03:27 PM EDT

PDB ID	:	5J4A
Title	:	CdiA-CT toxin from Burkholderia pseudomallei E479 in complex with cognate CdiI immunity protein
Authors	:	Goulding, C.W.; Johnson, P.M.
Deposited on	:	2016-03-31
Resolution	:	2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	.% 67% 5% 28%
1	C	161	.% 64% 7% • 28%
2	B	120	82% 5% 13%
2	D	120	73% 14% 13%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3537 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA nuclease CdiA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	116	873	554	151	166	2	0	0	0
1	C	116	873	554	151	166	2	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	156	MET	-	initiating methionine	UNP H9T8G6
A	285	ALA	ASP	engineered mutation	UNP H9T8G6
C	156	MET	-	initiating methionine	UNP H9T8G6
C	285	ALA	ASP	engineered mutation	UNP H9T8G6

- Molecule 2 is a protein called Immunity protein CdiI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	104	818	516	137	161	4	0	0	0
2	D	104	818	516	137	161	4	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	LYS	-	expression tag	UNP H9T8G7
B	110	THR	-	expression tag	UNP H9T8G7
B	111	SER	-	expression tag	UNP H9T8G7
B	112	LEU	-	expression tag	UNP H9T8G7
B	113	GLU	-	expression tag	UNP H9T8G7
B	114	HIS	-	expression tag	UNP H9T8G7
B	115	HIS	-	expression tag	UNP H9T8G7
B	116	HIS	-	expression tag	UNP H9T8G7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	117	HIS	-	expression tag	UNP H9T8G7
B	118	HIS	-	expression tag	UNP H9T8G7
B	119	HIS	-	expression tag	UNP H9T8G7
D	0	LYS	-	expression tag	UNP H9T8G7
D	110	THR	-	expression tag	UNP H9T8G7
D	111	SER	-	expression tag	UNP H9T8G7
D	112	LEU	-	expression tag	UNP H9T8G7
D	113	GLU	-	expression tag	UNP H9T8G7
D	114	HIS	-	expression tag	UNP H9T8G7
D	115	HIS	-	expression tag	UNP H9T8G7
D	116	HIS	-	expression tag	UNP H9T8G7
D	117	HIS	-	expression tag	UNP H9T8G7
D	118	HIS	-	expression tag	UNP H9T8G7
D	119	HIS	-	expression tag	UNP H9T8G7

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	40	40	40	0	0
			Total	O		
3	B	51	51	51	0	0
			Total	O		
3	C	35	35	35	0	0
			Total	O		
3	D	29	29	29	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: tRNA nuclease CdiA

Chain A:



- Molecule 1: tRNA nuclease CdiA

Chain C:

- Molecule 2: Immunity protein CdiI

Chain B:



- Molecule 2: Immunity protein CdiI

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α, β, γ	54.47Å 73.26Å 110.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.71 – 2.00 43.71 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (43.71-2.00) 95.6 (43.71-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	5.79 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.193 , 0.237 0.197 , 0.241	Depositor DCC
R_{free} test set	1932 reflections (6.67%)	DCC
Wilson B-factor (Å^2)	35.5	Xtriage
Anisotropy	0.756	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 39.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	3537	wwPDB-VP
Average B, all atoms (Å^2)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.48% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.38	0/888	0.53	0/1202
1	C	0.35	0/888	0.55	0/1202
2	B	0.41	0/833	0.54	0/1125
2	D	0.38	0/833	0.51	0/1125
All	All	0.38	0/3442	0.53	0/4654

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	873	0	879	5	0
1	C	873	0	879	11	0
2	B	818	0	799	4	0
2	D	818	0	799	11	0
3	A	40	0	0	0	0
3	B	51	0	0	1	0
3	C	35	0	0	3	0
3	D	29	0	0	0	0
All	All	3537	0	3356	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:ASP:OD2	2:B:2:ALA:HB2	1.84	0.77
1:C:229:ASP:OD2	2:D:2:ALA:HB2	1.86	0.76
1:C:246:LEU:HD13	1:C:284:LEU:HB3	1.71	0.73
1:C:246:LEU:HD11	1:C:268:PHE:CZ	2.30	0.66
1:A:289:LEU:O	1:A:294:LYS:NZ	2.30	0.65

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/161 (71%)	113 (99%)	1 (1%)	0	100	100
1	C	114/161 (71%)	112 (98%)	2 (2%)	0	100	100
2	B	102/120 (85%)	101 (99%)	1 (1%)	0	100	100
2	D	102/120 (85%)	101 (99%)	1 (1%)	0	100	100
All	All	432/562 (77%)	427 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/128 (72%)	91 (99%)	1 (1%)	80	83
1	C	92/128 (72%)	90 (98%)	2 (2%)	60	62
2	B	91/106 (86%)	91 (100%)	0	100	100
2	D	91/106 (86%)	91 (100%)	0	100	100
All	All	366/468 (78%)	363 (99%)	3 (1%)	86	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	273	SER
1	C	304	LEU
1	C	306	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	220	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	116/161 (72%)	-0.17	1 (0%)	85	86	26, 34, 47, 57	0
1	C	116/161 (72%)	0.14	1 (0%)	85	86	28, 41, 54, 64	0
2	B	104/120 (86%)	-0.09	0	100	100	24, 31, 42, 55	0
2	D	104/120 (86%)	-0.02	0	100	100	26, 36, 47, 58	0
All	All	440/562 (78%)	-0.03	2 (0%)	91	92	24, 36, 50, 64	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	201	ARG	2.4
1	C	287	ARG	2.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.