



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:53 AM GMT

PDB ID : 2J7W
Title : DENGUE VIRUS NS5 RNA DEPENDENT RNA POLYMERASE DOMAIN
COMPLEXED WITH 3'DGTP
Authors : Yap, T.L.; Xu, T.; Chen, Y.L.; Malet, H.; Egloff, M.P.; Canard, B.; Vasudevan,
S.G.; Lescar, J.
Deposited on : 2006-10-17
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	635	<div> <div>9%</div> <div>75%</div> <div>14%</div> <div>10%</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POLYPROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	571	Total	C	N	O	S	0	0	0
			4621	2926	828	837	30			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	374	GLU	GLY	CONFLICT	UNP Q6DLV0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			13	10	3		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	129	Total	O	0	0
			129	129		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: POLYPROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	163.72Å 180.92Å 58.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60 19.96 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.60) 99.7 (19.96-2.60)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 2.59Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.213 , 0.289 0.207 , 0.280	Depositor DCC
R_{free} test set	1340 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	62.5	Xtriage
Anisotropy	0.267	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 60.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 26889 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4772	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, ZN, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	2/4739 (0.0%)	0.58	1/6423 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	309	GLU	C-O	17.70	1.56	1.23
1	A	279	ARG	CZ-NH1	6.73	1.41	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	279	ARG	NE-CZ-NH2	-7.17	116.71	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4621	0	4493	52	0
2	A	2	0	0	0	0
3	A	13	0	0	0	0
4	A	7	0	10	0	0
5	A	129	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4772	0	4503	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:676:ASN:OD1	1:A:696:PRO:HB3	1.79	0.83
1:A:422:LYS:HA	1:A:422:LYS:HE2	1.62	0.81
1:A:802:GLN:H	1:A:802:GLN:HE21	1.27	0.80
1:A:704:GLN:HE21	1:A:715:GLU:H	1.34	0.76
1:A:346:THR:HB	1:A:347:PRO:HD3	1.69	0.72
1:A:718:MET:HG2	1:A:837:PRO:HG3	1.75	0.68
1:A:333:VAL:HA	5:A:2010:HOH:O	1.93	0.68
1:A:716:LEU:HD21	1:A:839:LEU:HD23	1.76	0.67
1:A:305:HIS:HB2	5:A:2061:HOH:O	1.94	0.66
1:A:308:TYR:O	1:A:309:GLU:HB2	1.94	0.66
1:A:863:ILE:HG22	1:A:864:PRO:HD3	1.80	0.63
1:A:792:ARG:HD3	1:A:795:TRP:CZ2	2.38	0.59
1:A:604:GLY:HA2	5:A:2034:HOH:O	2.04	0.58
1:A:809:MET:HA	1:A:809:MET:HE2	1.86	0.57
1:A:446:CYS:HB2	1:A:570:LEU:HD13	1.86	0.56
1:A:433:LEU:HD23	1:A:436:ARG:HH21	1.70	0.56
1:A:548:ASN:HD22	1:A:551:LYS:HE2	1.71	0.54
1:A:549:GLU:HB3	1:A:608:LEU:HD22	1.87	0.54
1:A:717:ILE:HD13	1:A:723:LYS:HG2	1.90	0.54
1:A:420:SER:HB3	5:A:2025:HOH:O	2.08	0.54
1:A:731:GLN:HG2	1:A:735:ILE:HD12	1.91	0.53
1:A:704:GLN:NE2	1:A:715:GLU:H	2.03	0.52
1:A:650:LYS:O	1:A:654:ARG:HG2	2.11	0.51
1:A:507:GLU:HG3	1:A:606:TYR:CE1	2.45	0.51
1:A:283:ILE:HG21	1:A:591:ILE:HD11	1.94	0.49
1:A:841:LYS:HG2	1:A:851:ILE:HD13	1.93	0.49
1:A:722:ARG:HD3	1:A:826:ASP:HB3	1.95	0.49
1:A:484:GLU:OE2	1:A:602:GLN:HB3	2.13	0.49
1:A:550:GLU:OE2	1:A:569:LYS:HE3	2.14	0.48
1:A:364:ARG:HG2	1:A:365:PRO:HD2	1.95	0.48
1:A:790:THR:O	1:A:791:SER:O	2.32	0.47
1:A:845:GLN:HG3	1:A:851:ILE:HD11	1.96	0.47
1:A:507:GLU:HG3	1:A:606:TYR:HE1	1.79	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:714:HIS:CG	1:A:839:LEU:HD21	2.50	0.46
1:A:716:LEU:O	1:A:723:LYS:HA	2.16	0.46
1:A:395:ARG:HD2	1:A:431:TRP:CD2	2.50	0.46
1:A:371:LYS:HE3	1:A:638:LEU:O	2.15	0.46
1:A:694:TRP:HB2	5:A:2084:HOH:O	2.15	0.45
1:A:497:PHE:HA	1:A:502:SER:HB3	1.98	0.45
1:A:333:VAL:HB	5:A:2011:HOH:O	2.16	0.45
1:A:583:THR:HA	1:A:584:PRO:HD3	1.87	0.45
1:A:481:ARG:HE	1:A:481:ARG:HA	1.81	0.44
1:A:290:THR:HB	1:A:309:GLU:H	1.80	0.44
1:A:565:ASN:O	1:A:569:LYS:HB2	2.17	0.44
1:A:841:LYS:HG2	1:A:851:ILE:CD1	2.48	0.43
1:A:298:PRO:HB2	1:A:355:LYS:HE3	2.00	0.43
1:A:537:TRP:CG	1:A:663:ASP:HB3	2.53	0.43
1:A:630:LYS:O	1:A:634:GLU:HG3	2.20	0.42
1:A:610:THR:HG23	1:A:660:ILE:HG22	2.03	0.41
1:A:481:ARG:HA	1:A:481:ARG:NE	2.34	0.41
1:A:334:VAL:HA	1:A:335:PRO:HD3	1.90	0.41
1:A:366:LEU:O	1:A:370:ARG:HG3	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	563/635 (89%)	520 (92%)	37 (7%)	6 (1%)	17 36

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	276	ILE
1	A	791	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	355	LYS
1	A	356	GLU
1	A	346	THR
1	A	405	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	488/551 (89%)	467 (96%)	21 (4%)	35 64

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	345	THR
1	A	389	ARG
1	A	425	VAL
1	A	440	LEU
1	A	474	TRP
1	A	476	MET
1	A	481	ARG
1	A	507	GLU
1	A	569	LYS
1	A	581	ARG
1	A	590	ASP
1	A	664	ASP
1	A	693	GLN
1	A	785	VAL
1	A	802	GLN
1	A	808	ASP
1	A	841	LYS
1	A	851	ILE
1	A	853	LEU
1	A	854	THR
1	A	863	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	339	GLN
1	A	350	GLN
1	A	548	ASN
1	A	555	GLN
1	A	562	GLN
1	A	645	GLN
1	A	682	ASN
1	A	693	GLN
1	A	704	GLN
1	A	742	GLN
1	A	760	GLN
1	A	768	HIS
1	A	802	GLN
1	A	835	ASN
1	A	861	GLN
1	A	869	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GTP	A	1886	-	8,12,34	0.94	1 (12%)	15,20,54	1.66	3 (20%)
4	PEG	A	1887	-	6,6,6	0.49	0	5,5,5	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTP	A	1886	-	-	0/12/12/38	0/0/0/3
4	PEG	A	1887	-	-	0/4/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1886	GTP	PG-O3G	2.06	1.62	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1886	GTP	PB-O3A-PA	-3.97	119.35	132.67
3	A	1886	GTP	PB-O3B-PG	-2.26	125.10	132.67
3	A	1886	GTP	O2G-PG-O1G	2.62	119.02	110.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	571/635 (89%)	0.44	59 (10%) 9 5	50, 60, 80, 84	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	346	THR	8.0
1	A	353	VAL	7.5
1	A	347	PRO	7.4
1	A	345	THR	7.3
1	A	349	GLY	7.1
1	A	289	SER	7.1
1	A	586	GLY	6.8
1	A	354	PHE	6.3
1	A	350	GLN	5.7
1	A	317	SER	5.4
1	A	587	THR	5.2
1	A	293	TYR	5.2
1	A	348	PHE	5.1
1	A	295	ASP	5.0
1	A	592	ILE	4.6
1	A	352	ARG	4.4
1	A	443	LEU	4.1
1	A	600	SER	4.1
1	A	601	GLY	4.0
1	A	285	GLU	3.6
1	A	303	ALA	3.6
1	A	590	ASP	3.5
1	A	743	GLY	3.4
1	A	344	ASP	3.3
1	A	423	ALA	3.3
1	A	291	TRP	3.2
1	A	588	VAL	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	318	SER	3.2
1	A	744	ALA	3.1
1	A	288	ASN	3.1
1	A	294	ASP	2.8
1	A	308	TYR	2.8
1	A	591	ILE	2.7
1	A	358	VAL	2.7
1	A	582	PRO	2.7
1	A	580	GLN	2.6
1	A	306	GLY	2.6
1	A	298	PRO	2.5
1	A	276	ILE	2.4
1	A	881	ASP	2.4
1	A	583	THR	2.4
1	A	420	SER	2.3
1	A	297	ASN	2.3
1	A	351	GLN	2.3
1	A	585	THR	2.3
1	A	742	GLN	2.3
1	A	589	MET	2.3
1	A	880	LEU	2.3
1	A	795	TRP	2.3
1	A	305	HIS	2.2
1	A	765	MET	2.2
1	A	360	THR	2.2
1	A	361	ARG	2.1
1	A	286	GLU	2.1
1	A	474	TRP	2.1
1	A	304	TYR	2.1
1	A	324	VAL	2.1
1	A	487	ALA	2.0
1	A	357	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GTP	A	1886	13/32	0.82	0.28	0.87	117,118,118,119	0
2	ZN	A	1884	1/1	0.99	0.03	-2.95	45,45,45,45	0
2	ZN	A	1885	1/1	0.97	0.06	-3.58	63,63,63,63	1
4	PEG	A	1887	7/7	0.88	0.22	-	77,77,79,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.